

# **ΠΑΝΕΠΙΣΤΗΜΙΟ ΘΕΣΣΑΛΙΑΣ**

ΣΧΟΛΗ ΕΠΙΣΤΗΜΩΝ ΥΓΕΙΑΣ  
ΤΜΗΜΑ ΚΤΗΝΙΑΤΡΙΚΗΣ

## **ΠΕΙΡΑΜΑΤΙΚΗ ΜΕΛΕΤΗ ΤΗΣ ΤΟΞΙΝΑΙΜΙΑΣ ΤΗΣ ΕΓΚΥΜΟΣΥΝΗΣ ΣΕ ΠΡΟΒΑΤΙΝΕΣ ΚΑΙ ΤΗΣ ΣΥΣΧΕΤΙΣΗΣ ΑΥΤΗΣ ΜΕ ΜΑΣΤΙΤΙΔΑ ΚΑΤΑ ΤΗΝ ΕΠΙΛΟΧΕΙΑ ΠΕΡΙΟΔΟ**

**ΜΑΡΙΑΝΑ Σ. ΜΠΑΡΜΠΑΓΙΑΝΝΗ**

*Κτηνίατρος*

### **ΔΙΔΑΚΤΟΡΙΚΗ ΔΙΑΤΡΙΒΗ**

που εκπονήθηκε στην Κλινική Μαιευτικής και Αναπαραγωγής  
του Τμήματος Κτηνιατρικής του Πανεπιστημίου Θεσσαλίας

Καρδίτσα 2016

# ΠΕΡΙΛΗΨΗ

Η παρούσα διατριβή εστιάζεται: (α) στην ανάπτυξη ενός έγκυρου προτύπου για τη μελέτη της τοξιναιμίας της εγκυμοσύνης, με ταυτόχρονη διατήρηση των απαιτήσεων κορεσμού των ζώων, (β) στην καταγραφή, με υπερηχογραφική εξέταση, των μεταβολών στο έμβρυο και στο μαστό των προβατίνων με τοξιναιμία της εγκυμοσύνης, (γ) στην αξιολόγηση των συνεπειών της τοξιναιμίας της εγκυμοσύνης στις προσβεβλημένες προβατίνες κατά την περιτοκεταία περίοδο και (δ) στην αξιολόγηση του ενδεχόμενου ρόλου της τοξιναιμίας της εγκυμοσύνης σε προδιάθεση των ζώων σε μαστίτιδα κατά την αμέσως μετά τον τοκετό περίοδο.

Η διατριβή χωρίζεται σε δύο κεφάλαια και ακολουθεί η Γενική Συζήτηση.

Στο Κεφάλαιο I ανασκοπείται η βιβλιογραφία σχετικά με την τοξιναιμία της εγκυμοσύνης.

Στο Κεφάλαιο II παρουσιάζεται η πραγματοποιηθείσα ερευνητική μελέτη. Το κεφάλαιο υποδιαιρείται σε δύο τμήματα.

Το τμήμα Α του κεφαλαίου περιγράφει την ερευνητική μελέτη και τα σχετικά αποτελέσματα μέχρι τον ενοφθαλμισμό του μαστού των ζώων, ο οποίος πραγματοποιήθηκε την 5η ημέρα μετά τον τοκετό. Στη μελέτη περιλήφθησαν 28 προβατίνες. Αρχικά, στα ζώα πραγματοποιήθηκε ανθελμινθική αγωγή και, στη συνέχεια, αυτά έλαβαν ένα μίγμα από μολύνουσες προνύμφες τριχοστρογγυλοειδών ελμίνθων. Μέχρι την 60ή ημέρα της εγκυμοσύνης, σε κάθε προβατίνα χορηγείτο καθημερινά 1,30 kg έτοιμης συμπυκνωμένης ζωοτροφής (περιεκτικότητα σε ενέργεια: 0,844 FU<sub>L</sub>) και 2,50 kg σανού τριφυλλιού. Από την 60ή μέχρι την 100ή ημέρα της εγκυμοσύνης, κάθε προβατίνα ελάμβανε καθημερινά 0,60 kg συμπυκνωμένης ζωοτροφής και 2,00 σανού τριφυλλιού. Μετά την 100ή ημέρα της εγκυμοσύνης, κάθε προβατίνα που κuoφορούσε ένα έμβρυο ελάμβανε καθημερινά 0,50 kg έτοιμης συμπυκνωμένης ζωοτροφής με μειωμένη περιεκτικότητα σε ενέργεια (0,748 FU<sub>L</sub>) και 0,50 kg σανού τριφυλλιού. Οι αντίστοιχες ποσότητες για προβατίνες που κuoφορούσαν δύο έμβρυα ήταν 0,60 kg και 0,50 kg, αντίστοιχα, και για προβατίνες που κuoφορούσαν τρία έμβρυα ήταν 0,80 kg και 0,50 kg, αντίστοιχα. Στις προβατίνες πραγματοποιούνταν οι παρακάτω εξετάσεις από την 120ή ημέρα της εγκυμοσύνης μέχρι τον τοκετό: παρασιτολογική εξέταση κοπράνων, βιοχημική εξέταση αίματος για μέτρηση της συγκέντρωσης β-υδροξυβουτυρικού οξέος και γλυκόζης, υπερηχογραφική εξέταση B-mode και Doppler εμβρύου/ων και μαστού. Από τον τοκετό μέχρι την 5η ημέρα μετά από αυτόν, πραγματοποιούνταν οι παρακάτω εξετάσεις: ζύγισμα των νεογέννητων αρνιών, εξέταση του γεννητικού συστήματος και μέτρηση της παραγόμενης ποσότητας γάλακτος. Σε 16 προβατίνες παρατηρήθηκαν αυξημένες συγκεντρώσεις β-υδροξυβουτυρικού οξέος στο αίμα ( $\geq 1,2 \text{ mmol L}^{-1}$  σε

τουλάχιστον δύο μετρήσεις) και αυτά τα ζώα κατανεμήθηκαν στην ομάδα A, ενώ στις υπόλοιπες 12 προβατίνες δεν παρατηρήθηκαν αυξημένες συγκεντρώσεις β-υδροξυβουτυρικού οξέος στο αίμα και κατανεμήθηκαν στην ομάδα B. Οι προβατίνες στην ομάδα A είχαν αυξημένο αριθμό αυγών παρασίτων (epg) στα κόπρανα και μικρότερες συγκεντρώσεις γλυκόζης στο αίμα από αυτές στην ομάδα B ( $P<0,025$ ,  $P=0,033$ , αντίστοιχα). Δεν υπήρχαν διαφορές μεταξύ των δύο ομάδων (A, B) στα ευρήματα της υπερηχογραφικής εξέτασης των εμβρύων, εκτός από τον όγκο αίματος στην ομφαλική αρτηρία, παράμετρο για την οποία οι τιμές στα ζώα της ομάδας B ήταν σημαντικά μεγαλύτερες από τις αντίστοιχες τιμές στα ζώα της ομάδας A. Μολαταύτα, όταν πραγματοποιήθηκε ανακατανομή των ζώων σε προβατίνες οι οποίες εκδήλωσαν δυστοκία και σε προβατίνες οι οποίες δεν εκδήλωσαν δυστοκία, παρατηρήθηκαν σημαντικές διαφορές ( $P<0,047$ ) μεταξύ των δύο κατηγοριών σε όλες τις αιμοδυναμικές παραμέτρους στην ομφαλική αρτηρία την 140ή και την 145η ημέρα της εγκυμοσύνης. Οι τιμές της έντασης των τόνων φωτεινότητας του γκρι χρώματος στο μαστικό παρέγχυμα ήταν μεγαλύτερη στα ζώα της ομάδας A από τα ζώα της ομάδας B ( $P=0,007$ ). Ο όγκος αίματος στον μαστικό αδένα ήταν μικρότερος στα ζώα της ομάδας A απ' ό,τι στα ζώα της ομάδας B ( $P<0,05$ ) και επιπλέον παρατηρήθηκαν διαφορές μεταξύ των δύο ομάδων στο δείκτη παλμικότητας και στη μέση ταχύτητα του αίματος ( $P=0,007$ ,  $P=0,036$ , αντίστοιχα) κατά τη διάρκεια των δύο τελευταίων εβδομάδων της εγκυμοσύνης. Η μέση διάρκεια της εγκυμοσύνης ήταν σημαντικά μικρότερης διάρκειας στις προβατίνες της ομάδας A (145,75 ημέρες) απ' ό,τι στις προβατίνες της ομάδας B (148,42 ημέρες) ( $P<0,001$ ). Παρατηρήθηκε επίσης μεγαλύτερο ποσοστό προσβολής από περιτοκεταία προβλήματα στα ζώα της ομάδας A. Ειδικότερα, το ποσοστό προσβολής από δυστοκία, περιγεννητικό θάνατο των εμβρύων/νεογέννητων ή επιλόχειες παθολογικές καταστάσεις του γεννητικού συστήματος ήταν 0,500, 0,375 ή 0,250, αντίστοιχα, στα ζώα της ομάδας A, ενώ στα ζώα της ομάδας B ήταν 0,083, 0,083 ή 0,000, αντίστοιχα ( $P=0,01$ ,  $P=0,039$  ή  $P=0,031$ , αντίστοιχα). Το μέσο σωματικό βάρος των νεογέννητων αρνιών από προβατίνες στην ομάδα A ήταν 3,5 kg και αυτό των αρνιών από προβατίνες στην ομάδα B ήταν 4,0 kg ( $P=0,016$ ). Βρέθηκε ότι υπήρχε σημαντική αρνητική συσχέτιση μεταξύ των συγκεντρώσεων του β-υδροξυβουτυρικού οξέος στο αίμα των έγκυων προβατίνων και του σωματικού βάρους των νεογέννητων αρνιών, καθώς και μεταξύ του παρασιτικού φορτίου των έγκυων προβατίνων και του σωματικού βάρους των νεογέννητων αρνιών ( $P=0,016$  και  $P=0,03$ , αντίστοιχα).

Το τμήμα B του κεφαλαίου περιγράφει την ερευνητική μελέτη και τα σχετικά αποτελέσματα μετά τον ενοφθαλμισμό του μαστού των ζώων, ο οποίος πραγματοποιήθηκε την 5η ημέρα μετά τον τοκετό. Οι προβατίνες στην ομάδα A χωρίστηκαν στις υποομάδες A1 (n=8) και A2 (n=8) και αυτές στην ομάδα B στις υποομάδες B1 (n=8) και B2 (n=4). Τα ζώα στις υποομάδες A1 και B1 ενοφθαλμίστηκαν την 5η ημέρα μετά τον τοκετό με εναπόθεση *Mannheimia haemolytica* στο

θηλαίο πόρο, ενώ αυτά στις υποομάδες A2 και B2 ήταν μάρτυρες. Πραγματοποιήθηκαν κλινική και υπερηχογραφική εξέταση B-mode και Doppler του μαστού και βακτηριολογική και κυτταρολογική εξέταση του μαστικού εκκρίματος, καθώς και ιστολογική εξέταση της ενοφθαλμισμένης θηλής (μετά από θηλεκτομή) και του παρεγχύματος των δύο μαστικών αδένων (μετά από λήψη ιστοτεμαχίου με βιοψία). Μαστίπδα εκδηλώθηκε σε όλες (8/8) τις προβατίνες της υποομάδας A1, σε 1/8 προβατίνα της υποομάδας A2, σε 4/8 προβατίνες της υποομάδας B1 και σε 0/4 προβατίνες της υποομάδας B2. Η συχνότητα απομόνωσης *M. haemolytica* από ζώα των υποομάδων A1 ή A2 ήταν μεγαλύτερη από την αντίστοιχη από ζώα των υποομάδων B1 ή B2 (A1 έναντι B1:  $P < 0,08$ , A2 έναντι B2:  $P > 0,3$ ). Επίσης, *M. haemolytica* απομονώθηκε συχνότερα από δείγματα ιστών ζώων των υποομάδων A1 (A1 έναντι B1:  $P = 0,008$ ) και A2 (A2 έναντι B2:  $P = 0,058$ ). Μετά τον ενοφθαλμισμό, ο θηλαίος κόλπος και ο γαλακτοφόρος κόλπος απεικονίζονταν ως ανηχογενείς κοιλότητες, με παρουσία υπερηχογενών σωματιδίων εντός αυτών. Η υπερηχογραφική εικόνα εντός του παρεγχύματος προοδευτικά παρουσίαζε ανομοιογένεια και αδρούς σχηματισμούς. Η διάμετρος της μαστικής αρτηρίας αυξήθηκε σημαντικά ήδη έξι ώρες μετά τον ενοφθαλμισμό, οπότε παρατηρήθηκε επίσης απότομη και σημαντική αύξηση στον όγκο αίματος στη μαστική αρτηρία στην ενοφθαλμισμένη πλευρά του μαστού (A1 έναντι B1:  $P < 0,06$ ), καθώς και στη μέση ταχύτητα του αίματος στο αγγείο. Επίσης, παρατηρήθηκε αύξηση στον όγκο αίματος στις ενοφθαλμισμένες θηλές. Οι χαρακτηριστικοί λεμφοειδείς σχηματισμοί στο όριο μεταξύ θηλαίου πόρου και θηλαίου κόλπου παρατηρήθηκαν σε 3/8 προβατίνες στην υποομάδα A1 και σε 7/8 στην υποομάδα B1 ( $P = 0,019$ ). Στην υποομάδα A1, οι σωρευτικές τιμές για τα μακροσκοπικά και ιστολογικά ευρήματα στη θηλή ήταν 18 και 23, αντίστοιχα και η σωρευτική τιμή για τα ιστολογικά ευρήματα στο μαστικό παρέγχυμα της ενοφθαλμισμένης πλευράς ήταν 24. Οι αντίστοιχες τιμές για την υποομάδα A2 ήταν 2, 9 και 5, για την υποομάδα B1 ήταν 5, 31 και 16 ( $P \leq 0.05$  έναντι της υποομάδας A1) και για την υποομάδα B2 όλες οι τιμές ήταν 0 ( $P > 0.05$  έναντι της υποομάδας A2).

Τα συμπεράσματα που προκύπτουν από τα ευρήματα αυτής της διατριβής είναι τα παρακάτω.

(α) Αναπτύχθηκε πρότυπο για πρόκληση τοξιναιμίας της εγκυμοσύνης σε προβατίνες για εφαρμογή σε πειραματικές μελέτες. Με το εν λόγω πρότυπο, καλύπτονταν οι απαιτήσεις κορεσμού των ζώων. Ο παρασιτισμός των ζώων ενδεχομένως συνέβαλε στην αποτελεσματική ανάπτυξη του προτύπου.

(β) Η υπερηχογραφική εξέταση των εμβρύων των προβατίνων με τοξιναιμία της εγκυμοσύνης έδειξε μειωμένο όγκο αίματος στην ομφαλική αρτηρία. Παρατηρήθηκαν σημαντικές μεταβολές στις αιμοδυναμικές ιδιότητες, ειδικά δε σε προβατίνες στις οποίες εκδηλώθηκε δυστοκία.

(γ) Περιγράφηκε η υπερηχογραφική εικόνα των μαστικών αδένων των προβατίνων κατά τη διάρκεια της γαλακτογένεσης. Τα ευρήματα της υπερηχογραφικής εξέτασης B-mode έδειξαν διαφορές μεταξύ υγιών προβατίνων και προβατίνων με τοξιναιμία της εγκυμοσύνης στην ανάπτυξη του μαστικού παρεγχύματος. Επιπλέον, παρατηρήθηκε μικρότερος όγκος αίματος στο μαστό προβατίνων με τοξιναιμία της εγκυμοσύνης.

(δ) Παρατηρήθηκε αυξημένο ποσοστό προσβολής από περιτοκεταϊά προβλήματα σε προβατίνες με τοξιναιμία της εγκυμοσύνης, καθώς και αυξημένη συχνότητα περιστατικών περιγεννητικής θνησιμότητας στα έμβρυα/νεογνά αυτών των προβατίνων.

(ε) Η τοξιναιμία της εγκυμοσύνης αποτελεί παράγοντα προδιάθεσης για μαστίτιδα στην επιλόχεια περίοδο. Πιθανόν, η μειωμένη λειτουργικότητα των λεμφοθλακίων στο όριο θηλαίου πόρου-θηλαίου κόλπου να οδηγεί σε μειωμένη προστασία του μαστικού αδένου.

**Με βάση την κείμενη νομοθεσία και μετά από σχετική απόφαση στη με αριθμό 27/03.11.2015 συνεδρίαση της Γενικής Συνέλευσης Ειδικής Σύγκλησης του Τμήματος Κτηνιατρικής του Πανεπιστημίου Θεσσαλίας, η συγγραφή της διατριβής έγινε στην αγγλική γλώσσα.**

### **Δημοσιεύσεις σχετιζόμενες με την παρούσα διατριβή**

Στις παρακάτω επιστημονικές δημοσιεύσεις παρουσιάζονται τμήματα της παρούσας διατριβής:

I. M.S. Barbagianni, E. Giannenas, E. Papadopoulos, I.G. Petridis, S.A. Spanos, P.G. Gouletsou, I. Valasi, G.C. Fthenakis (2015) "Pregnancy toxemia in ewes: development of an experimental model and potential interactions with gastrointestinal nematode infections" *Small Ruminant Research*, 133:102-107.

II. M.S. Barbagianni, P.G. Gouletsou, I. Valasi, I.G. Petridis, I. Giannenas, G.C. Fthenakis (2015). "Ultrasonographic findings in the ovine udder during lactogenesis in healthy ewes or ewes with pregnancy toxemia" *Journal of Dairy Research*, 82:293-303.

III. M.S. Barbagianni, V.S. Mavrogianni, A.I. Katsafadou, S.A. Spanos, V. Tsioli, A.D. Galatos, M. Nakou, I. Valasi, P.G. Gouletsou, G.C. Fthenakis (2015) "Pregnancy toxemia as risk factor for development of mastitis in sheep during the immediately post-partum period" *Small Ruminant Research*, 130:246-251.

**IV.** M.S. Barbagianni, S.A. Spanos, K.S. Ioannidi, N.G.C. Vasileiou, A.I. Katsafadou, I. Valasi, P.G. Gouletsou, G.C. Fthenakis (2015) "Increased incidence of peri-parturient problems in ewes with pregnancy toxemia" *Small Ruminant Research*, 132:111-114.

### **ΤΡΙΜΕΛΗΣ ΣΥΜΒΟΥΛΕΥΤΙΚΗ ΕΠΙΤΡΟΠΗ**

Γ.Χ. Φθενάκης, Καθηγητής	Επιβλέπων
Κλινική Μαιευτικής και Αναπαραγωγής, Τμήμα Κτηνιατρικής, Πανεπιστήμιο Θεσσαλίας	
Ε.Γ. Βαλάση, Επίκουρη καθηγήτρια	Μέλος Συμβουλευτικής Επιτροπής
Εργαστήριο Φυσιολογίας, Τμήμα Κτηνιατρικής, Πανεπιστήμιο Θεσσαλίας	
Π.Γ. Γκουλέτσου, Επίκουρη καθηγήτρια	Μέλος Συμβουλευτικής Επιτροπής
Κλινική Μαιευτικής και Αναπαραγωγής, Τμήμα Κτηνιατρικής, Πανεπιστήμιο Θεσσαλίας	

### **ΕΠΤΑΜΕΛΗΣ ΕΞΕΤΑΣΤΙΚΗ ΕΠΙΤΡΟΠΗ**

Γ.Σ. Αμοιρίδης, Καθηγητής	Τμήμα Κτηνιατρικής, Πανεπιστήμιο Θεσσαλίας
Γ.Χ. Φθενάκης, Καθηγητής	Τμήμα Κτηνιατρικής, Πανεπιστήμιο Θεσσαλίας
M. Caroprese, Αναπλ. καθηγήτρια	Dipartimento di Scienze Agrarie, degli Alimenti e dell'Ambiente, Università degli Studi di Foggia
Ε. Βαλάση, Επίκουρη καθηγήτρια	Τμήμα Κτηνιατρικής, Πανεπιστήμιο Θεσσαλίας
Π.Γ. Γκουλέτσου, Επίκουρη καθηγήτρια	Τμήμα Κτηνιατρικής, Πανεπιστήμιο Θεσσαλίας
Β.Σ. Μαυρογιάννη, Επίκουρη καθηγήτρια	Τμήμα Κτηνιατρικής, Πανεπιστήμιο Θεσσαλίας
Η. Γιάννενας, Λέκτορας	Τμήμα Κτηνιατρικής, Αριστοτέλειο Πανεπιστήμιο Θεσσαλονίκης

# **UNIVERSITY OF THESSALY**

SCHOOL OF HEALTH SCIENCES

FACULTY OF VETERINARY MEDICINE

## **EXPERIMENTAL STUDY OF PREGNANCY TOXAEMIA IN EWES AND ITS ASSOCIATION WITH MASTITIS IN THE POST-PARTUM PERIOD**

**MARIANA S. BARBAGIANNI**

*DVM (Thessaly)*

### **A THESIS SUBMITTED FOR THE DEGREE OF DOCTOR OF PHILOSOPHY**

Work carried out at the Department of Obstetrics and Reproduction  
of the Faculty of Veterinary Medicine of the University of Thessaly

Karditsa, Greece 2016



# ABSTRACT

Specific objectives of the present thesis were as follows: (i) the development of a valid model for studying pregnancy toxæmia, whilst maintaining appropriate satiation requirements in the experimental animals, (ii) the recording, by means of ultrasonographic examination, of changes occurring in the foetus and in the udder of ewes with pregnancy toxæmia, (iii) the evaluation of consequences of pregnancy toxæmia in affected ewes around the lambing period and (iv) the evaluation of the potential predisposing role of pregnancy toxæmia in development of mastitis in the immediately post-partum period.

The thesis is divided into two chapters followed by the General Discussion.

In Chapter I, the literature regarding pregnancy toxæmia is reviewed.

In the Chapter II, the research work performed is described in detail. The chapter is subdivided into two sections.

Section A of the chapter describes work performed and relevant results until challenge of the mammary gland of the experimental animals, which took place on the 5th day after lambing. In total, 28 ewes were included into the study. These initially had received effective anthelmintic treatment and, then, received a mixture of trichostrongylid infective larvae. Until the 60th day of pregnancy, per ewe daily ration was 1.30 kg of a concentrate feed (net energy: 0.844 FU<sub>L</sub>) plus 2.50 kg of alfalfa hay. From the 60th to 100th day of pregnancy, per ewe daily ration was 0.60 kg of the same concentrate plus 2.00 kg of hay. From the 100th day of pregnancy, per ewe daily ration was 0.50 kg of a reduced energy concentrate feed (net energy: 0.748 FU<sub>L</sub>) plus 0.50 kg of alfalfa hay for ewes with one foetus; during that period, respective figures for ewes with two foetuses were 0.60 kg and 0.50 kg and for ewes with three foetuses were 0.80 kg and 0.50 kg. In the experimental ewes, examinations performed, starting on the 120th day of pregnancy and until parturition, were: parasitological examinations of faecal samples, biochemical examination of blood samples for measurement of  $\beta$ -hydroxybutyrate and glucose concentrations, B-mode and Doppler ultrasonographic examinations of the foetus(es) and of the udder; examinations performed after lambing and until the 5th day after that were: weighing of lambs, examination of the genital tract of ewes and milk yield measurements of the ewes. In 16 ewes, increased  $\beta$ -hydroxybutyrate blood concentrations ( $\geq 1.2$  mmol L<sup>-1</sup> in at least two samplings) were detected and these animals were allocated into group A; the 12 ewes that did not, were allocated into group B. Ewes into group A had greater faecal epg counts and smaller blood concentrations of glucose

than ewes into group B ( $P<0.025$ ,  $P=0.033$ , respectively). There were no differences between ewes of the two groups (A, B) in findings of ultrasonographic examination of foetus(es), except for the blood input into the umbilical artery; in that parameter, values in ewes of group B were significantly greater than values in group A animals. However, there was clear evidence that when ewes were re-allocated into ones which developed and ones which did not develop dystocia, there were significant differences ( $P<0.047$ ) between them in all haemodynamic parameters in the umbilical artery on the 140th and the 145th day of pregnancy. Grey-scale intensity values of mammary parenchyma of ewes in group A were significantly greater than of those of ewes in group B ( $P=0.007$ ); blood input into the mammary gland was significantly greater in ewes of group B than in ewes of group A ( $P<0.05$ ) and differences between the two groups were also identified in pulsatility index and in mean blood velocity ( $P=0.007$ ,  $P=0.036$ , respectively) during the last fortnight of pregnancy. Mean duration of pregnancy was significantly shorter in group A ewes (145.75 days) than in group B ones (148.42 days) ( $P<0.001$ ). There was also a significantly higher incidence risk of peri-parturient problems in group A ewes; incidence risks of dystocia, of perinatal mortality of offspring and of post-partum genital disorders were 0.500, 0.375 and 0.250 in group A and 0.083, 0.083 and 0.000 in group B ewes, respectively ( $P=0.01$ ,  $P=0.039$  and  $P=0.031$ , respectively). Median bodyweight of lambs of ewes allocated into group A was 3.5 kg and that of ewes into group B 4.0 kg ( $P=0.016$ ); there was a significant reverse correlation between blood  $\beta$ -hydroxybutyrate concentrations and lamb birth bodyweight and between nematode epg faecal counts and lamb birth bodyweight ( $P=0.016$  and  $P=0.03$ , respectively).

Section B of the chapter describes work performed and relevant results after challenge of the mammary gland of the experimental animals. Group A ewes were allocated into subgroups A1 ( $n=8$ ) or A2 ( $n=8$ ) and group B ewes into B1 ( $n=8$ ) or B2 ( $n=4$ ). Ewes in A1 or B1 were challenged, on the 5th day after lambing, by deposition of *Mannheimia haemolytica* into one teat duct, whilst ewes in A2 and B2 were controls. Clinical and ultrasonographic examination of the udder and bacteriological and cytological examinations of the mammary secretion were performed, as well as pathological examination of the inoculated teat (after mammilectomy) and of both mammary parenchymas (after biopsy). Mastitis developed in 8/8 ewes of subgroup A1, in 1/8 ewe of subgroup A2, in 4/8 ewes in subgroup B1 and in 0/4 ewes in subgroup B2. Comparisons between subgroups revealed that isolations from A1 or A2 were greater than respective isolations from B1 or B2 (for A1 versus B1,  $P<0.08$ ; for A2 versus B2,  $P>0.3$ ). Also, bacteria were recovered more frequently from tissue samples from A1 than from B1 ( $P=0.008$ ) and from A2 than from B2 ( $P=0.058$ ). After challenge, the duct cistern and the gland cistern were observed as anechoic cavities, with presence of hyperechoic particles therein; progressively also, the ultrasonographic

pattern observed in the parenchyma became markedly heterogeneous, with presence of characteristically coarse formations within. After challenge, there was clear evidence for an increase in the diameter of the external pudendal artery already 6 h after challenge. There was also an abrupt and excessive increase in blood input into the mammary parenchyma of the challenged side (for differences between A1 and B1,  $P < 0.06$ ); mean velocity of blood also increased immediately after challenge; finally, blood input into the teat increased immediately after challenge. The characteristic lymphoid follicles at the border between teat duct and teat cistern were observed in 3/8 ewes in A1 and in 7/8 ewes in B1 ( $P = 0.019$ ). In A1, cumulative score for macroscopic and histological findings in the teat was 18 and 23, respectively; cumulative score for histological findings in the mammary parenchyma ipsilateral to the inoculated teat was 24, whilst scores for A2 were 2, 9 and 5, for B1 were 5, 31 and 16 ( $P \leq 0.05$  compared to results in A1) and for B2 were always 0 ( $P > 0.05$ , compared to results in A2).

The conclusions from the results of the present thesis are summarised herebelow.

- (a) A model has been developed to induce pregnancy toxæmia in ewes for use in experimental studies, at the same time covering satiation requirements of the animals. Parasitism might have further contributed to improving efficacy of the model.
- (b) Ultrasonographic examination of foetuses of ewes with pregnancy toxæmia revealed reduced blood input in the umbilical artery. There was a particularly intense modification of haemodynamic properties in ewes which later developed dystocia.
- (c) Ultrasonographic appearance of the mammary gland of ewes during lactogenesis has been described. Results of B-mode ultrasonographic examination indicated differences between healthy ewes and ewes with pregnancy toxæmia in the development of mammary parenchyma. Smaller blood input into the udder of ewes with pregnancy toxæmia was evident.
- (d) Increased incidence of peri-parturient problems was recorded in ewes with pregnancy toxæmia; there was also increased perinatal mortality in their offspring.
- (e) Pregnancy toxæmia can act as a potential predisposing factor for mastitis in the immediately *post-partum* period. Possibly, impairment of the lymphoid follicular structures present at the border between teat duct – teat cistern could have been the cause of reduced protection of the mammary gland.

## **Publications associated with the present thesis**

The following scientific papers presenting facets of the present thesis, are available:

- I.** M.S. Barbagianni, E. Giannenas, E. Papadopoulos, I.G. Petridis, S.A. Spanos, P.G. Gouletsou, I. Valasi, G.C. Fthenakis (2015) "Pregnancy toxemia in ewes: development of an experimental model and potential interactions with gastrointestinal nematode infections" *Small Ruminant Research*, 133:102-107.
- II.** M.S. Barbagianni, P.G. Gouletsou, I. Valasi, I.G. Petridis, I. Giannenas, G.C. Fthenakis (2015). "Ultrasonographic findings in the ovine udder during lactogenesis in healthy ewes or ewes with pregnancy toxemia" *Journal of Dairy Research*, 82:293-303.
- III.** M.S. Barbagianni, V.S. Mavrogianni, A.I. Katsafadou, S.A. Spanos, V. Tsioli, A.D. Galatos, M. Nakou, I. Valasi, P.G. Gouletsou, G.C. Fthenakis (2015) "Pregnancy toxemia as risk factor for development of mastitis in sheep during the immediately post-partum period" *Small Ruminant Research*, 130:246-251.
- IV.** M.S. Barbagianni, S.A. Spanos, K.S. Ioannidi, N.G.C. Vasileiou, A.I. Katsafadou, I. Valasi, P.G. Gouletsou, G.C. Fthenakis (2016) "Increased incidence of peri-parturient problems in ewes with pregnancy toxemia" *Small Ruminant Research*, 132:111-114.

## **ADVISORY COMMITTEE**

Professor G.C. Fthenakis	Supervisor
Department of Obstetrics and Reproduction, Faculty of Veterinary Medicine, University of Thessaly	
Assistant professor P.G. Gouletsou	Member of the advisory committee
Department of Obstetrics and Reproduction, Faculty of Veterinary Medicine, University of Thessaly	
Assistant professor I.G. Valasi	Member of the advisory committee
Department of Physiology, Faculty of Veterinary Medicine, University of Thessaly	

## **EXAMINATION BOARD**

Professor G.S. Amiridis	Faculty of Veterinary Medicine, University of Thessaly
Professor G.C. Fthenakis	Faculty of Veterinary Medicine, University of Thessaly
Associate professor M. Caroprese	Dipartimento di Scienze Agrarie, degli Alimenti e dell'Ambiente, Università degli Studi di Foggia
Assistant professor P.G. Gouletsou	Faculty of Veterinary Medicine, University of Thessaly
Assistant professor V.S. Mavrogianni	Faculty of Veterinary Medicine, University of Thessaly
Assistant professor I. Valasi	Faculty of Veterinary Medicine, University of Thessaly
Lecturer I. Giannenas	School of Veterinary Medicine, Aristotle University of Thessaloniki

# TABLE OF CONTENTS

	Page
<b>ΠΕΡΙΛΗΨΗ</b>	2
<b>ΣΥΜΒΟΥΛΕΥΤΙΚΗ ΕΠΙΤΡΟΠΗ - ΕΞΕΤΑΣΤΙΚΗ ΕΠΙΤΡΟΠΗ</b>	7
<b>ABSTRACT</b>	9
<b>ADVISORY COMMITTEE - EXAMINATION BOARD</b>	13
<b>TABLE OF CONTENTS</b>	14
<b>GENERAL INTRODUCTION</b>	16
Preface - Objectives of the thesis	17
Acknowledgments - Ευχαριστίες	18
<b>CHAPTER I</b>	21
<b>REVIEW OF THE LITERATURE</b>	
<b>PREGNANCY TOXAEMIA</b>	22
Introduction	22
Aetiology and predisposing factors	22
Pathogenetic pathways	23
Effects of the disease in immunity	27
Manifestations	27
Diagnosis	30
Control	30

<b>CHAPTER II</b>	<b>34</b>
<b>RESEARCH WORK</b>	
<b>A. BEFORE CHALLENGE OF THE MAMMARY GLAND</b>	
<b>OF THE EXPERIMENTAL EWES</b>	<b>35</b>
Materials and methods	35
Results	47
<b>B. AFTER CHALLENGE OF THE MAMMARY GLAND</b>	
<b>OF THE EXPERIMENTAL EWES</b>	<b>74</b>
Materials and methods	74
Results	81
<b>GENERAL DISCUSSION</b>	<b>107</b>
Introduction	108
Definitions regarding pregnancy toxaemia	109
The experimental model used for induction of pregnancy toxaemia	112
Doppler ultrasonographic examination of the foetus	113
Ultrasonographic examination of the mammary glands during lactogenesis	116
Peri-parturient problems in ewes with pregnancy toxaemia	119
Association of pregnancy toxaemia with mastitis during the subsequent lactation period	121
Epilogue	125
<b>REFERENCES</b>	<b>127</b>

# GENERAL INTRODUCTION



## Preface - Objectives of the thesis

The *puerperium* starts with completion of parturition. After lambing, the reproductive organs continue their increased role, although significance of the genital system obviously decreases and that of the mammary glands increases. Mastitis is the most significant disease of the udder, causing extensive financial losses in dairy sheep flocks. Moreover, mastitis has been recently described as the most significant welfare problem in sheep, independently of production type and management system (European Food Safety Authority 2014).

Although aetiology of mastitis has been clarified, there are still questionmarks regarding risk factors for the disease. Various factors (e.g., husbandry, genetic, environmental, diseases) have been described as predisposing sheep to mastitis. Inappropriate feeding practices may contribute to mastitis. Increased incidence risk of clinical and subclinical mastitis in ewes with vitamin A deficiency has been reported, possibly as the result of reduced integrity and functionality of the epithelial defences of the mammary gland in affected animals (Koutsoumpas et al. 2013). Similarly, selenium deficiency (Giadinis et al. 2011) or increased consumption of cottoncake meal (containing increased gossypol concentration) (Fthenakis et al. 2004) have been reported to contribute to development of mastitis in ewes, possibly as the result of impeded cellular defences of the affected ewes.

Pregnancy toxaemia is the most important metabolic disease of pregnant ewes. It occurs as the consequence of inappropriate management during pregnancy, more often during the last month of gestation. Clinical disease requires emergency intervention and increased medical attention, as it may lead to death of the affected animals (Brozos et al. 2011). The extent of the condition, in which ewes have increased  $\beta$ -hydroxybutyrate blood concentrations but do not develop clinical signs relevant to pregnancy toxaemia (potentially termed 'subclinical pregnancy toxaemia') is not known. Increased  $\beta$ -hydroxybutyrate blood concentrations in pregnant ewes are recognised as a risk factor for development of clinical signs relevant to pregnancy toxaemia and are used diagnostically for monitoring the situation within a flock (Brozos et al. 2011). Further, potential adverse effects of the problem have not been described in detail.

The present thesis focuses in experimentally induced pregnancy toxaemia in sheep and its potential effects in the mammary gland, with the general objective to increase relevant available knowledge. Specific objectives of the thesis are as follows.

- The development of a valid model for studying pregnancy toxaemia, whilst maintaining appropriate satiation requirements in the experimental animals.

- The recording, by means of ultrasonographic examination, of changes occurring in the foetus and in the udder of ewes with pregnancy toxaemia.
- The evaluation of consequences of pregnancy toxaemia in affected ewes around the lambing period.
- The evaluation of the potential predisposing role of pregnancy toxaemia in development of mastitis in the immediately post-partum period.

The present thesis has been carried out at the Department of Obstetrics and Reproduction of the Veterinary Faculty of the University of Thessaly. Research work started in 2012 and was carried out until the end of 2014; it was followed by analysis of results and writing up of the thesis. The thesis was financially supported by departmental funds.

A part of the work described in the thesis was carried out in collaboration with the Department of Surgery and the Department of Animal Husbandry and Nutrition of the Faculty; this latter collaboration, at subsequent stages of the work, continued with the Department of Nutrition of the Faculty of Veterinary Medicine of the Aristotle University of Thessaloniki. Parts of the work were also carried out at the private Medical Pathology Laboratory in Karditsa.

## **Acknowledgments - Ευχαριστίες**

Κατ' αρχήν, ευχαριστώ από καρδιάς τον επιβλέποντά μου, καθηγητή του Τμήματος Κτηνιατρικής του Πανεπιστημίου Θεσσαλίας κ. Γεώργιο Φθενάκη, ο οποίος μού έδωσε την δυνατότητα να εκπονήσω αυτή τη διατριβή και ήταν αρωγός, συνοδοιπόρος και υποστηρικτής σε κάθε φάση της διαδρομής μου. Η συμβολή του ήταν σημαντική στην επιλογή του θέματος, στην ανεύρεση εξοπλισμού και κατάλληλου ανθρώπινου δυναμικού για την εκπαίδευσή μου, καθώς επίσης και στη συγγραφή της διατριβής, όπου με τις συνεχείς εύστοχες και ουσιαστικές παρατηρήσεις του μού έδινε πάντα το έναυσμα για πιο διεξοδική διερεύνηση του θέματος. Επίσης, τον ευχαριστώ διότι μού δίδαξε τι θα πει υπομονή, επιμονή και αγάπη για την έρευνα, ενώ ταυτόχρονα φρόντιζε να με στηρίζει βρίσκοντας πάντα λύση στα φαινομενικά άλυτα προβλήματα, καθώς και εξασφαλίζοντάς συνεχώς την απαραίτητη χρηματοδότηση για την πραγματοποίηση της έρευνας. Η αφοσίωση και η εμπιστοσύνη του, μού χάρισαν την απαραίτητη αυτοπεποίθηση για την επιτυχή διεξαγωγή της μελέτης.

Ευχαριστώ την επίκουρη καθηγήτρια του Τμήματος Κτηνιατρικής του Πανεπιστημίου Θεσσαλίας κ. Ε. Βαλάση για την εκπαίδευσή μου στην υπερηχογραφία, κυρίως στην τεχνική της

διάγνωσης της εγκυμοσύνης, για την προθυμία της να βρίσκεται πάντα δίπλα μου και να ενισχύει με τις γνώσεις της την ερευνητική προσπάθειά μου και κυρίως για τις καθημερινές πολύτιμες για εμένα συμβουλές και συζητήσεις μας. Επίσης, ευχαριστώ την επίκουρη καθηγήτρια του Τμήματος Κτηνιατρικής του Πανεπιστημίου Θεσσαλίας κ. Π. Γκουλέτσου για την εκπαίδευσή μου στην υπερηχογραφία και κυρίως στην τεχνική εξέτασης των μαστών και θηλών, για την παρουσία της και τις χρήσιμες συμβουλές της κατά τη διάρκεια της πειραματικής μελέτης και για την αμέριστη ηθική συμπαράστασή της.

Ευχαριστίες επίσης απευθύνω:

Στην επίκουρη καθηγήτρια του Τμήματος Κτηνιατρικής του Πανεπιστημίου Θεσσαλίας κ. Β. Τσιώλη, για τη διεξαγωγή όλων των χειρουργικών επεμβάσεων, την υπομονή της και τις ατέλειωτες ώρες που αφιέρωσε για την επιτυχή διεξαγωγή της πειραματικής μελέτης.

Στον καθηγητή του Τμήματος Κτηνιατρικής του Πανεπιστημίου Θεσσαλίας κ. Γ.Σ. Αμοιρίδη, για τις πολύτιμες συμβουλές του, τη συμπαράστασή του σε όλη τη διαδικασία του πειραματικού μέρους και τη βοήθειά του σε θέματα υπερηχογραφικής εξέτασης.

Στην επίκουρη καθηγήτρια του Τμήματος Κτηνιατρικής του Πανεπιστημίου Θεσσαλίας κ. Β. Μαυρογιάννη, για την καθοδήγησή της στις μεθόδους κυτταρολογικής και ιστολογικής εξέτασης και για τη σημαντική συμβολή της στην αξιολόγηση των σχετικών ευρημάτων.

Στον διδάκτορα του Τμήματος Κτηνιατρικής του Πανεπιστημίου Θεσσαλίας κ. Ι. Πετρίδη, για τη βοήθειά του σε τεχνικά θέματα χειρισμού της συσκευής υπερηχογραφίας και για την εκπαίδευσή μου στην υπερηχογραφία Doppler.

Στον λέκτορα του Τμήματος Κτηνιατρικής του Αριστοτελείου Πανεπιστημίου Θεσσαλονίκης κ. Η. Γιάννενα, για την επιμέλεια της διατροφικής διαχείρισης των πειραματοζώων και για τον σχεδιασμό του κατάλληλου σιτηρεσίου για πρόκληση τοξιναιμίας εγκυμοσύνης με ταυτόχρονη διατήρηση της ευζωίας των ζώων.

Στον αναπληρωτή καθηγητή του Τμήματος Κτηνιατρικής του Αριστοτελείου Πανεπιστημίου Θεσσαλονίκης κ. Η. Παπαδόπουλο, για την επιμέλεια των παρασιτολογικών θεμάτων που αναφέρονται στη διατριβή.

Στον υπεύθυνο τεχνικό της εταιρείας Esaote κ. Ε. Σαμαρά, για την προθυμία του να αντιμετωπίζει όλα τα τεχνικά προβλήματα του υπερηχογράφου και να υποδεικνύει τρόπους, ώστε, μέσω της συσκευής, να εφαρμόζονταν οι θεωρητικές προσεγγίσεις.

Στον αναπληρωτή καθηγητή του Τμήματος Κτηνιατρικής του Πανεπιστημίου Θεσσαλίας κ. Α. Γαλάτο, για τον σχεδιασμό κατάλληλου αναισθητικού πρωτοκόλλου για τα πειραματόζωα, για την αναισθησιολογική επίβλεψή τους και για τη συνεχή εμπύχωσή του.

Στην επίκουρη καθηγήτρια του Τμήματος Κτηνιατρικής του Πανεπιστημίου Θεσσαλίας κ. Κ. Θεοδοσιάδου, για την παραχώριση χρήσης εξοπλισμού και για τη στήριξη και τη συνεχή ενθάρρυνση.

Στον επίκουρο καθηγητή του Τμήματος Ζωικής Παραγωγής του ΤΕΙ Θεσσαλίας κ. Σ. Παπαδόπουλο, για την παραχώρηση χρήσης εξοπλισμού.

Στον αναπληρωτή καθηγητή της Κτηνιατρικής Σχολής του Πανεπιστημίου του Liverpool κ. P. Cripps, για τις πολύτιμες ιδέες του στη διαχείριση και σωστή ανάλυση των αποτελεσμάτων.

Στον επίκουρο καθηγητή του Τμήματος Κτηνιατρικής του Πανεπιστημίου Θεσσαλίας κ. Ν. Σολωμάκο, για τη συμβολή του στη συλλογή μαστών από τα σφαγεία που αποτέλεσαν σημείο αναφοράς στην επιλογή κατάλληλης τεχνικής για τη θηλεκτομή.

Στη διδάκτορα του Τμήματος Ιατρικής του Πανεπιστημίου Θεσσαλίας κ. Μ. Νάκου, για την ετοιμασία των ιστοπαθολογικών παρασκευασμάτων.

Στον καθηγητή του Τμήματος Κτηνιατρικής του Πανεπιστημίου Θεσσαλίας κ. Χ. Μπιλλίνη, για την παραχώρηση χρήσης εξοπλισμού.

Στη συνάδελφο κ. Α. Κατσαφάδου, για την ουσιαστική βοήθειά της στις δειγματοληψίες και τη συμπαράστασή της στη διάρκεια της πειραματικής μελέτης.

Στους συναδέλφους Σ. Σπανό, Ν.Γ. Βασιλείου και Κ. Ιωαννίδη, για τη σημαντική βοήθειά τους στη φροντίδα και εξέταση των ζώων, όποτε αυτή χρειαζόταν.

Στην παρασκευάστρια του Τμήματος Ζωικής Παραγωγής του Τεχνολογικού Εκπαιδευτικού Ιδρύματος Ηπείρου κ. Κ. Φώτου, για την παρασκευή μικροβιολογικών υποστρωμάτων.

Στον κ. Α. Τρούκη και την κ. Φ. Ριζάβα, για τη φροντίδα των πειραματοζώων και τη βοήθειά τους στις δειγματοληψίες.

Τέλος, ευχαριστώ ιδιαίτερα τον σύντροφό μου για την ψυχολογική και οικονομική στήριξή του, την υπομονή του και ανοχή του να διασχίσει μαζί μου αυτήν την επίπονη διαδικασία. Ένα μεγάλο ευχαριστώ επίσης απευθύνω στους γονείς μου που με παρακίνησαν να ξεκινήσω αυτήν τη διατριβή, μοιράστηκαν μαζί μου όλες τις ανησυχίες και τους προβληματισμούς μου και με στήριξαν με κάθε δυνατό τρόπο, καθώς χωρίς αυτούς δεν θα είχα καταφέρει να ολοκληρώσω την όλη μελέτη.

# **CHAPTER I**

## **REVIEW OF THE LITERATURE**

# PREGNANCY TOXAEMIA

## Introduction

Pregnancy toxaemia is the most common metabolic disorder in sheep, which develops during late gestation. It was first described by McClymont and Setchell in 1955. Pregnancy toxaemia develops as a hypoglycaemic encephalopathy and hyperketonaemia and is a significant cause of deaths in peri-parturient ewes (Mavrogianni and Brozos 2008).

In affected ewes, of major welfare concern are the reduced feed intake occurring during the disease and the hunger associated with it. A further, less obvious, welfare effect is the depletion of body fat reserves during late gestation (Robinson 1986). Welfare concerns can also be raised in relation to the newborn lambs, which are born with suboptimal bodyweight (Lynch et al. 1990, Mukasa-Mugerwa et al. 1994), receive reduced quantity of colostrum of inferior quality (Banchemo et al. 2002, 2004) and are subjected to inappropriate mothering behaviours by their dams (Putu et al. 1988, Dwyer et al. 2003, Everett-Hincks et al. 2005).

## Aetiology and predisposing factors

Pregnancy toxaemia develops as the result of decreased energy intake by animals, in relation to their particularly increased requirements at the end of gestation. The disease is caused by abnormal metabolism of carbohydrates and fats during pregnancy (Mavrogianni and Brozos 2008, Brozos et al. 2011). It is exacerbated by the reduction of the space available in the rumen, as a consequence of the progressively increasing size of the gravid uterus (Guard 1995, Panousis et al. 2001).

Pregnancy toxemia occurs mainly in older, multiparous animals (Bickhardt et al. 1998), with most cases recorded during the last 14 days before lambing (Henze et al. 1998). Ewes bearing two or three foetuses are more susceptible to the disease (Pethick and Lindsay 1982, Everts and Kuiper 1983), as in these animals energy deficit is more common due to their increased energy requirements. Some sheep breeds are more susceptible to the disease, possibly associated with increased lipolysis and blood concentrations of non-esterified fatty acids genetically occurring in these breeds (Duehlmeier et al. 2011, 2013).

High-producing ewes are more susceptible to the disease (Rook 2000, Brozos et al. 2011), as they are at greater risk for negative energy balance. Moreover, bodyweight loss can also predispose ewes to the disease (Fthenakis et al. 2012); homeostatic mechanisms respond to changes in energy balance during pregnancy and, depending on the duration and severity of the energy deficit period, may lead to reductions in muscle and fat tissue (Chilliard et al. 2000, Faulconnier et al. 2001), which are responsible for bodyweight loss by animals.

Adverse weather conditions (e.g., low temperatures, windy periods) also predispose to the disease, as they increase energy requirements of animals. Hence, the incidence risk of the disease is increased during the winter months (Panousis et al. 2001).

Wasting diseases (e.g., parasitic infections, paratuberculosis) also predispose ewes to pregnancy toxemia by increasing energy depletion. Further, diseases that affect feed intake (e.g., dental problems, lameness) can also act as risk factors for development of pregnancy toxemia (Panousis et al. 2001).

Finally, factors not directly associated with the animals (e.g., financial situation of the farmer) can also predispose to development of the disease within a flock, as they may affect management practices necessary for prevention of the disease.

## **Pathogenetic pathways**

Approximately 60% of foetal growth takes place during the last stage of gestation (Twardock et al. 1973), during which period up to 40% of the available glucose in the pregnant female is directed to the foetoplacental unit, in order to satisfy energy demands (Hay et al. 1983, Rook 2000). During that stage, the female animal's metabolism, in order to meet growing energy demands of the developing foetus(es) increases hepatic gluconeogenesis, reduces glucose uptake by maternal peripheral tissues and enhances placental glucose transport capacity (Bell and Bauman 1997). Foetuses require glucose and amino acids, for energy and for protein deposition. If energy requirements are not met, it would lead to increased rate of lipolysis and an increase in rate of ketone body production. Further, glucose demands are increased if multiple foetuses are borne (Sargison et al. 1994), in which case increased hepatic entry rates of  $\beta$ -hydroxybutyrate (Pethick and Lindsay 1982), decreased glucose and insulin blood concentrations (Marteniuk and Herdt 1988, Sigurdsson 1988) and increased ketone body blood concentrations (Varnam et al. 1978) occur.

In sheep, glucose is the most important energy source, with insulin being the major regulator of energy partitioning (Hart 1983); therefore, glucose deficiency develops in case of significant malfunction of sheep metabolism. Bergman (1973) has reported that increased glucose synthesis was usually associated with decreased ketone body formation. Conversely, in sheep with reduced feed intake, which causes significantly reduced glucose turnover, ketonogenesis increases (Kronfeld and Simesen 1961). In sheep, glucose blood concentrations are physiologically smaller during pregnancy than during the lactation period (Bickhardt and Konig 1985, Henze et al. 1994), which explains the greater risk for development of the disease in pregnancy than in lactation. This may be the consequence of increased glucose utilisation by the growing foetus(es) (Rook 2000) or of impaired hepatic gluconeogenesis in pregnant animals (Schlumbohm and Harmeyer 2004). Further, hypocalcaemia, which may also be the result of imbalanced nutritional regime, depresses endogenous glucose production (Schlumbohm and Harmeyer 1990, 1999, 2003) and that way contributes to development of pregnancy toxemia (Simensen 1971, Bath 1983), with which it often coexists. The positive correlation between blood concentrations of glucose and insulin indicates that variations in insulin blood concentrations between reproductive stages can be the effect of variations in glucose blood concentrations, i.e., in cases of increased blood glucose values, there is increased glucose-stimulated pancreatic insulin secretion (Duehlmeier et al. 2011). The decreased blood concentrations of insulin in late pregnancy may also be the result of increasing lipolysis (Regnault et al. 2004); the observation of Boden and Shulman (2002) that long-term increased blood concentrations of non-esterified fatty acids inhibit secretion of insulin because of a 'lipotoxicity' of fatty acids in pancreatic-cells, lends support to the hypothesis.

It thus becomes evident that negative energy balance results in glucose deficiency, which is the initial factor leading in pregnancy toxemia (Rook 2000, Van Saun 2000, Kulcsar et al. 2006).

During the last stage of pregnancy, ewes may develop a decreased insulin sensitivity, which reduces glucose use by peripheral tissues and increases dependence on lipid and ketone metabolism; this can be exaggerated by undernutrition (Pettersen et al. 1993). Glucose for foetuses is transferred in a different way than in maternal tissues: the placental glucose transport takes place via the insulin independent glucose transporter 1 (GLUT1), while glucose uptake in the maternal skeletal muscles and adipose tissue is mediated by the insulin dependent glucose transporter 4 (Anderson et al. 2001).

Additionally, during negative energy balance, various endocrinological parameters are modified. Schwartz et al. (2000) and Woods (2005) have found that various factors (e.g., insulin,



leptin, insulin like growth factor-1, ghrelin) involved in regulation of feed intake and energy usage are affected by the energy imbalance. Under-nourished pregnant ewes have been found with smaller concentrations of leptin (Thomas et al. 2001, Bispham et al. 2003) and insulin (Wallace et al. 1999, Luther et al. 2007) compared to ewes receiving balanced rations; their deficiency may further affect feed intake and nutrient mobilisation (Schwartz et al. 2000, Woods and Seeley 2000). Further, intracerebroventricular infusion of sheep with leptin has been shown to decrease their feed intake (Henry et al. 2001, Morrison et al. 2001); similar findings have been reported after intracerebroventricular infusion of insulin (Foster et al. 1991).

Long-standing undernutrition has been found to lead to suppression of the hypothalamic-pituitary-adrenal axis and reduced cortisol blood concentrations in pregnant ewes (Jaquiery et al. 2006, Verbeek et al. 2012). Potentially, cortisol could play an indirect role, through its influence on leptin, as leptin and cortisol mutually regulate each other and affect energy balance (Leal-Cerro et al. 2001). Possibly, insulin like growth factor-1 plays an important role in carbohydrate and protein metabolism, as well as in cell replication and differentiation (Gluckman et al. 1987, Jones and Clemmons 1995); furthermore, it is essential in regulation of growth of the placenta and foetus (Wathes et al. 1998, Fowden 2003) and maternal nutrition may influence foetal development through this (Bassett et al. 1990, Brameld et al. 2000), directly or through partitioning of nutrients between tissues (Fowden 2003).

In order to cover energy demands, pregnant ewes mobilise fatty acids from peripheral tissues for transport to the liver and extra-hepatic tissues. Uptake of fatty acids by the liver is proportional to their concentration in the blood. In the liver, fatty acids are stored as triglycerides and may be transformed to ketone bodies, be oxidised or be incorporated into lipoproteins in the Golgi device for release into the general circulation (Lean 2002). The mobilisation of long chain fatty acids from adipose tissue leads to increase of blood concentrations of non-esterified fatty acids and of ketone bodies. Therefore, blood concentrations of non-esterified fatty acids reflect fat mobilisation and those of  $\beta$ -hydroxybutyrate reflect fat oxidation in the liver (LeBlanc 2010); that way also, measurement of glucose and  $\beta$ -hydroxybutyrate blood concentrations can also be used to assess the nutritional status of pregnant ewes in late gestation (Russel 1984). As ketones are metabolic acids, acidosis may also develop in cases of pregnancy toxemia (Lean 2002).

The ketone bodies are oxidised, that way providing fuel for energy production. Many peripheral tissues (e.g., heart, skeletal muscles, kidneys, developing mammary glands) gain energy from the oxidation of ketone bodies (Harmeyer and Schlumbohm 2006). At the end, incorporation of acetyl coenzyme A into the Krebs cycle is inhibited and leads in accumulation of triglycerids into the hepatic cells; this lead to transformation of the acetyl coenzyme A to acetate,

$\beta$ -hydroxybutyrate and acetone. Lack of oxaloacetate leads to incomplete break-down of non-esterified fatty acids, leading to enhanced ketonogenesis and consequently increased  $\beta$ -hydroxybutyrate blood concentrations, which inhibit gluconeogenesis and contribute to further hypoglycaemia of the pregnant animal; in turn, this enhances the energy deficit and continues the production of ketone bodies in a vicious circle, concluding in pregnancy toxaemia (Schlumbohm and Harmeyer 2004).

Possibly, additional endocrinological factors may be involved in development of the disease (Schlumbohm and Harmeyer 2004). Cortisol blood concentrations can have an effect in lipid mobilisation and ketonaemia during late gestation (Reid 1960), whilst increased blood concentrations of progesterone and oestradiol have been recorded in ewes that received reduced energy after the 100th day of gestation (Lemley et al. 2014).

Further, the insufficient oxidation of fatty acids leads in accumulation of triglycerides in hepatic cells (Lean 2002), which, given that mechanisms of their metabolism in or export from the liver are impaired, results in hepatic lipidosis. This can be a further factor contributing to pregnancy toxaemia. In the absence of propionate influx from the rumen, there is a reliance on amino acids and lactate to provide carbon chains for gluconeogenesis. Sheep more susceptible to pregnancy toxaemia have been found with hepatocytes producing smaller amounts of glucose from glucogenic precursors, including propionate, lactate and alanine. The impaired liver function observed before development of pregnancy toxaemia has been attributed to a diversion of the metabolism of polyunsaturated fatty acids from phospholipids to triacylglycerols; this could explain, at least partly, the degenerative changes in membrane and subcellular organelle structure within the organ (Henderson et al. 1982). Recently, Chen et al. (2012) have reported that hepatic expression of *angiopoietin-like protein 3* gene (which encodes a member of a family of proteins expressed predominantly in the liver) was significantly down-regulated in animals with pregnancy toxaemia (Oike et al. 2005). Angiopoietin-like protein 3 is a secretory glycoprotein expressed only in the liver (Conklin et al. 1999), which can mediate in lipolysis of adipocytes (Shimamura et al. 2003) and contribute in carbohydrate metabolism and lipid metabolism (Robciuc et al. 2010).

The foetus has also a strong demand for amino acids, which are primarily used for provision of energy (rather than coverage of structural needs). This demand is met from dietary sources and mobilisation of protein reserves. Sheep having received, in early gestation, increased energy before subsequent exposure to an energy deficit were found to be able to maintain optimum gluconeogenesis than animals which had received an energy deficient nutritional regime at the start of gestation. Anyway, after protein reserves would be depleted, there is little possibility

for pregnant animals to produce glucose if the diet failed to provide sufficient protein or propionate (Lean 2002).

Hence, the reduced ability of ewes in late gestation to utilise ketone bodies is another key factor in the pathogenesis of pregnancy toxaemia. Hyperketonaemia is a characteristic feature of pregnancy toxaemia and explains, at least in part, why the disease occurs much more often in the last stage of pregnancy rather than in the early lactation period when energy output would be greater.

### **Effects of the disease in immunity**

The increased blood concentrations of glucocorticoids and ketone bodies present in pregnancy toxaemia have been considered to contributing to the suppression of cell-mediated immunity in affected animals. Sartorelli et al. (1999, 2000) have reported that ovine neutrophils after incubation with  $\beta$ -hydroxybutyrate showed impairment of particle uptake, chemotaxis and bactericidal activity. Lacetera et al. (2001) have commented that pregnancy toxaemia could alter cell-mediated immunity of ewes, as the reduced concentration of glucose coupled with the potentially toxic effects of  $\beta$ -hydroxybutyrate and non-esterified fatty acids had negative effects in the proliferation of mononuclear cells of sheep; they have also indicated that antigen-specific IgG production in ewes with pregnancy toxaemia was also negatively affected.

## **Manifestations**

### **Clinical features**

Initial clinical features of pregnancy toxaemia may not be noticed. An observation leading to suspicion of pregnancy toxaemia is that pregnant ewes would be standing away from the flock, rejoining it if prompted to, or staying in a corner of the animal house, often leaning on a wall, especially if enclosed into a pen. Later, mild neurological signs develop, including depression or suppression; affected animals have only instinctive responses to stimuli applied to the ocular, auditory and/or olfactory exteroceptors. However, often these initial symptoms may not be observed, with animals appearing to be clinically normal at that time, with more pronounced

symptoms developing later. The hyperketonaemia, which develops progressively, exerts a depressive action on appetite of affected animals, which becomes selective; animals would gradually stop eating concentrate feed, hay and, finally, straw (Scharrer 1999, Rossi et al. 2000, Panousis et al. 2001, Sargison 2007, 2008).

At a later stage of the disease, sheep would show marked depression and little or no reaction to approach by people. Ocular and/or auditory reflexes could be depressed or absent at this stage and the sheep would often walk into other animals or objects. With further depression of consciousness, the affected animals would remain standing, often with the head against a wall, for many hours at a time with no attempt to move. They would show no reaction to humans or other situations that usually are stressful for a flock. Neuromuscular disturbances may also be observed, varying in intensity and degree of spread; these include myoclonic twitching of the ears, periorbital muscles and muzzle and clonic intermittent contractions of muscle groups of the head, back and limbs, lasting for seconds to few minutes, at a rate of one to three per second. These usually develop suddenly, with rapid spread over the muscle groups. Respiration rate and rectal temperature are not modified (Panousis et al. 2001, Sargison 2007, 2008).

Eventually, affected ewes would go in sternal or lateral recumbency and usually make no attempt to rise on stimulation, having complete lack of reaction to the environment, except for lower reflexes. Some sheep may pass through short periods during which they might be hypersensitive to stimuli, standing quivering and reacting with a violent start to any sudden stimulus, although changes in consciousness are observed. Paddling movements of the limbs are often recorded in ewes in lateral recumbency. Chewing movements, manifested by teeth grinding and vigorous licking movements, can be observed (Panousis et al. 2001, Sargison 2007, 2008).

Animals show loss of eye reservation reflex (response by blinking and/or movement of the head to a hand moved suddenly towards the eye), of pupillary light reflex (response of pupil, by dilation, to closing of the lid and shading of the eye, then, by constriction, to sudden exposure to bright light) and frequently of auditory reflex (response by movement of the head or ears to sudden shouting near the ears). If food is put in the mouth of sheep, it may be chewed and swallowed (Panousis et al. 2001, Sargison 2007, 2008).

It is noteworthy that animals can show a marked clinical improvement, if foetuses die *in utero*. However, this would be often followed by deterioration of the clinical condition after one to two days. Death is the most common outcome, if treatment would not be instigated at an early stage of the disease. In any case, all sheep do not necessarily show all the symptoms or pass uniformly through all stages of depression of consciousness.

## Paraclinical findings

The salient paraclinical finding is the increased blood concentration of  $\beta$ -hydroxybutyrate, which characterises the disease. In animals with clinical signs of the disease, values of the parameter often are above 2.5 to 3.0 mmol L<sup>-1</sup> (Sargison et al. 1994). Acetate blood concentrations are also above 0.2 mmol L<sup>-1</sup>. In such cases, there is also marked hyperketonuria that becomes evident even by using semi-quantitative measuring methods (Hindson and Winter 1990, Panousis et al. 2001, Sargison 2007, 2008).

Hypoglycaemia is another salient feature of the disease, with values of glucose blood concentrations being <50 mg dL<sup>-1</sup> or even <35 mg dL<sup>-1</sup> in severe cases of the disease. It is however noteworthy that, if foetuses die *in utero*, glucose values may be within the reference range (Panousis et al. 2001).

Hypocalcaemia can be a feature of the disease, as decreased feed intake reduces intestinal calcium absorption, lowering blood concentration of calcium (Harmeyer and Schlumbohm 2006). Increased aspartate aminotransferase activity and decreased albumin blood concentration may be the result of impaired liver function, due to fat infiltration of that organ (Radostits et al. 2007, Smith and Sherman 2009).

As the disease can lead to impairment of renal function with glomerular lesions (Ferris et al. 1969), renal dysfunction and decreased creatinine clearance may occur. These can be manifested with increased blood concentrations of creatinine and blood urea nitrogen in affected ewes (McCausland et al. 1974).

Haematological findings can include increased (due to dehydration) (Smith and Sherman 2009) or decreased (due to rupture of erythrocytes) (Tharwat and Al-Sobayil 2014) haematocrit and haemoglobin concentration values. The rupture of erythrocytes would be the result of endotoxaemia (Smith and Sherman 2009). Anaemia in pregnant ewes would lead to increased pulmonary oxygen utilisation, hence decreased partial oxygen pressure in the blood vessels, including the umbilical artery. Increased blood concentration of cardiac troponin I has also been indicated to occur in cases of the disease (Tharwat et al. 2012).

Changes in cerebrospinal fluid include reduced glucose concentrations (Scott et al. 1995), but no alterations in other parameters (Scott 1992).

## Post-mortem findings

Carcasses are often in poor condition up to emaciated, as well as dehydrated. Liver lipidosis may be evident; the liver is usually swollen, pale and friable. Histologically, there may be distension of the parenchymatous cells with presence of fatty globules. Fat content in the liver can be increased up to 30% w/w, indicating triglyceride saturation (Sargison 2007).

The adrenal glands can be hypertrophied and pale or hyperaemic. Lesions in the lungs are usually associated with the prolonged recumbency. Dead fetuses, often decomposed, found in the uterus are a principal feature of the disease. Histological lesions in the brain include nuclear swelling, hypertrophy and proliferation and cerebrocortical necrosis (Panousis et al. 2001, Sargison 2007, 2008).

## Diagnosis

The clinical diagnosis takes into account the gestational stage of the animal and the combination of the relevant clinical findings. The post-mortem examination provides little diagnostic support as findings are non-specific. Confirmation of clinical diagnosis of pregnancy toxemia is achieved by measurement of  $\beta$ -hydroxybutyrate blood concentration and identification of the increased values of the parametre; availability of rapid measurement equipment makes estimation of the parametre possible even in field conditions (Panousis et al. 2012). Measurement of glucose blood concentration is also of help, whilst measurement of calcium blood concentration does not provide significant diagnostic support, but should be performed, as hypocalcaemia is often a simultaneous problem (Panousis et al. 2001, Sargison 2007).

## Control

## Prognosis

In cases of pregnancy toxemia, prognosis should always be cautious. The efficacy of the treatment depends upon timely diagnosis of the problem and early start of the treatment regime. Treatment of the disorder should be based on two general principles: (i) administration of energy

sources and (ii) removal of factors increasing energy requirements of affected animals, which may not be always applied successfully (Brozos et al. 2011).

However, treatment may still fail, even after a timely start. In animals with signs of the terminal stage of the disease (neurological signs, blindness, recumbency), it often leads to transient improvement of the general condition of the animal, which could subsequently deteriorate; finally, the animal would die. Sargison (1995) has reported that, despite a full course of treatment of toxæmic ewes, only one third of affected ewes would likely survive. Evaluation of glucose blood concentration in animals with pregnancy toxæmia may be used as an indicator of viability of foetuses (Lima et al. 2012).

In such cases, for welfare reasons, euthanasia of affected animals should be considered before instigation of treatment. Sub-standard welfare of sick animals adds to the financial constraints of treatment, which can be expensive, but often fruitless. Further, even if treated intensively, ewes may die after parturition (Henze et al. 1998). Hence, possibly euthanasia might be the best approach for the welfare of affected animals and depending on the circumstances (Caroprese et al. 2016).

## Treatment of affected ewes and their newborn lambs

Treatment of cases of pregnancy toxæmia has been recently reviewed in detail by Brozos et al. (2011). In general, it is based in the administration of energy sources (e.g., glucose, propylene glycol, dextrose), for which various schemes and routes of administration have been established (Buswell et al. 1986, Andrews 1997, Rook 2000, Sargison 2007, Brozos et al. 2011). Administration of various other agents (e.g., recombinant bovine somatotropin, insulin, flunixin meglumine) as part of the therapeutic regime has also been reported (Andrews 1998, Henze et al. 1998, Zamir et al. 2009), but these have not been widely employed in clinical conditions.

Removal of foetuses by induction of lambing or caesarean section is also part of the entire therapeutic scheme, as it leads to decrease of energy requirements of the pregnant animal (Sargison 2007, 2008, Brozos et al. 2011). Usually, after removal of foetuses, the general condition of the animal is improved. Often however, it may deteriorate again, especially if the foetus(es) had died *in utero*.

Newborn lambs from ewes with pregnancy toxæmia require increased care. These usually have a suboptimal birth weight, are stressed and may be premature. If their cardiac or respiratory function is weak, their body should be massaged to induce respiration. Doxapram hydrochloride (5-10 mg per lamb) should be administered by intravenous or subcutaneous injection or by

sublingual dropping (Monin 1994). If the dam cannot produce adequate amount of colostrum, then the newborns should be given colostrum from another ewe in the flock or from the 'colostrum bank' of the farm at a dose of 50 mL kg<sup>-1</sup> bw, 4 times in the first 24 hours of life. Subsequently, it should be evaluated whether the ewe/doe would be able to produce enough milk for feeding the newborns; if this is not considered to be possible, artificial feeding should be undertaken (Brozos et al. 2011).

### Care for other ewes in a flock with affected animals

Pregnancy toxemia should be considered as a flock problem. If clinical cases of the disease are diagnosed in a flock, then clinically health animals in that flock are potentially at risk to also develop the disease (Brozos et al. 2011). Therefore, appropriate measures must be taken for these animals. Risk factors, at individual and flock level, should be evaluated at the time of attending any ill animals for initiating remedial actions.

A broad-spectrum anthelmintic treatment course should be administered in all animals if a diagnosis of gastrointestinal parasitism can be supported. Then, pregnant animals could be grouped according to body condition score and to stage of pregnancy, in order to improve feeding appropriately and according to the needs of each group, as well as to avoid wasteful feeding to animals. If grouping of animals is not possible, high-energy supplementary feed (e.g., vegetable fat, molasses) should be provided to all pregnant animals in the flock. Administration of a propionic salt (sodium or calcium) is also beneficial (Brozos et al. 2011).

It is noteworthy that as clinical cases had already been diagnosed in the flock, the flock had already been characterised as an at risk flock, so individual animals need to be identified in order to prevent development of the disease, as well as for early instigation of treatment if clinical signs develop. For this, measurement of  $\beta$ -hydroxybutyrate blood concentration in ewes at the last month of gestation can be performed. If the number of foetuses borne by the ewe has not been identified, the value of 0.8 mmol L<sup>-1</sup> should be considered to indicate that the animal is at increased risk of developing clinical disease; if the number of foetuses had been determined, then the measurement should be applied only in animals carrying multiple foetuses and a threshold of 1.2 mmol L<sup>-1</sup> should be considered (Sargison 2007, Braun et al. 2010, Fthenakis et al. 2012). Animals found to have  $\beta$ -hydroxybutyrate blood concentration above those thresholds should be separated from other animals and monitored closely for development of early signs of the disease, in which case treatment should be instigated immediately. If the financial or labour constraints preclude the above approach, then examination of 10 to 15% of animals in the flock is of value



only in order to assess progressive changes in the  $\beta$ -hydroxybutyrate blood concentration on population basis.

Recent reports have indicated that the administration of butaphosphan and cyanocobalamin in animals of flocks with clinical cases of the disease may be of value to prevent further cases (Pereira et al. 2013, Temizel et al. 2015). Butaphosphan is an organic source of phosphorus that acts in gluconeogenesis in hepatic carbohydrate metabolism and plays a significant role in ATP synthesis (Rollin et al. 2010) and cyanocobalamin is the synthetic analogue of vitamin B<sub>12</sub>, which can be used as gluconeogenetic substrate (Pereira et al. 2013). Reports have indicated the efficacy of the combination in the prevention of ketosis in cows (Furll et al. 2010, Rollin et al. 2010), but its clinical application in sheep has been thusfar limited.

# **CHAPTER II**

## **RESEARCH WORK**

## A. BEFORE CHALLENGE OF THE MAMMARY GLAND OF THE EXPERIMENTAL EWES

### Materials and methods

#### Experimental overview

In total, 28 3- to 5-year old Chios-cross ewes were included into the study. Conditions prescribed by legislation of the European Union in relation to animal experimentation procedures (Council Directive 86/809/EEC) were met during this work. Throughout the study, ewes were housed.

Animals were included into the study in the summer, when they were at the end of their lactation period. Ewes with no history of mastitis were considered. Moreover, a general clinical examination, with special reference to their udder, was initially performed. Milk samples were collected for bacteriological and cytological examinations.

All animals into the study were initially drenched with a broad-spectrum anthelmintic, specifically netobimin (HAPADDEX® or. dr., Merck Animal Health, Summit, USA; dose rate: 20 mg kg<sup>-1</sup> bodyweight) 30 days before mating. Then, 21 days later, the ewes received orally 2 mL of phosphate-buffer-saline containing 5,000 third-stage larvae of a trichostrongylid helminth mixture of local strains of *Teladorsagia* spp., *Trichostrongylus* spp., *Cooperia* spp., *Haemonchus* spp. and *Oesophagostomum* spp., which had been kindly prepared by Dr E. Papadopoulos.

Four rams of known fertility were penned in a box adjoining to the one with the ewes for one month before their introduction into the females. They were introduced into ewes in early August, at which time sheep were in the reproductive season and the experimental ewes were into the dry-period. Animals were observed daily and matings were recorded; all ewes were mated within 20 days after ram introduction and no repeat matings were observed. For subsequent management of the experimental animals, first day of pregnancy (P0) was considered to be the day of their mating.

Ewes were subsequently monitored throughout their pregnancy. Repeated ultrasonographic examinations were carried out to confirm pregnancy and number of foetuses borne. Further, ultrasonographic examination of the mammary parenchyma was performed.

Blood samples were regularly collected from all ewes at regular time-points subsequently to the 100th day of pregnancy (P100), for measurement of  $\beta$ -hydroxybutyrate and glucose concentrations. Faecal samples were also collected for parasitological examinations.

Ewes with clinical signs relevant to pregnancy toxemia were recognised for treatment. All ewes lambed; their parturition (L0) was monitored. Possible obstetrical problems were recorded; details of the situation and outcome were noted. Further, possible post-partum genital disorders of the ewes were monitored.

### Clinical examination of the udder of the experimental ewes – Sample collection

Before admission into the study, all ewes were clinically examined, with special attention paid to their mammary glands and teats (Fthenakis 1994, Saratsis et al. 1998, Mavrogianni et al. 2005). In all cases, samples were collected from both teats and both mammary glands of each ewe.

A thorough disinfection was carried out by using povidone iodine scrub solution on the teat apex and the lower (1 cm) part of the teat skin. A fine (20 G), plastic, sterile catheter (Abbocath®; Abbott, Abbott Park, IL, USA) was used for sampling the teat duct and collecting teat duct material. The stylet was taken out and the catheter was cut with a sterile blade to a length of 2 mm. In order to ensure accurate and consistent cutting of the catheter at the desired length, a sterilised ruler was always placed beside the catheter. The whole procedure was carried out under aseptic conditions. The catheter was held from the cannula hub and was inserted into the teat, rolled around the internal teat wall, in order to sample the mucosa, and then withdrawn. Description and validation details of the method have been presented previously (Mavrogianni et al. 2006a). Milk samples were then obtained. The first two squirts of secretion were drawn onto the palm of the gloved hand of the investigator and examined for the presence of abnormal signs; then, 10 to 15 mL of secretion were carefully collected into a sterile container.

Subsequently, clinical examinations of the udder were performed on each occasion when ultrasonographic examination of the mammary parenchyma was carried out during the last stage of pregnancy.

## Bacteriological and cytological examinations of samples from the experimental ewes

Samples of material collected on the tip of the catheter ('teat duct material') and milk samples were plated onto Columbia 5% sheep blood agar; the media were incubated aerobically at 37 °C for up to 72 h. Throughout this study, all bacteria isolated were identified by using conventional techniques (Barrow and Feltham 1993, Euzeby 1997).

The California Mastitis Test (CMT) was carried out in milk samples, as described by Fthenakis (1995) for ewes' milk, by using a reagent (Jorgen Kruuse A/S, Marslev, Denmark). Five degrees of reaction scores ('negative', 'trace', '1', '2', '3'), were recognised, according to the standards of Schalm et al. (1971) and Fthenakis (1995) for ewes' milk. Finally, leucocyte subpopulations were identified by direct microscopy after Giemsa stain of milk films; in each case, 100 cells were observed and counted.

## Feeding of the experimental ewes

Starting 40 days before ram introduction and until the 60th day of pregnancy (P60), the ration provided per ewe was 0.65 kg of a commercial concentrate feed in mash form plus 1.25 kg of alfalfa hay, offered twice daily. Barley straw was provided into the pens thrice daily, whilst water was available *ad libitum*. The concentrate feed was based on cereal grains, cereal bran and oil by-products; details are in Table II.i. From P60 to P100, the ration provided per ewe was 0.30 kg of the same concentrate plus 1.00 kg of alfalfa hay, offered twice daily. Barley straw and water were also provided.

On P100, ewes were allocated into separate pens according to the number of foetuses borne (one, two or three), as indicated in the ultrasonographic examination. In ewes with one foetus, the ration provided per ewe was 0.25 kg of a specially formulated and prepared concentrate feed in mash form plus 0.25 kg of alfalfa hay, offered twice daily. Respective figures for ewes with two foetuses were 0.30 kg and 0.25 kg and for ewes with three foetuses were 0.40 kg and 0.25 kg. Barley straw was provided into the pens thrice daily, whilst water was available *ad libitum*. The concentrate feed was based on cereal grains, cereal bran and oil by-products; details are in Table II.i.

**Table II.i.** Details of the concentrate feed (in mash form) provided to ewes during the study.

	Concentrate feed provided until 100th day of pregnancy	Concentrate feed provided after 100th day of pregnancy
<b>Feed composition (g kg<sup>-1</sup>)</b>		
Maize grains	200.0	200.0
Barley grains	120.0	10.0
Wheat grains	40.0	21.5
Wheat bran	280.0	355.5
Cottoncake meal	120.0	160.0
Sunflower meal	100.0	121.0
Lucerne meal	60.0	52.0
Soybean meal	30.0	30.0
Soy oil	2.5	5.0
Limestone	32.5	30.0
Monocalcium phosphate	1.5	1.5
Magnesium oxide	1.0	1.0
Vitamin premix <sup>1</sup>	1.0	1.0
Trace mineral premix <sup>2</sup>	1.5	1.5
Salt	10.0	10.0
<b>Total</b>	<b>1000 g</b>	<b>1000 g</b>
<b>Feed proximate analysis<sup>3</sup> (g kg<sup>-1</sup>)</b>		
Dry matter	882.2	888.1
Crude protein (N×6.25)	142.2	174.2
Crude fat (ether extract)	53.3	46.3
Crude fibre	93.0	128.1
Ash	75.9	78.1
Starch	306.1	203.3
<b>Feed calculated analysis</b>		
Calcium (g kg <sup>-1</sup> )	11.9	11.5
Phosphorus (total) (g kg <sup>-1</sup> )	7.2	7.5
Sodium (g kg <sup>-1</sup> )	4.1	4.1
Chloride (g kg <sup>-1</sup> )	6.4	6.4
Lysine (g kg <sup>-1</sup> )	6.1	6.5
Methionine+cystine (g kg <sup>-1</sup> )	5.4	5.8
Net energy (FU <sub>L</sub> )	0.884	0.748

1 Supplied per kg of concentrate feed: retinol 10,000 IU, cholecalciferol 2.500 IU, tocopherol 30 IU.

2 Supplied per kg of concentrate feed: zinc 100 mg, manganese 100 mg, iron 80 mg, cobalt 0.2 mg, iodine 2 mg, selenium 0.2 mg.

3 According to Association of Official Analytical Chemists (1995).

Feed was provided in troughs allowing a space of 40 cm per ewe. Transition from the standard concentrate feed to the specially formulated feed was carried out as follows: for 3 days a mixture of 3:1 standard feed:special feed was given to ewes, for another 3 days a mixture of 1:1 standard feed:special feed was given and, finally, for another 3 days a mixture of 1:3 standard feed:special feed was given.

### Parasitological examinations of faecal samples from the experimental ewes

Faecal samples were initially collected on the day of challenge with the trichostrongylid helminth larvae mixture, 21 days later (9 days before ram introduction into the flock of ewes) and 28 days after ram introduction into the ewes. Then, samples were again collected on P100, P120, P140. Faecal samples were collected directly from the rectum of each animal, placed into an isothermic box and transferred to the laboratory for epg counting. Each sample was divided in three lots, as follows. One lot was processed for trichostrongylid epg counting according to the modified McMaster technique with saturated NaCl solution; the second lot was processed for *Dicrocoelium dendriticum* epg counting according to the modified McMaster technique with ZnSO<sub>4</sub> (sp.g. 1.40); finally, the third lot was processed for *Fasciola* spp. and *Paramphistomum cervi* epg counting by using the Telemann sedimentation technique (acid - ether) (Ministry of Agriculture, Fisheries and Food 1986, Rehbein et al. 1999, Otranto and Traversa 2002, Taylor 2010).

### Biochemical examinations of blood samples from the experimental ewes

Starting on P100 and every 5 days thereafter, a blood sample was collected from each ewe into the study, for measurement of  $\beta$ -hydroxybutyrate and glucose concentrations. On each occasion, samples were collected 4 to 5 hours after the morning feeding of the animals. A drop of blood was placed on an appropriate strip, which was subsequently inserted into an automated reader (Precision Xceed Meter<sup>®</sup>; Abbott Laboratories, Abbott Park, IL, USA), validated for measurement of  $\beta$ -hydroxybutyrate or glucose concentration in sheep blood (Panousis et al. 2012, Pichler et al. 2014). Different strips were used for measurements of  $\beta$ -hydroxybutyrate and glucose concentrations.

## Ultrasonographic examinations of the foetus(es) of the experimental ewes

Starting on P40, in the morning of each test day, a standard ultrasonographic examination of the uterus of each ewe was performed in order to confirm pregnancy, as well as to measure number of foetuses. The examination was repeated on P70, P100 and P120 to confirm continuation of pregnancy. After P120, animals were examined every five days until lambing (P125, P130, P135, P140, P145). On each of these days (P120-P145), the diameter of the abdomen of the foetus and a Doppler ultrasonographic examination of the umbilical artery were also performed.

The examination was performed with the animal on the standing position and restrained inside a crate, using the support of an assistant. For examination, hair on the abdomen and the udder had been fully clipped.

The standard procedure of ultrasonographic examination of ewes for diagnosis of pregnancy was performed with an ultrasound scanner (MyLab<sup>®</sup> 30; ESAOTE SpA, Genova, Italy) fitted with a convex transducer; 7.5 MHz imaging frequency and 120 mm scanning depth were used for this procedure. Coupling gel was applied. The transducer was placed on the skin of the abdomen, at a location depending on the day of gestation (more caudally at early gestation, moving cranially to the udder as gestation advanced). The diameter of the abdomen of the foetus was measured by addressing the ultrasound beam at 90 ° angle to the abdominal wall and parallel to the longitudinal axis of the foetus in the region of the umbilical cord (Ali and Hayder 2007).

Doppler measurements were taken at the umbilical artery; 2.5 MHz imaging frequency and 120 mm scanning depth were used for this procedure. A skilled assistant was helping when measuring the blood flow velocity, in order to make adjustments necessary for optimal quality of colour-flow images. Animals were stressed as little as possible during the examination, with the objective to remaining still for the period of time necessary to apply correctly the sample gate and to take the spectral display after localising the vessel to be studied. Each examination did not last over 30 to 40 s, after which the procedure was repeated only after 50 to 60 s. Each foetus was imaged in a cross section of its abdomen, at the point where the entrance of the umbilical artery therein, was seen. The umbilical artery was detected and visualised. At the start, a cross-section of the vessel, at the point of its entry into the foetus, was taken by using colour Doppler, with the objective to measure its diameter. Then, flow waveforms were obtained from the vessel at the midcord part of the free-floating umbilical cord; pulse wave Doppler was applied; the sample gate was positioned inside the vessel and at its centre, whilst taking care not to include the vessel wall



and to avoid artifacts. Adjustments necessary for optimal colour-flow image quality were made. A 'Doppler angle' [i.e., the angle at which the ultrasound beams intersect the path of flowing blood, also termed 'angle of insonation' (Ginther 2007, Petridis et al. 2014)] of 0 ° to 10 ° was used in the examination. No data were collected during foetal movements or transient cardiac arrhythmias.

Each examination session in an animal was completed within 10 min. In all cases, images (B-mode or Doppler) were frozen and saved on the equipment hard-disk for performing subsequently appropriate measurements and data analysis.

### **Ultrasonographic examinations of the udder of the experimental ewes**

Starting on the 120th day after the recorded mating (P120), in the morning of each test day, an ultrasonographic examination of the udder was performed. The examination was repeated every five days until lambing (P125, P130, P135, P140, P145).

The examination was performed with the animal on the standing position and restrained inside a crate, using the support of an assistant. For examination, hair on the abdomen and the udder had been fully clipped. Coupling gel was applied. Initially, the left side of the udder was imaged and subsequently, the whole procedure was repeated for the right side of the udder.

B-mode ultrasonographic examination of the mammary parenchyma was performed with an ultrasound scanner (MyLab® 30; ESAOTE SpA, Genova, Italy) fitted with a linear transducer. The transducer was placed on the caudal surface of the udder and moved around it; 10.0 MHz imaging frequency and 60 mm scanning depth were used for this procedure. The transducer was placed in a position perpendicular to the long axis and dorsal B-mode sections of the mammary parenchyma were taken, starting from the upper part downwards. In each mammary gland, three images were saved for further processing; first image was taken before the branching of the external pudental artery (*arteria pudenda externa*), second when distance between branches of the external pudental artery was ~1 cm and third image was taken immediately before the gland cistern (*sinus lactiferous*) became visible.

Doppler measurements were taken at the external pudental artery; 6.6 MHz imaging frequency and 60 mm scanning depth were used for this procedure. A skilled assistant was helping when measuring the blood flow velocity, in order to make adjustments necessary for optimal quality of colour-flow images. Animals were stressed as little as possible during the examination, with the objective to remaining still for the period of time necessary to apply correctly the sample gate and to take the spectral display after localising the vessel to be studied. The

external pudendal artery was detected and visualised. At the start, a cross-section of the vessel was taken before its branching, with the objective to measure its diameter. Then, flow waveforms were obtained from the vessel; pulse wave Doppler was applied; the sample gate was positioned inside the vessel and at its centre, whilst taking care not to include the vessel wall and to avoid artifacts. Adjustments necessary for optimal colour-flow image quality were made. A 'Doppler angle' of 50 ° to 60 ° was used in the examination.

Each examination session in an animal was completed within 20 min. In all cases, images (B-mode or Doppler) were frozen and saved on the equipment hard-disk for performing subsequently appropriate measurements and data analysis.

### **Weighing of lambs**

Lambs of the experimental ewes were weighed on the day of their birth by using automated scales (Jorgen Kruuse A/S, Marslev, Denmark).

### **Examination of the genital tract of ewes post-partum**

Immediately after lambing (day L0), the genital tract of each ewe was examined clinically with the animal in the standing position. The external genitalia were observed. Presence of vaginal discharge was evaluated. Two gloved fingers of the investigator were inserted into the vagina, in order to evaluate possible intra-vaginal presence of foetal membranes. The examination was performed daily for at least five days after parturition (animals with no problems) or until any clinical signs had subsided.

### **Milk yield measurements of the experimental ewes**

Milk yield of the ewes was estimated by measurements performed on the 3rd (L3) and the 5th (L5) days after lambing. Ewes were injected intramuscularly with 10 i.u. oxytocin and milked out. Lambs were taken away from the ewes for four hours, when ewes were injected again with 10 i.u. oxytocin and once more milked out. The milk was collected into a plastic container. To avoid misreading by foam formation and for accuracy of results, the milk was slowly poured into a volumetric glass container graduated to 2 mL and allowed still for five minutes before reading (Fthenakis and Jones 1990b, Fthenakis et al. 2005).

## Data management and analysis

### *Start of pregnancy of the experimental ewes*

Start of pregnancy for the experimental ewes was considered to be the day of their mating.

### *Retrospective grouping of experimental ewes depending on the number of fetuses borne*

Retrospectively and based on (i) results of ultrasonographic examination detecting the number of fetuses and (ii) number of lambs born, animals were grouped according to number of fetuses borne (one, two or three).

### *Retrospective allocation of experimental ewes into groups depending on the blood concentration of $\beta$ -hydroxybutyrate concentration*

Retrospectively and based on results of blood  $\beta$ -hydroxybutyrate concentrations, animals were allocated into one of two groups: group A included ewes with  $\beta$ -hydroxybutyrate concentrations  $>1.2 \text{ mmol L}^{-1}$  in two samples collected after the 129th day of pregnancy, whilst group B included all other ewes into the study.

### *Retrospective grouping of experimental ewes depending on the development of dystocia*

Specifically for analysis of the results of the Doppler ultrasonographic examination of the umbilical artery, animals were grouped according to development of dystocia, i.e., independently of number of fetuses borne and blood  $\beta$ -hydroxybutyrate concentrations.

### *Results taken into account during the ultrasonographic examination of the umbilical artery*

During the ultrasonographic examination of the foetus, four sets of images were recorded: (i) fetuses, (ii) images of cross-section of the foetus (iii) images of cross-sections of umbilical artery and (iv) spectral waveforms of umbilical artery. Data management was performed on images that had been saved at the time of examination.

Presence of fetuses and numbers borne by the ewe under examination were recorded. In stored images of cross-section of fetuses. The diameter of the foetus was calculated directly on the image as the distance from entrance of the umbilical cord into the foetus to the antipodal point of the circular section image.

Stored images of cross-sections of umbilical artery were processed by means of MyLab software (ESAOTE SpA, Genova, Italy), which, after pointing out the internal boundaries of the vessel, calculated the internal diameter of the vessel. Results were expressed as cm.

Spectral waveforms of the umbilical artery were processed by means of MyLab software (ESAOTE SpA, Genova, Italy). On each occasion, waveforms from three consecutive cardiac cycles of the animal under examination were considered for calculations. The software, based on the outline of the waveform, calculated directly the below haemodynamic parameters in that vessel (Maulik 2005, Ginther 2007, Wood et al. 2010).

- Resistance index:  $[(PSV-EDV)/PSV]$  (PSV: peak systolic velocity, EDV: end diastolic velocity) indicating the effect of the vessel under examination resisting blood flow.
- Pulsatility index:  $[(PSV-EDV)/TAMV]$  (TAMV: time-averaged maximum velocity) measuring the systolic-diastolic differential of the velocity pulse in the vessel under examination.
- Systolic:diastolic velocity ratio:  $[ASF/ADF]$  (ASF: average diastolic flow, ADF: average systolic flow) delineating systolic and diastolic phases of a blood flow waveform in the vessel under examination.
- General pressure:  $[P_{syst}-P_{diast}]$  ( $P_{syst}$ : systolic pressure,  $P_{diast}$ : diastolic pressure) measuring the change in pressure from the diastolic level to the systolic level in the vessel under examination (mm Hg).
- Mean pressure:  $[\frac{1}{3}\times P_{syst}+\frac{2}{3}\times P_{diast}]$  measuring the average blood pressure over time by proprietary pulse dynamics pattern-recognition algorithms in the vessel under examination (mm Hg).
- Mean velocity: indicating blood speed across the lumen of the vessel under examination at a given instance ( $m\ s^{-1}$ ).
- Systolic acceleration: indicating blood acceleration across the lumen of the vessel under examination ( $m\ s^{-2}$ ).
- Blood input: indicating the volume of blood entering into the foetus per unit of time ( $mL\ min^{-1}$ ).

*Results taken into account during the ultrasonographic examination of the udder of the experimental ewes*

During the ultrasonographic examination of the udder, three sets of images were recorded for each side of the udder: (i) images of mammary parenchyma, (ii) images of cross-sections of external pudendal artery and (iii) spectral waveforms of external pudendal artery. All images were initially evaluated visually for presence of abnormal structures therein. Then, data management was performed on images that had been saved at the time of examination. Data for left and right mammary gland of the same animal were initially considered separately.

Stored images of mammary parenchyma were processed by means of ImageJ software (National Institutes of Health, Rockville Pike, MD, USA), which can edit, process and analyse grey-scale images, by calculating area and pixel value statistics to produce intensity values (National Institutes of Health 2013). In an image processing context, grey-scale analysis refers to the image's overall pixel grey intensity values (Ojala et al. 2002). For analysis of grey-scale, intensity values of each of the three images stored from each mammary gland on each occasion were considered together. Areas with vessels or ductal formations were not taken into account for the grey-scale analysis. Results were expressed on a 0 (black) to 255 (white) scale.

Stored images of cross-sections of the external pudendal artery were processed by means of MyLab software (ESAOTE SpA, Genova, Italy), which, after pointing out the internal boundaries of the vessel, calculated the internal diameter of the vessel. Results were expressed as cm.

Spectral waveforms of the external pudendal artery were processed by means of MyLab software (ESAOTE SpA, Genova, Italy). On each occasion, waveforms from three consecutive cardiac cycles of the animal under examination were considered for calculations. The software, based on the outline of the waveform, calculated directly the following haemodynamic parameters in that vessel (Maulik 2005, Ginther 2007, Wood et al. 2010): resistance index, pulsatility index, systolic:diastolic velocity ratio, general pressure, mean pressure, mean velocity and systolic acceleration. Further, the below haemodynamic parameter was also calculated.

- Blood input: indicating the volume of blood entering into the mammary gland per unit of time ( $\text{mL min}^{-1}$ ).

#### *Duration of pregnancy*

The duration of pregnancy was calculated as the length of time from the day of start of pregnancy (P0) to the day of lambing (L0) and was expressed in days.

#### *Statistical computations*

All data were entered into Excel spreadsheets (Microsoft Corporation, Redmond, WA, USA). Initially, appropriate descriptive statistics for all parameters were performed.

In parameters, in which sequential measurements were taken, repeated measures mixed effect linear regression models were used to determine whether outcomes changed over the course of the study period. Models were adjusted for repeated measures within animals. Independent variables (fixed effects) included (i) number of fetuses borne (one, two or three) or inclusion into group A or B (based on results of blood  $\beta$ -hydroxybutyrate concentrations), (ii) day of pregnancy and (iii) a day of pregnancy by number of fetuses borne or group A/B interaction.

For ultrasonographic measurements in the foetus, repeated measures mixed effect linear regression models were used to determine whether outcomes in the two groups of the study changed over the course of the study period. Effect of experimental subjects (animals) was included as random effect in the model. Models were adjusted for repeated measures within animals. Independent variables (fixed effects) included experimental group (A or B), time of the study (i.e., P120, P125, P130, etc.) and a time of the study by experimental group interaction. In total two separate analyses were performed. Analysis of data obtained during the last stage of pregnancy (6 time-points: P120, P125, P130, P135, P140, P145) was initially performed, followed by analysis of data obtained during the last week of pregnancy (2 time-points: P140, P145).

For ultrasonographic measurements in the mammary parenchyma, at first, repeated measures mixed effect linear regression models were used to determine whether outcomes in the two mammary glands of each animal changed over the course of the study period. Models were adjusted for repeated measures within animals. Variables (fixed effects) included mammary gland (i.e., left or right) and time-point of the study (i.e., P120, P125, P130, etc.). In all cases, no significant differences were evident between the left and the right glands of the experimental animals within each group (for all parameters,  $P > 0.63$ ). In view of that, the results of all measurements for left and right glands obtained from each animal at each time-point were averaged for between group comparisons. Then, repeated measures mixed effect linear regression models were used to determine whether outcomes in the two groups of the study changed over the course of the study period. Effect of experimental subjects (animals) was included as random effect in the model. Models were adjusted for repeated measures within animals. Independent variables (fixed effects) included experimental group (A or B), time of the study (i.e., P120, P125, P130, etc.) and a time of the study by experimental group interaction. In total four separate analyses were performed. Analysis of data obtained during the last stage of pregnancy (6 time-points: P120, P125, P130, P135, P140, P145) was initially performed, followed by analysis of data obtained during the last fortnight of pregnancy (3 time-points: P135, P140, P145) and analysis of data obtained during the first week of lactation (2 time-points: L3, L5). Finally, an analysis that took into account all data (8 time-points) was carried out. Subsequently, a different set of analyses was performed, where, for each parent, results obtained on three time-points (P120, P140, L5) were compared between them.

The same model as for ultrasonographic measurements in the foetus was employed for evaluation of differences in blood  $\beta$ -hydroxybutyrate or glucose concentrations during pregnancy (P120-P145). For these two parameters, only analysis of all data obtained during the last stage of pregnancy (6 time-points: P120, P125, P130, P135, P140, P145) was performed.

The same model as for ultrasonographic measurements in the foetus was employed for parasitological findings during pregnancy (P100-P140). However, for this parameter only analysis of all data obtained during pregnancy (3 time-points: P100, P120, P140) was performed.

Durations of pregnancy, birth bodyweights of lambs born from ewes into the study and measurements of milk collected on each sampling occasion (L3, L5) were compared between groups A and B by using Student's t-test.

Analysis of correlation between blood  $\beta$ -hydroxybutyrate concentrations and faecal epg counts / blood glucose concentrations / lamb birth bodyweight and between faecal epg counts and lamb birth bodyweight was performed.

For evaluation of potential correlations between milk quantities obtained from ewes and grey-scale intensity / blood input values, results of milk quantities collected on each of L3 or L5 were correlated with the grey-scale / blood input values obtained from each ewe on the respective occasion, i.e., 32 pairs of values for group A results and 24 pairs of values for group B results. Then, averaged results of milk quantities obtained on L3 and L5 were correlated with the grey-scale / blood input values obtained from each ewe (mammary parenchyma) on each of P130 to P145, i.e., 28 pairs in total tested on four separate occasions.

Incidence of animals with dystocia or with post-partum genital disorders among ewes in group A and B was calculated. Moreover, incidence of animals in which perinatal mortality of offspring (i.e., foetus(es) dead at lambing or newborn lamb dead by 2nd day of life) was noted, among ewes in group A and B was calculated. In all cases, respective confidence intervals were also computed. Initially, Fisher-exact tests were performed for comparisons of proportions between the two groups; then, z ratio for the significance of difference of two proportions was determined.

An electronic data management tool was employed (Lowry 2012, 2015). Significance level was set at  $P \leq 0.05$ .

## **Results**

### **Findings before enrolling the experimental ewes into the study**

All ewes that were introduced into the study had no history of mastitis. No mammary abnormalities were clinically evident in any of the experimental ewes during clinical udder examination before enrolment into the study. Moreover, no bacteria were isolated from any teat

duct material or milk samples collected before start of the study. California Mastitis Test scores in milk samples were negative (scores 'negative' or 'trace') and observation of Giemsa-stained milk films revealed only scarce presence of macrophages therein (on average, one cell per 10 fields with the 10× objective lens).

### Findings of pregnancy diagnosis in the experimental ewes

All ewes into the study conceived. During the ultrasonographic examination, of them, 7 were found to bear one foetus, 17 were found to bear two foetuses and 4 were found to bear three foetuses. These findings were confirmed at parturition when ewes lambed the number of offspring imaged during ultrasonographic examination for pregnancy diagnosis.

### Blood concentrations of $\beta$ -hydroxybutyrate in the experimental ewes

Ewes with three foetuses had greater median blood  $\beta$ -hydroxybutyrate concentrations than ewes with two or one foetuses. In ewes with one foetus, there was no evidence that changes in blood  $\beta$ -hydroxybutyrate concentrations during last stage of pregnancy were significant ( $P=0.52$ ); however, in ewes with two or three foetuses, progressive increase in those concentrations was significant ( $P<0.005$ ).

In total, 16 ewes were allocated into group A, i.e. 57% of the ewes that received reduced energy feed. These included 2 ewes with one foetus (28% of ewes bearing singles), 10 ewes with two foetuses (59% of ewes bearing twins) and 4 ewes with three foetuses (100% of ewes bearing triplets). Of these 16 ewes, 4 (25%; i.e., 14% of all ewes into the study) showed clinical signs relevant to pregnancy toxaemia. Group B included 12 ewes (5 with one foetus, i.e., 72% of ewes bearing singles, and 7 with two foetuses, i.e. 41% of ewes bearing twins).

Ewes into group A had greater blood  $\beta$ -hydroxybutyrate concentrations than ewes into group B ( $P<0.001$ ). Progressive increase of blood  $\beta$ -hydroxybutyrate concentrations in ewes of group A was significant ( $P<0.001$ ), whilst there was no evidence of significant changes in blood  $\beta$ -hydroxybutyrate concentrations in ewes of group B ( $P=0.855$ ). Finally, there was a significant interaction of day of pregnancy and group ( $P<0.001$ ). Detailed results of blood  $\beta$ -hydroxybutyrate concentrations are in Table II.ii.



**Table II.ii.** Median (min.-max.) blood  $\beta$ -hydroxybutyrate concentrations (mmol L<sup>-1</sup>) in ewes, which had been receiving reduced feed energy during the final stage of pregnancy.

	Day of pregnancy									
	100th	105th	110th	115th	120th	125th	130th	135th	140th	145th
Ewes with 1 foetus	0.50 (0.5-0.7)	0.45 (0.4-0.8)	0.60 (0.4-0.7)	0.60 (0.4-1.5)	0.60 (0.4-0.7)	0.70 (0.5-2.3)	0.80 (0.5-2.1)	0.70 (0.4-2.4)	0.70 (0.5-2.8)	0.60 (0.4-2.4)
Ewes with 2 foetuses	0.70 (0.4-1.5)	0.80 (0.4-2.3)	1.00 (0.5-2.5)	0.90 (0.4-1.6)	1.00 (0.4-1.4)	1.10 (0.5-3.4)	0.80 (0.4-2.4)	1.70 (0.5-3.3)	2.20 (0.4-4.8)	2.50 (0.8-6.2)
Ewes with 3 foetuses	0.70 (0.5-0.9)	0.70 (0.5-0.9)	1.40 (1.0-1.4)	1.50 (1.1-2.1)	2.30 (2.2-2.8)	3.15 (2.3-3.6)	3.25 (1.7-3.9)	2.60 (2.1-6.8)	4.40 (1.8-6.3)	4.55 (2.0-6.1)
Ewes into group A	0.80 (0.5-1.5)	0.90 <sup>a</sup> (0.4-2.3)	1.30 <sup>a</sup> (0.4-2.5)	1.10 <sup>a</sup> (0.4-2.1)	1.25 <sup>a</sup> (0.5-2.8)	1.90 <sup>a</sup> (0.5-3.6)	2.00 <sup>a</sup> (0.5-3.9)	2.40 <sup>a</sup> (1.2-6.8)	2.80 <sup>a</sup> (1.8-6.3)	3.70 <sup>a</sup> (1.7-6.2)
Ewes into group B	0.60 (0.4-1.4)	0.60 <sup>a</sup> (0.4-0.8)	0.65 <sup>a</sup> (0.5-0.9)	0.65 <sup>a</sup> (0.4-1.5)	0.60 <sup>a</sup> (0.4-1.2)	0.70 <sup>a</sup> (0.5-1.4)	0.60 <sup>a</sup> (0.4-1.1)	0.70 <sup>a</sup> (0.4-1.4)	0.60 <sup>a</sup> (0.4-1.1)	0.80 <sup>a</sup> (0.4-1.1)

Group A: ewes with  $\beta$ -hydroxybutyrate concentrations >1.2 mmol L<sup>-1</sup> in two samples collected after the 129th day of pregnancy - Group B: all other ewes into the study.

<sup>a</sup> within the same column:  $P < 0.05$ .

## Parasitological findings in the experimental ewes

Before start of the mating period, faecal counts for nematodes and trematodes in all animals were 0 epg. Ewes allocated into group A had more increased nematode epg counts towards the end of pregnancy than ewes allocated into group B ( $P<0.025$ ). Progressive increase of nematode epg counts in faecal samples throughout pregnancy was significant ( $P<0.001$ ), whilst there was a significant interaction of day of pregnancy and group ( $P<0.001$ ). Detailed results of nematode epg counts in faecal samples are in Table II.iii. In contrast, faecal counts for trematodes in all animals were consistently 0 epg throughout the study.

There was a significant correlation between blood  $\beta$ -hydroxybutyrate concentrations and faecal epg counts ( $r=0.556$ ,  $P<0.001$ ) during the last stage of pregnancy.

**Table II.iii.** Median (min.-max.) nematode epg counts in faecal samples from ewes, which had been receiving reduced feed energy during the final stage of pregnancy.

	on day of challenge	21 days after challenge	28 days after ram introduction	Day of pregnancy		
				100th	120th	140th
Ewes with 1 foetus	0 (0-0)	0 (0-50)	250 (100-400)	350 (150-500)	550 (500-700)	650 (600-850)
Ewes with 2 foetuses	0 (0-0)	0 (0-50)	250 (150-400)	35 (200-450)	600 (350-850)	800 (500-1100)
Ewes with 3 foetuses	0 (0-0)	0 (0-50)	250 (150-450)	375 (250-550)	700 (500-850)	975 (800-1250)
Ewes into group A	0 (0-0)	0 (0-50)	250 (150-450)	350 (200-550)	650 (450-850)	850 <sup>a</sup> (500-1250)
Ewes into group B	0 (0-0)	0 (0-50)	250 (100-400)	350 (150-500)	550 (350-800)	675 <sup>a</sup> (500-950)

Group A: ewes with  $\beta$ -hydroxybutyrate concentrations  $>1.2$  mmol L<sup>-1</sup> in two samples collected after the 129th day of pregnancy - Group B: all other ewes into the study.

<sup>a</sup> within the same column:  $P<0.05$ .

## Findings of ultrasonographic examination of the diametre of the abdomen of the foetus and findings of Doppler ultrasonographic examination of the umbilical artery

There was clear evidence that, throughout the last stage of pregnancy, the progressive increase in the diametre of the foetus and in the blood input in the umbilical artery was significant ( $P<0.005$  and  $P<0.001$ , respectively) in both groups, A or B (Fig. II.1.). Progressive significant changes were evident during the last 10 days of the pregnancy in three haemodynamic

parameters (resistance index, pulsatility index, systolic:diastolic velocity ratio;  $P=0.002$ ,  $P=0.019$ ,  $P<0.001$ , respectively) only in group A ewes, whilst in group B no significant modifications were recorded ( $P>0.25$ ) (Fig. II.2.). In umbilical artery diameter, progressive significant increase during the last 10 days of the gestation was evident in both groups, A or B ( $P=0.025$ ,  $P<0.001$ , respectively) (Figs II.2. and II.3.). No significant changes were seen in the other haemodynamic parameters throughout the study in either group ( $P<0.07$ ).


In general, there were no differences between ultrasonographic findings in ewes of the two groups (A, B) when comparisons took place on a day by day basis, except for the blood input into the umbilical artery (Fig. II.1.); in that parameter, values in ewes of group B were significantly greater than values in group A animals from 120th day of gestation. However, there was clear evidence that when ewes were allocated into ones which subsequently developed and ones that did not develop dystocia, there were significant differences ( $P<0.047$ ) between them in all haemodynamic parameters in the last two measurements during pregnancy (P140, P145).

Further, when data obtained during the last two measurements (140th and 145th day of pregnancy) were considered together and compared (i.e., 32 observations for group A ewes and 24 observations for group B ewes), a difference between group A and B became clear in the results of umbilical artery diameter, resistance index, pulsatility index and systolic:diastolic velocity ratio, all being greater in ewes into group B than in ewes into group A ( $P<0.001$ ,  $P=0.040$ ,  $P=0.029$ ,  $P=0.007$ , respectively). For ewes distinguished into ones which subsequently developed and ones that did not develop dystocia, again differences in all haemodynamic parameters were significant ( $P<0.02$ ).

Detailed results of ultrasonographic examinations are in Tables II.iv and II.v.


**Figure II.1.** Time-series (last stage of pregnancy) graphs of (a) the diametre of the abdomen of the foetus (cm) and (b) the blood input into the umbilical artery ( $\text{mL min}^{-1}$ ) in ewes with (group A, red line) or without (group B, blue line) increased  $\beta$ -hydroxybuturate blood concentration during the last stage of pregnancy.

(a)




**Figure II.1.** (continued).

(b)




**Figure II.2.** Time-series (last stage of pregnancy) graphs of (a) the diameter (cm) of the umbilical artery and (b) the systolic:diastolic velocity ratio of the umbilical artery in ewes with (group A, red line) or without (group B, blue line) increased  $\beta$ -hydroxybuturate blood concentration during the last stage of pregnancy.


(a)



(b)



**Figure II.3.** Spectral waveforms (140th day of pregnancy) of the umbilical artery (at the point of entrance of the artery into the abdomen of the foetus) of ewes, with (group A, left picture) or without (group B, right picture) increased  $\beta$ -hydroxybuturate blood concentration during the last stage of pregnancy. Images taken and processed on a MyLab<sup>®</sup> 30 ultrasonography system with convex transducer, imaging frequency: 2.5 MHz - scanning depth: 120 mm.



**Table II.iv.** Quantitative results (mean±standard error of the mean) of ultrasonographic examination of the foetus in ewes, which had been receiving reduced feed energy during the final stage of pregnancy.

	Day of pregnancy					
	120th	125th	130th	135th	140th	145th
<b>Group A</b>						
Diametre of foetus (cm)	6.77±0.18	7.01±0.20	7.27±0.19	7.52±0.22	7.69±0.20	8.03±0.21
Umbilical artery diametre (cm)	0.63±0.02	0.64±0.03	0.65±0.02	0.66±0.01	0.65±0.02**	0.71±0.01*
Resistance index	0.71±0.01	0.66±0.02	0.68±0.01	0.65±0.02	0.62±0.02	0.46±0.05
Pulsatility index	1.24±0.04	1.16±0.07	1.23±0.04	1.25±0.07	0.97±0.05*	0.75±0.08
Systolic:diastolic velocity ratio	3.06±0.41	2.68±0.30	3.98±0.32	2.56±0.28	2.26±0.30	1.40±0.31
General pressure (mm Hg)	1.16±0.14	0.84±0.11**	1.06±0.11**	1.18±0.09	1.91±0.47	1.36±0.23
Mean pressure (mm Hg)	0.53±0.05	0.45±0.04	0.48±0.04	0.52±0.04	0.99±0.20	0.41±0.05
Mean velocity (m s <sup>-1</sup> )	0.32±0.02	0.32±0.02**	0.31±0.01**	0.33±0.01	0.44±0.05	0.28±0.00
Systolic acceleration (m s <sup>-2</sup> )	4.21±0.15	3.72±0.29	3.37±0.15	3.90±0.25	3.82±0.35	2.25±0.36
Blood input (mL min <sup>-1</sup> )	534.8±34.1**	639.7±34.2	602.2±30.2*	679.8±44.5*	658.5±40.9**	861.0±41.2*
<b>Group B</b>						
Diametre of foetus (cm)	6.67±0.24	6.90±0.30	7.16±0.30	7.74±0.34	7.96±0.38	8.33±0.33
Umbilical artery diametre (cm)	0.66±0.02	0.71±0.03	0.70±0.03	0.72±0.03	0.79±0.02**	0.80±0.03*
Resistance index	0.68±0.01	0.66±0.02	0.68±0.01	0.68±0.03	0.69±0.02	0.62±0.02
Pulsatility index	1.15±0.03	1.06±0.04	1.18±0.04	1.22±0.12	1.19±0.05*	0.99±0.07
Systolic:diastolic velocity ratio	3.28±0.11	3.01±0.17	3.17±0.08	3.25±0.33	3.33±0.20	2.78±0.22
General pressure (mm Hg)	1.15±0.15	1.40±0.08**	1.60±0.13**	2.15±0.37	2.08±0.27	2.56±0.73
Mean pressure (mm Hg)	0.51±0.06	0.60±0.03	0.65±0.07	0.90±0.19	0.85±0.16	1.22±0.33
Mean velocity (m s <sup>-1</sup> )	0.31±0.02	0.36±0.01**	0.36±0.02**	0.41±0.05	0.42±0.04	0.49±0.06
Systolic acceleration (m s <sup>-2</sup> )	3.89±0.23	4.06±0.32	4.02±0.24	4.90±0.51	4.46±0.27	4.15±0.61
Blood input (mL min <sup>-1</sup> )	726.3±53.1**	755.4±79.2	891.2±89.9*	859.7±60.4*	1222.1±133.6**	1210.6±136.0*



**Table II.iv.** (continued).

	Day of pregnancy					
	120th	125th	130th	135th	140th	145th
<b>Ewes which subsequently developed dystocia</b>						
Diametre of foetus (cm)	6.87±0.24	7.13±0.19	7.57±0.21	7.87±0.30	8.00±0.22	7.81±0.19
Umbilical artery diametre (cm)	0.65±0.02	0.66±0.03	0.65±0.02	0.65±0.01	0.63±0.02*	0.70±0.02*
Resistance index	0.70±0.01	0.63±0.03	0.66±0.01	0.66±0.03	0.55±0.01**	0.30±0.03**
Pulsatility index	1.22±0.04	1.02±0.07	1.21±0.08	1.30±0.11	0.89±0.01**	0.48±0.01**
Systolic:diastolic velocity ratio	3.36±0.15	2.84±0.21	3.34±0.28	3.04±0.18	2.51±0.12*	1.56±0.01**
General pressure (mm Hg)	1.27±0.15	1.00±0.14	1.24±0.10	1.34±0.07	1.10±0.12**	0.53±0.01**
Mean pressure (mm Hg)	0.52±0.08	0.46±0.04	0.53±0.05	0.50±0.02	0.47±0.05*	0.27±0.02*
Mean velocity (m s <sup>-1</sup> )	0.32±0.03	0.32±0.02	0.31±0.02	0.33±0.01	0.31±0.01*	0.28±0.01*
Systolic acceleration (m s <sup>-2</sup> )	4.25±0.05	3.64±0.47	3.53±0.23	3.82±0.37	2.95±0.18**	1.27±0.03**
Blood input (mL min <sup>-1</sup> )	540.7±43.2	670.4±47.0	623.7±38.5	656.2±51.5	678.0±70.5*	822.3±55.6*
<b>Ewes which subsequently did not develop dystocia</b>						
Diametre of foetus (cm)	6.66±0.18	6.89±0.22	7.04±0.21	7.44±0.23	7.56±0.26	8.35±0.26
Umbilical artery diametre (cm)	0.63±0.02	0.68±0.03	0.69±0.02	0.70±0.02	0.75±0.02*	0.77±0.02*
Resistance index	0.70±0.01	0.67±0.01	0.69±0.01	0.66±0.02	0.68±0.02**	0.62±0.02**
Pulsatility index	1.19±0.04	1.21±0.06	1.21±0.03	1.21±0.08	1.14±0.05**	1.00±0.05**
Systolic:diastolic velocity ratio	3.55±0.18	3.14±0.12	3.29±0.08	3.19±0.21	3.19±0.16*	2.75±0.15**
General pressure (mm Hg)	1.21±0.11	1.30±0.10	1.37±0.11	1.81±0.24	2.52±0.37**	2.17±0.50**
Mean pressure (mm Hg)	0.52±0.04	0.51±0.04	0.55±0.05	0.74±0.12	1.06±0.17*	1.03±0.23*
Mean velocity (m s <sup>-1</sup> )	0.32±0.02	0.34±0.02	0.34±0.02	0.38±0.03	0.46±0.04*	0.45±0.05*
Systolic acceleration (m s <sup>-2</sup> )	3.98±0.19	3.93±0.22	3.64±0.20	4.48±0.34	4.47±0.26**	3.89±0.43**
Blood input (mL min <sup>-1</sup> )	655.0±45.1	708.5±58.2	765.6±67.6	778.2±49.4	996.3±106.3*	1089.8±94.4*

**Table II.iv.** (continued).

Group A: ewes with  $\beta$ -hydroxybutyrate concentrations  $>1.2 \text{ mmol L}^{-1}$  in two samples collected after the 129th day of pregnancy - Group B: all other ewes into the study.

\*  $P < 0.05$ , \*\*  $P < 0.01$  between the two groups at respective day for the respective parameter.

**Table II.v.** Quantitative results (mean±standard error of the mean) of Doppler ultrasonographic examination of the umbilical artery in ewes, which had been receiving reduced feed energy during the final stage of pregnancy; combined results of 140th and 145th day of pregnancy.

Haemodynamic parametre	Group A	Group B
Umbilical artery diametre (cm)	0.68±0.01*	0.79±0.02*
Resistance index	0.56±0.03*	0.66±0.02*
Pulsatility index	0.87±0.05*	1.10±0.05*
Systolic:diastolic velocity ratio	2.00±0.22*	3.08±0.15*
General pressure (mm Hg)	1.74±0.29	2.30±0.36
Mean pressure (mm Hg)	0.78±0.12	1.02±0.17
Mean velocity (m s <sup>-1</sup> )	0.39±0.03	0.45±0.04
Systolic acceleration (m s <sup>-2</sup> )	3.25±0.28	4.30±0.31
Blood input (mL min <sup>-1</sup> )	755.3±33.7*	1215.7±92.5*

Haemodynamic parametre	Ewes which subsequently developed dystocia	Ewes which subsequently did not develop dystocia
Umbilical artery diametre (cm)	0.66±0.03*	0.76±0.01*
Resistance index	0.45±0.07*	0.66±0.01*
Pulsatility index	0.68±0.10*	1.08±0.04*
Systolic:diastolic velocity ratio	2.13±0.26*	3.01±0.11*
General pressure (mm Hg)	0.87±0.18*	2.38±0.29*
Mean pressure (mm Hg)	0.39±0.07*	1.05±0.13*
Mean velocity (m s <sup>-1</sup> )	0.30±0.02*	0.46±0.03*
Systolic acceleration (m s <sup>-2</sup> )	2.28±0.45*	4.23±0.24*
Blood input (mL min <sup>-1</sup> )	735.7±88.8*	1047.9±69.8*

Group A: ewes with  $\beta$ -hydroxybutyrate concentrations  $>1.2$  mmol L<sup>-1</sup> in two samples collected after the 129th day of pregnancy - Group B: all other ewes into the study.

\* between results in the same row:  $P<0.05$ .

## Blood concentrations of glucose in the experimental ewes

Ewes with three foetuses had smaller median blood concentrations of glucose than ewes with two or one foetuses ( $P=0.048$  for comparisons between ewes with three foetuses and ewes with one or two foetuses).

After P130, of the 16 ewes in group A, 10 had blood glucose concentration  $<55$  mg dL<sup>-1</sup> and 5 ewes had blood glucose concentration  $<35$  mg dL<sup>-1</sup> in at least one measurement. During the same period, all group B ewes had blood glucose concentration  $>55$  mg dL<sup>-1</sup> in at least one measurement. After P130, there was evidence that ewes into group A had smaller blood glucose concentrations than ewes into group B ( $P=0.033$ ). Detailed results of blood glucose concentrations are in Table II.vi.

Correlation coefficients ( $r$ ) between blood concentrations of  $\beta$ -hydroxybutyrate and glucose in ewes with one, two or three foetuses were -0.669, -0.519 and -0.811, respectively ( $P<0.003$ ).

Correlation coefficients ( $r$ ) between blood concentrations of  $\beta$ -hydroxybutyrate and glucose in ewes in group A or B were -0.469 ( $P < 0.001$ ) and 0.046 ( $P = 0.772$ ), respectively.

**Table II.vi.** Median (min.-max.) blood glucose concentrations (mmol L<sup>-1</sup>) in ewes, which had been receiving reduced feed energy during the final stage of pregnancy.

	Day of pregnancy									
	100th	105th	110th	115th	120th	125th	130th	135th	140th	145th
Ewes with 1 foetus	58 (57-60)	60 (56-61)	62 (57-65)	48 (44-66)	48.5 (40-71)	58 (44-71)	49 (34-65)	60 (27-65)	60 (30-88)	68 (30-93)
Ewes with 2 foetuses	65 (64-65)	69 (49-75)	63 (44-93)	62 (40-88)	59 (45-77)	57 (44-73)	58 (42-78)	65 (32-92)	64 (36-86)	62 (25-106)
Ewes with 3 foetuses	60 (55-65)	55 (53-60)	63.5 (62-67)	58 (45-63)	46 (37-52)	39.5 (33-50)	45 (27-57)	51 (20-54)	39 (28-60)	36 (28-58)
Ewes into group A	63 (55-65)	55 (49-75)	64 (60-67)	56 (43-81)	54 (37-68)	53.5 (33-68)	48 <sup>a</sup> (27-74)	52.5 <sup>a</sup> (20-85)	52 <sup>a</sup> (28-86)	53.5 <sup>a</sup> (25-80)
Ewes into group B	59 (57-65)	60 (51-69)	62 (44-93)	66 (40-88)	59 (40-77)	58 (44-73)	61 <sup>a</sup> (46-78)	65 <sup>a</sup> (41-92)	63 <sup>a</sup> (43-88)	71 <sup>a</sup> (57-106)

Group A: ewes with  $\beta$ -hydroxybutyrate concentrations >1.2 mmol L<sup>-1</sup> in two samples collected after the 129th day of pregnancy - Group B: all other ewes into the study.

<sup>a</sup> within the same column:  $P < 0.05$ .

## Findings of clinical examination of the udder of the experimental ewes

No mammary abnormalities were clinically evident in any of the experimental ewes at any of the periodic clinical examination during the last stage of pregnancy.

## Findings of ultrasonographic examination of the udder of the experimental ewes

Anatomic structures within the mammary parenchyma had a reduced to medium echogenicity. Blood vessels and lactiferous ducts could be readily imaged in all the animals during the study. No abnormal structures were evident. As pregnancy advanced (nearing parturition), the mammary parenchyma was seen more heterogeneous, with simultaneous appearance of a lobular image (consisting of the alveolar areas with reduced echogenicity) and the ductal part of the gland and the blood vessels therein (imaged as anechoic antra).

Throughout the last stage of pregnancy, mean ( $\pm$ standard error of the mean) grey-scale intensity values in group A were  $69.5\pm 4.5$  on P120,  $81.9\pm 6.4$  on P130,  $95.3\pm 3.3$  on P145; respective values in group B were  $64.5\pm 4.9$ ,  $71.3\pm 4.1$  and  $79.3\pm 2.0$  (for differences between groups,  $P=0.007$ ). There was also clear evidence that, throughout the last stage of pregnancy, the progressive increase in grey-scale intensity values was significant ( $P<0.001$ ) in both groups, A or B. Then, after lambing, on L3, grey-scale intensity values decreased sharply compared to those on P145: by 20% or 15% for group A or B, respectively, which was a significant decrease ( $P=0.003$  for group A,  $P=0.005$  for group B). During the first week of lactation, changes over time were not significant ( $P>0.6$ ), although differences between the two groups remained significant ( $P=0.046$ ) (Figs II.4. and II.5.).

**Figure II.4.** Serial B-mode ultrasonographic appearance of mammary parenchyma of ewes with (group A, left picture in each pair of images) or without (group B, right picture in each pair of images) increased  $\beta$ -hydroxybuturate blood concentration during the last stage of pregnancy: (a) on the 120th/125th day of pregnancy (P120/P125), (b) on the 145th day of pregnancy (P145), (c) on the 5th day of the subsequent lactation period (L5). Images taken at a level after the branching of the external pudendal artery (when distance between the two branches was  $\sim 0.5$  cm) on a MyLab<sup>®</sup> 30 ultrasonography system with linear transducer, imaging frequency: 10.0 MHz - scanning depth: 60 mm.

(a)



(b)





Figure II.4. (continued).

(c)




Figure II.5. Time-series (last stage of pregnancy and initial stage of lactation period) graphs of the grey-scale intensity values (0-255) of the mammary gland of ewes with (group A, red line) or without (group B, blue line) increased  $\beta$ -hydroxybuturate blood concentration during the last stage of pregnancy.






Throughout the last stage of pregnancy and during the first week of lactation, the progressive increase of the diameter of the external pudendal artery was significant in both groups, A or B ( $P<0.001$ ). However, the diameter of the vessel did not differ significantly between the two groups ( $P=0.374$  in pregnancy and  $P=0.591$  in lactation) (Fig. II.6.).

**Figure II.6.** Time-series (last stage of pregnancy and initial stage of lactation period) graphs of the diameter (cm) of the external pudendal artery of ewes with (group A, red line) or without (group B, blue line) increased  $\beta$ -hydroxybuturate blood concentration during the last stage of pregnancy.



The spectral display in Doppler mode was observed as a broad band structure (Fig. II.7.). Throughout the last stage of pregnancy, the progressive changes in the haemodynamic parameters studied were significant ( $P<0.02$ ) in both groups, A or B, bar changes in resistance index or in blood systolic acceleration ( $P>0.4$ ). During the first week of lactation, changes in all the haemodynamic parameters studied were not significant in both groups ( $P>0.12$ ), bar changes in blood input in group B ewes, which decreased significantly from L3 to L5 ( $P=0.009$ ).

**Figure II.7.** Spectral waveforms (145th day of pregnancy) of the external pudendal artery of ewes with (group A, left picture) or without (group B, right picture) increased  $\beta$ -hydroxybuturate blood concentration during the last stage of pregnancy. Images taken and processed on a MyLab® 30 ultrasonography system with linear transducer, imaging frequency: 6.6 MHz - scanning depth: 60 mm.




Blood input was significantly greater in ewes of group B ( $169 \text{ mL min}^{-1}$  on P130,  $397 \text{ mL min}^{-1}$  on P145,  $649 \text{ mL min}^{-1}$  on L3) than in ewes of group A ( $119 \text{ mL min}^{-1}$ ,  $262 \text{ mL min}^{-1}$ ,  $413 \text{ mL min}^{-1}$  on respective days) during the last stage of pregnancy and the first week of lactation ( $P < 0.05$ ) (Fig. II.8.). Differences between group A and group B in all other haemodynamic parameters studied were not significant throughout the last stage of pregnancy ( $P > 0.25$ ). However, when data obtained only during the last fortnight of pregnancy or during the first week of lactation were considered, differences in pulsatility index (Fig. II.8.) and in mean blood velocity showed evidence of significance between the two groups ( $P = 0.007$  and  $P = 0.036$ , respectively). Differences in results for these two parameters were also significant during the first week of lactation ( $P < 0.04$ ), whilst for other parameters, they were not ( $P > 0.06$ ).

Detailed results of ultrasonographic examinations are in Table II.vii.


**Figure II.8.** Time-series (last stage of pregnancy and initial stage of lactation period) graphs of the Doppler ultrasonography results of the external pudendal artery of ewes with (group A, red line) or without (group B, blue line) increased  $\beta$ -hydroxybuturate blood concentration during the last stage of pregnancy and initial stage of lactation (a) blood input ( $\text{mL min}^{-1}$ ), (b) pulsatility index.

(a)



**Figure II.8.** (continued).

(b)



**Table II.vii.** Quantitative results (mean±standard error of the mean) of ultrasonographic (B-mode or Doppler) examination of the udder in ewes, which had been receiving reduced feed energy during the final stage of pregnancy; results of both mammary glands of ewes are considered together, as no statistical difference ( $P>0.63$ ) was evident between left and right mammary glands.

	Day of pregnancy						Day of lactation	
	120th	125th	130th	135th	140th	145th	3rd	5th
Group A								
Mammary parenchyma grey-scale	69.5±4.5	76.0±1.9*	81.9±6.4*	82.1±5.5*	89.9±4.7 <sup>a</sup>	95.3±3.3 <sup>**</sup>	76.7±4.5*	75.5±5.0 <sup>a</sup>
External pudendal artery diametre (cm)	0.35±0.02 <sup>ab</sup>	0.38±0.02	0.38±0.02	0.39±0.02	0.44±0.02 <sup>ac</sup>	0.46±0.02	0.53±0.02	0.56±0.04 <sup>bc</sup>
Resistance index	0.62±0.02 <sup>ab</sup>	0.60±0.01	0.60±0.02	0.62±0.01	0.56±0.02 <sup>a</sup>	0.58±0.01	0.59±0.00	0.58±0.01 <sup>b</sup>
Pulsatility index	1.31±0.04 <sup>ab</sup>	1.03±0.03	1.02±0.05	1.08±0.05	0.88±0.04 <sup>**ac</sup>	0.95±0.04 <sup>*</sup>	0.99±0.03 <sup>*</sup>	0.98±0.04 <sup>**bc</sup>
Systolic:diastolic velocity ratio	2.98±0.13 <sup>ab</sup>	2.74±0.14	2.61±0.13±	2.68±0.10	2.44±0.14 <sup>a</sup>	2.86±0.14	2.48±0.05	2.42±0.07 <sup>b</sup>
General pressure (mm Hg)	0.94±0.06 <sup>ab</sup>	1.17±0.14	1.48±0.16	1.37±0.20	1.73±0.21 <sup>ac</sup>	1.55±0.09	4.45±0.39	4.49±0.37 <sup>bc</sup>
Mean pressure (mm Hg)	0.32±0.02 <sup>ab</sup>	0.43±0.05	0.56±0.07	0.60±0.10	0.86±0.10 <sup>ac</sup>	0.61±0.05	1.86±0.20	1.86±0.16 <sup>bc</sup>
Mean velocity (m s <sup>-1</sup> )	0.24±0.01 <sup>ab</sup>	0.35±0.04	0.33±0.02	0.34±0.03	0.44±0.02 <sup>*ac</sup>	0.48±0.02 <sup>*</sup>	0.62±0.04 <sup>*</sup>	0.62±0.03 <sup>bc</sup>
Systolic acceleration (m s <sup>-2</sup> )	5.17±0.28 <sup>a</sup>	5.31±0.53	5.76±0.65	5.78±0.47	5.83±0.40 <sup>b</sup>	6.10±0.20	9.51±1.04	9.50±0.83 <sup>ab</sup>
Blood input (mL min <sup>-1</sup> )	83.0±10.9 <sup>ab</sup>	109.2±10.6 <sup>*</sup>	119.5±19.3 <sup>*</sup>	161.4±23.7 <sup>*</sup>	233.2±35.4 <sup>*ac</sup>	262.3±29.0 <sup>**</sup>	413.6±44.0 <sup>**</sup>	378.8±39.0 <sup>**bc</sup>

**Table II.vii.** (continued).

	Day of pregnancy						Day of lactation	
	120th	125th	130th	135th	140th	145th	3rd	5th
<b>Group B</b>								
Mammary parenchyma grey-scale	64.5±4.9 <sup>ab</sup>	66.9±4.3 <sup>*</sup>	71.3±4.1 <sup>*</sup>	70.6±2.6 <sup>*</sup>	74.1±2.0 <sup>*ac</sup>	79.3±2.0 <sup>**</sup>	67.6±3.2 <sup>*</sup>	66.8±2.7 <sup>bc</sup>
External pudendal artery diametre (cm)	0.35±0.01 <sup>ab</sup>	0.38±0.01	0.38±0.01	0.39±0.01	0.44±0.01 <sup>ac</sup>	0.46±0.01	0.53±0.01	0.56±0.01 <sup>bc</sup>
Resistance index	0.66±0.02	0.58±0.01	0.62±0.03	0.61±0.02	0.61±0.02	0.60±0.02	0.58±0.02	0.61±0.02
Pulsatility index	1.24±0.09	0.97±0.04	1.07±0.05	1.06±0.04	1.06±0.05 <sup>**a</sup>	1.05±0.02 <sup>*</sup>	1.1±0.03 <sup>*</sup>	1.20±0.04 <sup>**a</sup>
Systolic:diastolic velocity ratio	2.96±0.17 <sup>ab</sup>	2.35±0.07	2.63±0.24	2.70±0.22	2.53±0.13 <sup>a</sup>	2.46±0.09	2.39±0.09	2.54±0.17 <sup>b</sup>
General pressure (mm Hg)	1.03±0.11 <sup>ab</sup>	1.33±0.11	1.28±0.25	1.50±0.20	1.63±0.13 <sup>ac</sup>	2.31±0.32	4.00±0.51	3.51±0.48 <sup>bc</sup>
Mean pressure (mm Hg)	0.34±0.05 <sup>ab</sup>	0.65±0.15	0.45±0.18	0.52±0.09	0.70±0.05 <sup>ac</sup>	0.84±0.12	1.47±0.24	1.29±0.24 <sup>bc</sup>
Mean velocity (m s <sup>-1</sup> )	0.26±0.02 <sup>ab</sup>	0.38±0.03	0.29±0.05	0.33±0.03	0.36±0.03 <sup>*ac</sup>	0.44±0.03 <sup>*</sup>	0.52±0.05 <sup>*</sup>	0.51±0.05 <sup>bc</sup>
Systolic acceleration (m s <sup>-2</sup> )	6.00±0.57 <sup>a</sup>	4.94±0.63	4.86±0.65	4.92±0.63	5.00±0.42 <sup>b</sup>	5.71±0.36	8.76±0.87	8.97±0.46 <sup>ab</sup>
Blood input (mL min <sup>-1</sup> )	114.6±11.2 <sup>*ab</sup>	191.4±34.3 <sup>*</sup>	169.4±22.5 <sup>*</sup>	241.8±34.9 <sup>*</sup>	282.9±30.2 <sup>*ac</sup>	397.0±17.7 <sup>**</sup>	649.0±50.2 <sup>**</sup>	482.0±42.0 <sup>**bc</sup>

Group A: ewes with  $\beta$ -hydroxybutyrate concentrations  $>1.2$  mmol L<sup>-1</sup> in two samples collected after the 129th day of pregnancy - Group B: all other ewes into the study.

\*  $P<0.05$ , \*\*  $P<0.01$  between the two groups at respective day for the respective parametre.

<sup>a</sup>, <sup>b</sup>, <sup>c</sup> for numbers marked with same superscript within a parametre (same row):  $P<0.05$ .

## Development of clinical signs relevant to pregnancy toxaemia in the experimental ewes

In four group A ewes (1 with two foetuses and 3 with three foetuses) (14% of ewes into the study, 25% of ewes into group A), clinical signs relevant to pregnancy toxaemia were recorded after the 140th day of pregnancy. These included depression (n=4), selective or reduced appetite (n=4) and recumbency (n=2).

Treatment was instigated immediately after detection of clinical signs. Treatment included administration of propylene glycol *per os* (30 g, *b.i.d.* × 3 days) and, in ewes with three foetuses, glucose intravenously (2 g, *q.i.d.* × 2 days), as well as twice daily 10-min long walks of the affected animals. In all affected animals, clinical signs subsided two to four days after instigation of treatment.

## Obstetrical problems in the experimental ewes

In group A ewes, dystocia was diagnosed in eight ewes [0.500, 95% confidence intervals (CI): 0.280-0.720]; in all cases, veterinary intervention was necessary to treat that. These eight ewes included the four animals with clinical signs relevant to pregnancy toxaemia. In two ewes, postural abnormalities of the foetus were evident (1 case: shoulder flexion, 1 case: lateral head deviation). In the other six ewes, expulsive deficiency was considered as the likely diagnosis.

In group B ewes, dystocia was diagnosed in one ewe only (0.083, 95% CI: 0.015-0.354); veterinary intervention was necessary. Postural abnormality of the foetus (shoulder flexion) was evident.

## Duration of pregnancy in the experimental ewes

Duration of pregnancy was significantly shorter in ewes with increased  $\beta$ -hydroxybuturate blood concentration during the last stage of pregnancy ( $P<0.001$ ). Mean ( $\pm$ standard error of the mean) duration of pregnancy of ewes into group A was  $145.75\pm 0.35$  days. Mean duration of pregnancy of ewes into group B was  $148.42\pm 0.38$  days.

## Birth bodyweight of lambs of the experimental ewes

Median birth bodyweight of lambs of ewes with one foetus was 5.00 kg (4.0-5.9 kg), of those ewes with two foetuses was 3.90 kg (2.0-5.3 kg) and of those with three foetuses was 2.95 kg (2.7-4.7 kg) ( $P < 0.001$ ). Median bodyweight of lambs of ewes allocated into group A was 3.5 kg (2.0-5.3 kg) and that of ewes in group B 4.0 kg (2.7-5.9 kg) ( $P = 0.016$ ).

There was a significant reverse correlation between nematode epg faecal counts and lamb birth bodyweight ( $r = -0.361$ ,  $P = 0.03$ ). There was also a significant reverse correlation between blood  $\beta$ -hydroxybutyrate concentrations and lamb birth bodyweight ( $r = -0.405$ ,  $P = 0.016$ ).

## Perinatal mortality of lambs of the experimental ewes

In five animals into group A, a dead foetus was found at lambing. Moreover, lambs born from two ewes died by the 2nd day of life (in one of these, a dead foetus had been found at lambing). Thus, perinatal mortality in offspring of group A ewes occurred in totally six animals (0.375, 95% CI: 0.185-0.614).

In group B, in the animal with obstetrical problems, a dead foetus was found (0.083 of group B ewes, 95% CI: 0.015-0.354).

## Post-partum disorders of the genital tract in the experimental ewes

In group A ewes, post-partum genital disorders were evident in four ewes (all had dystocia) (0.250, 95% CI: 0.102-0.495); of these, three animals developed mild metritis and one animal showed retention of foetal membranes. Immediately after diagnosis of the disorder, each ewe received an intramuscular injection of oxytocin (5 i.u.). The disorder resolved by the 3rd day after lambing (L3) with no further complications.

In group B ewes, no post-partum genital disorders were recorded (0.000, 95% CI: 0.000-0.243).

A summary of the obstetrical and post-partum problems of the experimental ewes is in Table II.viii.



**Table II.viii.** Incidence risk (95% confidence intervals) of peri-parturient problems diagnosed in ewes with (group A) or without (group B) increased  $\beta$ -hydroxybuturate blood concentration during the last stage of pregnancy.

Peri-parturient problem	Group A	Group B
Dystocia	0.500 <sup>a</sup> (0.280-0.720)	0.083 <sup>a</sup> (0.015-0.354)
Perinatal mortality of offspring	0.375 <sup>a</sup> (0.185-0.614)	0.083 <sup>a</sup> (0.015-0.354)
Post-partum disorders	0.250 <sup>a</sup> (0.102-0.495)	0.000 <sup>a</sup> (0.000-0.243)

<sup>a</sup>: between results in the same row  $P < 0.05$ .

### Milk yield of the experimental ewes

Mean collected quantity of milk from group A ewes on L3 was 78.1±4.2 mL and on L5 was 86.4±3.8 mL; respective quantities for group B ewes were 89.5±3.1 mL and 117.6±4.8 mL ( $P=0.019$  and  $P<0.001$ , respectively).

There was a significant reverse correlation between grey-scale intensity values in L3/L5 and milk quantities collected from these animals on the same days (for group A:  $r=-0.409$ ,  $P=0.010$ ; for group B:  $r=-0.4378$ ,  $P=0.016$ ); finally, there was also a significant reverse correlation between grey-scale intensity values on P145 and milk quantities collected from animals after lambing ( $r=-0.443$ ,  $P=0.015$ ). There was also a significant correlation between blood input values in L3/L5 and milk quantities collected from these animals on the same days (for group A:  $r=0.2709$ ,  $P=0.067$ ; for group B:  $r=0.4083$ ,  $P=0.024$ ); further, there was a significant correlation between blood input values on P130/P135/P140/P145 and milk quantities collected from these animals after lambing ( $r>0.41$ ,  $P<0.035$ ).

## B. AFTER CHALLENGE OF THE MAMMARY GLAND OF THE EXPERIMENTAL EWES

### Materials and methods

#### Experimental overview

As described previously, of the 28 ewes, in 16 increased  $\beta$ -hydroxybutyrate ( $>1.2$  mmol L<sup>-1</sup>) concentrations were recorded in blood samples during the last stage of pregnancy; these animals had been allocated into group A. In the other 12 ewes,  $\beta$ -hydroxybutyrate concentrations in blood samples during the last stage of pregnancy remained consistently  $<1.0$  mmol L<sup>-1</sup>; these animals had been allocated into group B. Conditions prescribed by legislation of the European Union in relation to animal experimentation procedures (Council Directive 86/809/EEC) were met during this work; the experiment was carried out under a licence for experimental procedures obtained from the Greek Ministry of Agriculture.

After lambing, animals within each group (A or B) were further allocated at random (at the toss of a coin) in subgroups (A1 or A2, B1 or B2). On the 3rd day post-partum, measurement of  $\beta$ -hydroxybutyrate blood concentrations was performed in all samples as described previously; it was confirmed that all ewes into subgroups A1 or A2 still had concentrations  $>1.2$  mmol L<sup>-1</sup> and all those in B1 or B2 still had concentrations  $<1.0$  mmol L<sup>-1</sup>. Then, on the 5th day post-partum, ewes in subgroups A1 or B1 were challenged by deposition into the teat duct of *Mannheimia haemolytica*, strain VSM08L (Mavrogianni et al. 2005, 2006b), which has been isolated from the teat duct of a ewe and is of known pathogenicity for the mammary gland of this species.

#### Feeding of the experimental ewes

After lambing, in ewes suckling one lamb, the ration provided per ewe was 0.30 kg of the specially formulated and prepared concentrate feed in mash form (Table II.i) plus 0.35 kg of alfalfa hay, offered twice daily. Respective figures for ewes suckling two lambs were 0.35 kg and 0.35 kg and for ewes suckling three lambs were 0.50 kg and 0.35 kg. Then, on the 5th day of the lactation period, quantities of concentrate feed to ewes were as follows: 0.35, 0.40 and 0.55 kg for ewes

suckling one, two or three lambs, respectively, offered twice daily. Barley straw was provided into the pens thrice daily, whilst water was available *ad libitum*.

## Preparation of inoculum and inoculation procedure of the experimental ewes

The *M. haemolytica* strain was grown on Columbia blood agar and checked for purity; then it was inoculated into Soy-broth (BioMerieux S.A., Marcy-l'Etoile, Γαλλία) and incubated aerobically at 37 °C for 5 h. Serial dilutions of the broth culture into sterile phosphate-buffer-saline pH 7.3 (PBS Dulbecco-Biochrom AG, Berlin, Germany) were carried out; finally, 0.2 mL of the desired dilution was withdrawn with a syringe. The inoculum contained 1200 to 1250 c.f.u., as estimated by the method of Miles and Misra (1938).

To ensure sterile conditions, on the day before inoculation, the hairs of the teats of the experimental ewes were clipped by using fine scissors and the skin of the udder was scrubbed by using chlorhexidine. Ewes were challenged as follows; initially, the teat was disinfected by using iodine povidone solution; then, a sterile plastic fine catheter (Abbocath®) 20 G, 2 mm-long was inserted into the teat; the syringe was attached to the catheter and the bacterial suspension was deposited inside the teat. Of the ewes in subgroup A1 or B1, half were challenged into the left teat and half into the right teat (animals were allocated for challenge into the left or right teat at random). The same technique was used to inject 0.2 mL of PBS into the contralateral teat of each ewe. PBS was also injected into the respective site of both teats (i.e., 2 mm deep) of ewes in subgroup A2 or B2. After challenge, lambs were kept away from their dams for 2 hours. Details are in Table II.ix.

**Table II.ix.** Summary of the experimental design.

Subgroup	n	β-hydroxybutyrate blood concentrations at late gestation	<i>M. haemolytica</i> deposition into the teat duct	Days after challenge, when tissue samples were collected and ewes were removed of the study
A1	8	>1.2 mmol L <sup>-1</sup>	Yes	D1 (n=2), D2 (n=2), D3 (n=2), D4 (n=2)
A2	8	>1.2 mmol L <sup>-1</sup>	No	D1 (n=2), D2 (n=2), D3 (n=2), D4 (n=2)
B1	8	<1.0 mmol L <sup>-1</sup>	Yes	D1 (n=2), D2 (n=2), D3 (n=2), D4 (n=2)
B2	4	<1.0 mmol L <sup>-1</sup>	No	D1 (n=1), D2 (n=1), D3 (n=1), D4 (n=1)

D0: day of challenge

## Pre- and post-challenge procedures in the experimental ewes

### *Clinical examination of the udder - Sample collection*

A general clinical examination of the experimental ewes, with special reference to their udder, was performed before challenge, on the 3rd and the 5th days after lambing (D-2: 2 days before challenge, D0: day of challenge, respectively). Samples of teat duct material and milk samples were then collected on those occasions.

After challenge, detailed examinations of the animals (including their udder) and collection of samples were carried out 12 h after challenge (D0+12 h) and thereafter on D1 (1 day after challenge), D2, D3 and D4 (unless of course, a ewe had been removed from the study before that).

### *Ultrasonographic examination of the udder*

Starting on the 3rd day of the lactation period, i.e. two days before challenge (D-2), in the morning of each test day, an ultrasonographic examination of the udder was performed; the examination was repeated on D0, before challenge. After challenge, the examination was performed 3 h after challenge (D0 + 3 h; only subgroups A1 and B1 - only Doppler examination), 6 h after challenge (D0 + 6 h; only subgroups A1 and B1 - only Doppler examination), 12 h after challenge (D0 +12 h) and thereafter on D1, D2, D3 and D4 (unless of course, a ewe had been removed from the study before that).

Before each examination, lamb(s) of the respective ewe had been removed from the dam for one hour. The examination was performed with the animal on the standing position and restrained inside a crate, using the support of an assistant. For examination, hair on the abdomen and the udder had been fully clipped. Coupling gel was applied. Initially, the left side of the udder was imaged and subsequently, the whole procedure was repeated for the right side of the udder.

B-mode and Doppler examination of the mammary parenchyma was performed by the same equipment (MyLab<sup>®</sup> 30; ESAOTE SpA, Genova, Italy) and by using the same techniques as described in detail already. In B-mode examination, 10.0 MHz imaging frequency and 60 mm scanning depth were used for this procedure. In Doppler examination, 5.0 to 6.6 MHz imaging frequency and 60 mm scanning depth were used for this procedure.

Subsequently, ultrasonographic evaluation was performed in the teat of the ewes. Ample quantity of coupling gel was applied. B-mode ultrasonographic examination was initially performed. The transducer was placed longitudinally to the teat; 12.0 MHz imaging frequency and 20 to 30 mm scanning depth were used for this procedure. All the teat was imaged in one section. Then,

colour Doppler evaluation was performed in the subcutaneous vessels of the teat. Adjustments necessary for optimal colour image quality were made, with the pulse repetition frequency maintained constant, in order to ensure comparability of images taken; 6.6 MHz imaging frequency and 20 to 30 mm scanning depth were used for this procedure

In all cases, images (B-mode or Doppler) were frozen and saved on the equipment hard-disk for performing subsequently appropriate measurements and data analysis.

#### *Bacteriological and cytological examinations of udder samples*

Conventional bacteriological examination of samples was carried out (Barrow and Feltham 1993, Euzeby 1997). The California Mastitis Test (CMT) was then carried out and leucocyte subpopulations were identified in milk samples.

#### *Mammilectomy and mammary parenchyma biopsy*

Two of the ewes in each of subgroups A1, A2 and B1 were subjected to mammilectomy of the inoculated teat and mammary parenchyma biopsy of both sides of the udder on each of D1, D2, D3 and D4. One ewe in subgroup B2 was subjected to the same procedures on each of the above days.

The operation was carried out under sedation and cranial epidural block; opioids were also administered for analgesia (Galatos 2011). All surgical procedures were performed under strict aseptic conditions. Initially, the teat was ligated at its base and excised. At the same time, a biopsy of the parenchyma of both mammary glands was also performed (Fthenakis and Jones 1990b); two cubes of tissue, approximately 1×1×1 cm, were removed from the ventral part of each gland. After obtaining the tissue samples and repairing the wounds, ewes were removed from the experiment.

#### *Procedures in udder tissue samples*

The skin of each excised teat and the subcutaneous tissues were incised with a sterile blade; initially, the mucosa of the teat cistern (*sinus papillaris*) was exposed and subsequently the teat duct (*ductus papillaris*) was incised and its mucosa was exposed. A new blade was used for scraping the mucosa of the teat cistern, whilst another one was used to scrape the mucosa of the teat duct. An electronic cutimetre was used to measure 2 mm from the teat orifice, in order to determine the precise site of the teat, where the inoculum had been deposited (Mavrogianni et al. 2005). The scrapings from the inside of the teat and one of the cubes of tissue from each

mammary gland were used for bacteriological examination, by applying conventional techniques (Barrow and Feltham 1993, Euzeby 1997).

Longitudinal sections, involving all the structures of the teat, were carried out for histopathological examination. The second tissue sample from each mammary parenchyma was also processed. Tissue samples were fixed in 10% neutral-buffered formalin and embedded in paraffin wax. Haematoxylin and eosin (HE) standard staining procedures were performed for histopathological study of tissue samples.

## Data management and analysis

### *Definitions*

Clinical mastitis was defined as the presence of clinically evident abnormalities in the mammary gland or the mammary secretion. Subclinical mastitis was defined as the simultaneous isolation of bacteria from milk samples and the detection of positive CMT scores, with no clinically evident abnormalities.

### *Cytological examination of milk samples*

In the California Mastitis Test, reactions scored  $\geq 1$  were considered to be indicative of increased cellular content in milk (Fthenakis 1995)

### *Results taken into account during the ultrasonographic examination of the udder of the experimental ewes*

During the ultrasonographic examination of the udder, three sets of images were recorded for each side of the udder: (i) images of mammary parenchyma, (ii) images of cross-sections of external pudendal artery and (iii) spectral waveforms of external pudendal artery. All images were initially evaluated visually for presence of abnormal structures therein. Then, data management was performed on images that had been saved at the time of examination. Data for left and right mammary gland of the same animal were considered separately.

Stored images of mammary parenchyma were processed to produce intensity values for analysis of grey-scale, with results were expressed on a 0 (black) to 255 (white) scale. Stored images of cross-sections of the external pudendal artery were also processed for calculation of the internal diameter of the vessel and of the same haemodynamic parameters of the external pudendal artery, i.e., resistance index, pulsatility index, systolic:diastolic velocity ratio, general pressure, mean pressure, mean velocity, systolic acceleration and blood input.

Blood input in teat vessels was presented by scoring the amount of colour on each image in a 1 to 4 scale, compared to 0, which was assigned on each teat before challenge; hence, scores 1 or 2 were given based on relative increase of diameter ( $\leq 50\%$  increase for score 1 and  $>50\%$  for score 2) of already imaged vessels; scores 3 or 4 were given only if 'new' (i.e., not visible in previous images) vessels ( $\leq 2$  vessels for score 3 and  $>2$  vessels for score 4) were imaged.

Finally, results from a ewe in subgroup A2, which developed spontaneous subclinical mastitis during the study were not taken into account in the analysis.

#### *Scoring of pathological findings in tissue samples*

A scoring system previously devised (Fragkou et al. 2007) was used and numerical values were assigned for the macroscopic or histological pathological findings in the teats or in the mammary parenchyma of the experimental animals. The system is detailed in Table II.x.

**Table II.x.** Description of scores given for pathological findings in the teats or the mammary parenchyma of ewes inoculated with *M. haemolytica* (Fragkou 2007).

Score	Description
<b>Teat – Macroscopic pathological findings</b>	
0	Normal
1	Presence of folds on the mucosa of the teat
2	Hyperaemia of the mucosa of the teat
3	Thickening of the mucosa of the teat, with increased number of folds and presence of petechiae
4	Extreme thickening of the mucosa of the teat, with loss of the separation of the compartments of the teat
<b>Teat – Histopathological findings</b>	
0	Normal
1	Presence of a few, scattered leucocytes
2	Presence of increased numbers of leucocytes clustered under the epithelium of the teat
3	Presence of high numbers of leucocytes evenly distributed under the epithelium of the teat
4	Presence of high numbers of leucocytes, plus hyperplasia of lymphoid tissue at the teat duct - teat cistern border
<b>Parenchyma – Histopathological findings</b>	
0	Normal
1	Presence of a few, scattered leucocytes
2	Presence of increased numbers of leucocytes clustered in the intra- and inter-alveolar area
3	Diffuse presence of leucocytes, extravasation and destruction of epithelial cells
4	Haemorrhages, destruction of alveoli and loss of the internal architecture of the parenchyma

### *Statistical computations*

All data were entered into Excel spreadsheets (Microsoft Corporation, Redmond, WA, USA). Initially, descriptive statistics for all parameters were performed.

The proportions of ewes with clinical or subclinical mastitis, of bacterial isolations from teat duct material or mammary secretion and of CMT positive results, as well as the proportion of bacterial isolations from tissue samples were calculated. Proportions were compared between groups (A *versus* B) and between inoculated or non-inoculated animals (A1 plus B1 *versus* A2 plus B2), as well as between corresponding subgroups (A1 *versus* B1, A2 *versus* B2). Separate analyses were carried out for the inoculated and the contralateral side. Finally, comparisons were made between the above proportions in samples from inoculated or contralateral sides within the same subgroup. Chi-square tests were performed for comparisons of proportions between respective groups / subgroups.

Wilcoxon Signed Rank tests were performed to evaluate differences in pathology score medians between groups (A *versus* B) and between inoculated or non-inoculated animals (A1 and B1 *versus* A2 and B2), as well as between corresponding subgroups (A1 *versus* B1, A2 *versus* B2). They were also used to test differences between pathology scores in the mammary parenchyma ipsilaterally or contralaterally to the inoculated teat within the same subgroup.

For the study of the results of the ultrasonographic examinations, repeated measures mixed effect linear regression models were used to determine whether outcomes in the four subgroups of the study changed over the course of the study period. Effect of experimental subjects (animals) was included as random effect in the model. Models were adjusted for repeated measures within animals. Independent variables (fixed effects) included mammary gland, experimental subgroup (A1, A2, B1, B2), time of the study (i.e., D-2, D0, D0+3 h, D0+6 h, D0+12 h, D1, D2, D3, D4) and a time of the study by mammary gland or a time of the study by experimental subgroup interaction. For each parameter, an analysis that took into account all data (9 time-points) was carried out. Subsequently, a more simplified analysis was performed, which accounted data in the three time-periods of the study (i.e., 'Before challenge', 'Day of challenge', 'After day of challenge').

An electronic data management tool was employed (Lowry 2012, 2015). Significance level was set at  $P \leq 0.05$ .



## Results

### Findings in examinations before challenge of the experimental ewes

The mammary glands and the teats of all ewes were clinically healthy in both examinations performed before challenge. The teats were soft with no external abnormalities. Moreover, no bacteria were isolated from any teat duct material or milk samples obtained. California Mastitis Test scores in milk samples were negative (scores 'negative' or 'trace') and observation of Giemsa-stained milk films revealed only scarce presence of macrophages therein (on average, one cell per 10 fields with the 10× objective lens).

No abnormal findings were evident during the observation of ultrasonographic images of the mammary parenchyma and the teat. The mammary parenchyma appeared mildly heterogeneous. Presence of milk was evident in images of both the parenchyma and the teat.

### Clinical, bacteriological and cytological findings after challenge of the experimental ewes

#### *Inoculated (or corresponding) side of the udder*

In subgroup A1, three ewes developed (D1 to D3) signs of clinical mastitis (changes in the mammary secretion, which became serous or sero-haemorrhagic and contained flakes, and presence of increased size and temperature of the teat and the mammary parenchyma). The other five animals in the subgroup developed subclinical mastitis, i.e., in total, 8/8 animals developed mastitis. *M. haemolytica* was isolated in pure culture from teat duct material and milk samples of all ewes (8/8) (in total, 28/28 and 21/28 isolations, respectively), starting 12 h post-inoculation and until D4. The CMT increased (>'1') in all ewes (27/28 positive samples). Leucocytes were seen in Giemsa-stained secretion films; their great majority ( $\geq 95\%$ ) consisted of neutrophils, with some macrophages and lymphocytes also present.

In subgroup A2, one ewe developed subclinical mastitis on D2; *M. haemolytica* was isolated on two occasions (D2, D3) from the teat duct material and the milk sample of that ewe (in total, 2/28 and 2/28 isolations, respectively). The CMT was increased in milk samples (D2, D3) from that animal (2/28 positive samples in total). Neutrophils were mainly seen in the secretion films from that animal, whilst scarce macrophages were observed in the samples from the other ewes.

In subgroup B1, four ewes developed subclinical mastitis ( $P=0.021$  when compared to animals that developed mastitis in subgroup A1). *M. haemolytica* was isolated in pure culture from teat duct material samples of all ewes (8/8), as well as from milk samples of four ewes, starting 12 h post-challenge and until D4 (in total, 25/28 and 11/28 isolations, respectively). The CMT increased ( $>'1'$ ) in all ewes (27/28 positive samples). Leucocytes were seen in Giemsa-stained secretion films; their great majority ( $\geq 80\%$ ) consisted of neutrophils, with macrophages and lymphocytes also evident.

In subgroup B2, no abnormal findings were recorded ( $P=0.460$  when compared to animals that developed mastitis in subgroup A2). No bacteria were isolated from teat duct material or milk samples from these ewes (in total, 0/14 and 0/14 isolations) and the CMT remained negative (0/14 positive samples). Macrophages were seen scarcely in the milk samples.

In total, there was a tendency for more bacterial isolations from the inoculated side of ewes with increased  $\beta$ -hydroxybuturate blood concentration during the last stage of the preceding pregnancy (group A) than from that of ewes with no increased  $\beta$ -hydroxybuturate blood concentration during the last stage of the preceding pregnancy (group B) (53/112 *versus* 36/84, respectively), which, however, was not statistically corroborated ( $P=0.535$ ). Ewes that were inoculated (A1 plus B1), yielded more bacterial isolations than uninoculated controls (A2 plus B2) (85/112 *versus* 4/84, respectively;  $P<0.001$ ). Comparisons between subgroups revealed that isolations from inoculated (or corresponding) side of A1 or A2 were greater than isolations from respective side of B1 or B2 (for isolations from teat duct material or mammary secretion samples from subgroup A1 *versus* B1,  $P=0.075$  and  $P=0.007$ , respectively; for isolations from subgroup A2 *versus* B2,  $P=0.305$ ).

When comparing CMT scores, significantly more positive CMT results were evident in samples from inoculated side of ewes than from uninoculated controls ( $P<0.001$ ), whilst all other comparisons did not reveal significant differences ( $P>0.2$ ). Detailed results are in Table II.xi.

**Table II.xi.** Clinical findings, sequential isolation of *M. haemolytica* and results of California Mastitis Test in the udder of ewes with one teat inoculated or not with the organism.

	12 h post-inoculation				D1				D2				D3				D4			
	A1	A2	B1	B2	A1	A2	B1	B2	A1	A2	B1	B2	A1	A2	B1	B2	A1	A2	B1	B2
<b>Clinical signs</b>																				
Presence	0/8	0/8	0/8	0/4	3/8	0/8	0/8	0/4	3/6	0/6	0/6	0/3	2/4	0/4	0/4	0/2	0/2	0/2	0/2	0/1
<b><i>M. haemolytica</i> isolations</b>																				
Teat duct material	8/8	0/8	8/8	0/4	8/8	0/8	8/8	0/4	6/6	1/6	5/6	0/3	4/4	1/4	2/4	0/2	2/2	0/2	2/2	0/1
Secretion	1/8	0/8	2/8	0/4	8/8	0/8	2/8	0/4	6/6	1/6	4/6	0/3	4/4	1/4	2/4	0/2	2/2	0/2	1/2	0/1
<b>California Mastitis Test results</b>																				
Positive	7/8	0/8	8/8	0/4	8/8	0/8	8/8	0/4	6/6	1/6	6/6	0/3	4/4	1/4	4/4	0/2	2/2	0/2	1/2	0/1

D1, D2, D3, D4 = days after inoculation

A1, A2, B1, B2 = subgroups (A\*: ewes with increased  $\beta$ -hydroxybuturate blood concentration during the last stage of the preceding pregnancy, B\*: ewes with no increased  $\beta$ -hydroxybuturate blood concentration during the last stage of the preceding pregnancy; \*1: ewes challenged with deposition of *M. haemolytica* at the teat duct on the 5<sup>th</sup> day post-partum, \*2: uninfected control ewes).

n/m=positive results out of total animals sampled

#### *Non-inoculated side of the udder*


In subgroup A1, *M. haemolytica* was isolated on two occasions (12 h post-inoculation and D1) from milk samples, but not from teat duct material samples, of the contralateral side of one ewe (in total, 0/28 and 2/28 isolations from teat duct material and milk samples, respectively). Increased CMT scores were also recorded in milk samples from that animal. In all other subgroups, no bacteria were isolated from the contralateral side of the udder (teat duct material and milk samples) of the experimental ewes. Comparisons performed did not reveal any significant difference ( $P>0.2$ ).

When proportions of bacterial isolations or positive CMT scores in samples from inoculated or contralateral side were compared, significance was recorded in subgroups A1 or B1 ( $P<0.001$ ), but not in subgroups A2 or B2 ( $P>0.14$ ).


#### Findings of ultrasonographic examination of the udder of the experimental ewes

After D1, the duct cistern (*sinus papillaris*) and the gland cistern (*sinus lactiferous*) were observed as anechoic cavities, with presence of hyperechoic particles therein, initially in the duct cistern (Fig. II.9.) and later in the gland cistern. Progressively also (after D2), the ultrasonographic pattern observed in the parenchyma became markedly heterogeneous, with presence of characteristically coarse formations within (Figs II.10. and II.11.).


**Figure II.9.** B-mode ultrasonographic appearance of the teat of a ewe (subgroup A1), before challenge with *M. haemolytica* into the teat duct showing the teat cistern (*sinus papillaris*) as an anechoic cavity (left picture) and 2 days after challenge (D2) with presence of hyperechoic particles therein, obvious in a 'snow-like' appearance at the right part of the cavity (i.e., at the lower part of the cistern) (right picture). Images taken on a MyLab® 30 ultrasonography system with linear transducer, imaging frequency: 12.0 MHz - scanning depth: 20 to 30 mm.



**Figure II.10.** B-mode ultrasonographic appearance of the parenchyma of the inoculated side of the udder of a ewe (subgroup A1), before challenge with *M. haemolytica* into the teat duct (left picture) and 4 days after challenge (D4) (right picture). Images taken immediately before the gland cistern on a MyLab® 30 ultrasonography system with linear transducer, imaging frequency: 10.0 MHz - scanning depth: 60 mm.




**Figure II.11.** B-mode ultrasonographic appearance of the parenchyma of the inoculated side of the udder of a ewe (subgroup A1) 4 days after challenge with *M. haemolytica* into the teat duct (D4) (left picture) and of the parenchyma of the contralateral side of the udder (right picture). Images taken immediately before the gland cistern on a MyLab® 30 ultrasonography system with linear transducer, imaging frequency: 10.0 MHz - scanning depth: 60 mm.




After challenge, median grey-scale intensity values in the inoculated side of subgroup A1 and B1 temporarily decreased, but no evidence of significant differences was produced. Recorded values were 66.9 'Before challenge' to 64.8 on 'Day of challenge' to 79.4 'After day of challenge' for subgroup A1 and 73.7 'Before challenge' to 69.1 to 'Day of challenge' to 71.2 'After day of challenge' for subgroup B1 ( $P < 0.19$  for subgroup A1 and  $P < 0.055$  for subgroup B1) (for differences between A1 and B1,  $P < 0.17$ ). A similar pattern was evident in the contralateral side of subgroup B1 ewes only ( $P = 0.002$  on 'Day of challenge' compared to 'Before challenge'), but the opposite trend was noticed in the respective side of subgroup A1 ewes ( $P = 0.022$  on 'Day of challenge' compared to 'Before challenge') (Fig. II.12.).

**Figure II.12.** Time-series (before and after challenge with *M. haemolytica* into the teat duct) graphs of the grey-scale intensity values (0-255) of the mammary gland of ewes in subgroup A1 (red lines) or B1 (blue lines), which had one teat inoculated (solid lines) or not (dotted lines).



After challenge (subgroups A1 and B1), there was clear evidence for an increase in the diameter of the external pudendal artery already 6 h after challenge ( $P < 0.02$ ), which later (D1) decreased to pre-challenge figures ( $P > 0.12$ ). Some increase was also observed in the external pudendal artery of the contralateral side of the udder in both subgroups, although evidence indicated a significant difference between the two sides only in subgroup B1 ( $P = 0.28$  for subgroup A1,  $P = 0.025$  for subgroup B1) (Fig. II.13.). Similar changes were not observed in subgroups A2 and B2 ( $P > 0.11$ ).

**Figure II.13.** Time-series (before and after challenge with *M. haemolytica* into the teat duct) graphs of the diameter (cm) of the external pudendal artery of ewes in subgroup A1 (red lines) or B1 (blue lines), which had one teat inoculated (solid lines) or not (dotted lines).




The spectral display in Doppler mode was observed as a broad band structure. After challenge, there was an abrupt and excessive increase in blood input into the mammary parenchyma of the challenged side in ewes in subgroups A1 and B1, as follows; median input in

subgroup A1 ewes was 400 mL min<sup>-1</sup> 'Before challenge' to 513 mL min<sup>-1</sup> on 'Day of challenge' to 318 mL min<sup>-1</sup> after that ( $P<0.001$ ) and respective values in subgroup B1 were 527 mL min<sup>-1</sup>, 709 mL min<sup>-1</sup> and 418 mL min<sup>-1</sup> ( $P<0.003$ ) (for differences between A1 and B1,  $P=0.006$  on 'Day of challenge' and  $P=0.055$  'After day of challenge'). There was some, but not significant, increase of blood input into the contralateral gland: 400 mL min<sup>-1</sup>, 452 mL min<sup>-1</sup> and 332 for subgroup A1 and 523 mL min<sup>-1</sup>, 538 mL min<sup>-1</sup> and 321 mL min<sup>-1</sup> for subgroup B1, respectively ( $P<0.065$  for subgroup A1,  $P<0.42$  for subgroup B1) and evidence showed that blood input into the parenchyma after challenge was greater in the inoculated than the contralateral side ( $P<0.03$ ) (Figs II.14. and II.15.). No such changes were evident in the uninfected animals.

Mean velocity of blood increased immediately after challenge; in ewes of subgroup A1, it returned to previous values later than in ewes of subgroup B1 (comparisons between values obtained 'Before challenge' to values obtained on 'Day of challenge':  $P=0.001$  for subgroup A1,  $P<0.001$  for subgroup B1; comparisons between values obtained 'Before challenge' to values obtained 'After day of challenge':  $P=0.004$  for subgroup A1,  $P=0.242$  for subgroup B1; comparisons between values obtained on 'Day of challenge' to values obtained 'After day of challenge':  $P=0.38$  for subgroup A1,  $P<0.001$  for subgroup B1). A similar pattern, but of smaller magnitude, was recorded in the contralateral mammary glands of these ewes ( $P$  values varying from  $<0.001$  to 0.124) (Fig. II.16.). No such changes were evident in the uninfected animals ( $P$  values varying from  $<0.135$  to 0.5).


**Figure II.14.** Time-series (before and after challenge with *M. haemolytica* into the teat duct) graphs of the Doppler ultrasonography results of the external pudendal artery of ewes in subgroup A1 (red lines) or B1 (blue lines), which had one teat inoculated (solid lines) or not (dotted lines): (a) blood input (mL min<sup>-1</sup>), (b) mean velocity (m s<sup>-1</sup>).





**Figure II.14.** (continued).

(b)



**Figure II.15.** Serial (before and after challenge with *M. haemolytica* into the teat duct; from left to right: picture on D-2, D0+12 h, D2) spectral waveforms of the external pudendal artery of (a) a ewe with increased  $\beta$ -hydroxybuturate blood concentration during the last stage of the preceding pregnancy, which was subsequently inoculated into the ipsilateral teat (on D0) (subgroup A1), (b) a ewe with no increased  $\beta$ -hydroxybuturate blood concentration during the last stage of the preceding pregnancy, which was subsequently inoculated into the ipsilateral teat (on D0) (subgroup B1) or (c) a ewe with no increased  $\beta$ -hydroxybuturate blood concentration during the last stage of the preceding pregnancy, which was subsequently inoculated into the contralateral teat (on D0) (subgroup B1). Images taken and processed on a MyLab<sup>®</sup> 30 ultrasonography system with linear transducer, imaging frequency: 5.0 to 6.6 MHz - scanning depth 60 mm.

(a)






Figure II.15. (continued).

(b)



(c)



In both subgroups A1 and B1, blood input into the teat increased immediately after challenge and remained increased during the study ( $P < 0.001$  for all comparisons) in the inoculated teat. A similar increase was evident in the contralateral teat, but of smaller magnitude ( $P$  values varying from 0.023 to 0.49), with a significant difference between the two sides of the same animal ( $P < 0.002$  for all comparisons) (Fig. II.16.).

Detailed results are in Tables II.xii. and II.xiii.

**Figure II.16.** Serial (before and after challenge with *M. haemolytica* into the teat duct; from left to right: picture on D-2, D0+12 h, D2) colour Doppler images of the subcutaneous vessels of the teat of (a) a ewe with increased  $\beta$ -hydroxybuturate blood concentration during the last stage of the preceding pregnancy, which was subsequently inoculated into that teat (on D0) (subgroup A1), (b) a ewe with no increased  $\beta$ -hydroxybuturate blood concentration during the last stage of the preceding pregnancy, which was subsequently inoculated into that teat (on D0) (subgroup B1) or (c) a ewe with no increased  $\beta$ -hydroxybuturate blood concentration during the last stage of the preceding pregnancy, which was subsequently inoculated into the contralateral teat (on D0) (subgroup B1). Images taken and processed on a MyLab<sup>®</sup> 30 ultrasonography system with linear transducer; imaging frequency: 6.6 MHz - scanning depth: 20 to 30 mm.

(a)






Figure II.16. (continued).

(b)



(c)



**Table II.xii.** Quantitative results [median (min.-max.)] of ultrasonographic (B-mode or Doppler) examination of the udder in ewes with one teat inoculated or not with *M. haemolytica*; results of the two sides of the udder are considered separately.

	Before challenge	Day of challenge	After day of challenge	Before challenge	Day of challenge	After day of challenge
Subgroup A1	Inoculated side of the udder			Contralateral side of the udder		
Mammary parenchyma grey-scale	73.7 <sup>a</sup> (65.9-96.1)	69.1 <sup>a,m</sup> (38.3-89.0)	71.2 <sup>z</sup> (37.8-99.2)	75.5 (6.4-86.4)	72.8 <sup>m</sup> (42.5-109.6)	74.3 (47.3-101.0)
External pudendal artery diameter (cm)	0.55 <sup>a</sup> (0.51-0.64)	0.57 <sup>a</sup> (0.52-0.75)	0.55 <sup>m</sup> (0.50-0.68)	0.55 <sup>a</sup> (0.51-0.60)	0.57 <sup>a,b</sup> (0.49-0.63)	0.55 <sup>b,m</sup> (0.47-0.66)
Resistance index	0.62 <sup>a</sup> (0.53-0.64)	0.54 <sup>a,b,m,x</sup> (0.44-0.61)	0.59 <sup>b,z</sup> (0.44-0.76)	0.59 <sup>a</sup> (0.55-0.64)	0.57 <sup>a,b,m,x,z</sup> (0.47-0.68)	0.60 <sup>b</sup> (0.49-0.72)
Pulsatility index	1.02 <sup>a,z</sup> (0.82-1.20)	0.83 <sup>a,b,m,z</sup> (0.64-0.99)	0.98 <sup>b,z</sup> (0.69-1.69)	1.00 <sup>a</sup> (0.86-1.16)	0.91 <sup>a,b,m</sup> (0.68-1.28)	1.06 <sup>b</sup> (0.71-1.48)
Systolic:diastolic velocity ratio	2.40 <sup>a,z</sup> (2.20-2.73)	2.51 <sup>b,z</sup> (2.17-3.33)	2.40 <sup>a,b,z</sup> (1.97-2.61)	2.35 (2.13-2.78)	2.32 <sup>x</sup> (2.00-3.10)	2.33 <sup>x</sup> (1.95-3.19)
General pressure (mm Hg)	4.40 <sup>a,b</sup> (3.30-7.50)	6.80 <sup>a,c,z</sup> (3.40-7.50)	7.50 <sup>b,c,m,x,z</sup> (3.70-7.50)	4.60 <sup>a,x</sup> (3.30-7.50)	7.00 <sup>a,x,z</sup> (3.90-7.50)	5.10 <sup>m,x,z</sup> (3.20-7.50)
Mean pressure (mm Hg)	1.90 <sup>a,b,z</sup> (1.50-3.00)	3.00 <sup>a,z</sup> (1.80-3.70)	2.60 <sup>b,x,z</sup> (1.20-3.90)	1.80 <sup>a</sup> (1.50-2.90)	2.80 <sup>a,b,x,z</sup> (1.50-5.60)	2.00 <sup>b,x,z</sup> (0.80-5.50)
Mean velocity (m s <sup>-1</sup> )	0.65 <sup>a,b,z</sup> (0.54-0.84)	0.75 <sup>a,z,x</sup> (0.59-0.98)	0.80 <sup>b,m,x,z</sup> (0.51-0.97)	0.61 <sup>a,b,z</sup> (0.49-0.75)	0.71 <sup>a,x,z</sup> (0.60-0.94)	0.67 <sup>b,m,x,z</sup> (0.43-0.94)
Systolic acceleration (m s <sup>-2</sup> )	9.24 <sup>a</sup> (4.30-15.71)	9.55 <sup>b,z</sup> (4.35-15.78)	10.25 <sup>a,b,x,z</sup> (6.42-20.76)	9.53 (2.47-15.36)	9.39 <sup>a,x,z</sup> (1.95-11.97)	9.83 <sup>a,z</sup> (7.51-14.58)
Blood input into parenchyma (mL min <sup>-1</sup> )	400 <sup>a,b,z</sup> (296-572)	513 <sup>a,c,m,x,z</sup> (325-1043)	318 <sup>b,c,x</sup> (232-737)	400 <sup>a,z</sup> (292-656)	452 <sup>b,m,z</sup> (242-728)	332 <sup>a,b</sup> (203-508)
Blood input at teat (score 0-4)	0 <sup>a,b</sup> (0-2)	3 <sup>a,b,c,m,x</sup> (1-4)	2 <sup>b,c,m,x,z</sup> (0-4)	0 <sup>z</sup> (0-3)	1 <sup>m</sup> (0-2)	1 <sup>m,z</sup> (0-3)

**Table II.xii.** (continued).

	Before challenge	Day of challenge	After day of challenge	Before challenge	Day of challenge	After day of challenge
<b>Subgroup A2</b>	<b>Ipsilateral to inoculated side of the udder</b>			<b>Contralateral side of the udder</b>		
Mammary parenchyma grey-scale	75.2 (53.7-82.5)	67.2 (53.8-94.4)	73.2 (50.0-127.5)	74.2 (46.0-88.9)	58.2 (43.7-104.0)	71.7 (53.7-123.7)
External pudendal artery diametre (cm)	0.53 (0.37-0.63)	0.52 (0.45-0.71)	0.53 (0.37-0.76)	0.55 (0.39-0.70)	0.53 (0.42-0.63)	0.54 (0.40-0.66)
Resistance index	0.58 (0.53-0.67)	0.65 <sup>x</sup> (0.47-0.70)	0.56 (0.47-0.73)	0.59 (0.52-0.68)	0.61 <sup>x</sup> (0.56-0.70)	0.61 (0.49-0.72)
Pulsatility index	0.99 (0.90-1.12)	0.92 (0.63-1.08)	1.00 (0.75-1.63)	0.98 (0.74-1.39)	0.95 <sup>a</sup> (0.78-1.08)	1.03 <sup>a</sup> (0.74-1.66)
Systolic:diastolic velocity ratio	2.42 (1.75-3.06)	2.82 (2.10-3.38)	2.27 (1.87-3.76)	2.40 (2.20-3.13)	2.57 <sup>x</sup> (2.30-3.28)	2.51 <sup>x</sup> (1.97-3.56)
General pressure (mm Hg)	4.30 (3.60-7.50)	4.50 (2.70-7.50)	4.10 <sup>x</sup> (1.70-7.50)	4.10 <sup>x</sup> (3.30-4.80)	3.40 <sup>x</sup> (2.90-4.10)	4.00 <sup>x</sup> (1.40-7.50)
Mean pressure (mm Hg)	1.80 (1.30-3.50)	2.00 (0.80-3.60)	2.00 <sup>x</sup> (0.60-3.40)	1.70 (1.40-3.00)	1.20 <sup>x</sup> (0.60-2.70)	1.30 <sup>x</sup> (0.50-3.40)
Mean velocity (m s <sup>-1</sup> )	0.63 (0.52-0.78)	0.60 <sup>x</sup> (0.55-0.80)	0.60 <sup>x</sup> (0.49-0.70)	0.60 (0.54-0.66)	0.61 <sup>x</sup> (0.52-0.63)	0.59 <sup>x</sup> (0.41-0.68)
Systolic acceleration (m s <sup>-2</sup> )	9.45 <sup>m</sup> (6.20-11.99)	7.55 (4.49-10.6)	8.58 <sup>x</sup> (5.11-14.19)	9.51 <sup>a,m</sup> (6.92-14.30)	7.18 <sup>a,x</sup> (5.35-10.74)	8.21 (5.03-17.89)
Blood input into parenchyma (mL min <sup>-1</sup> )	401 <sup>a</sup> (288-795)	335 <sup>x</sup> (148-512)	267 <sup>a,x</sup> (120-392)	384 <sup>a</sup> (268-995)	246 (165-624)	258 <sup>a</sup> (120-996)
Blood input at teat (score 0-4)	0 (0-2)	0 <sup>x</sup> (0-2)	0.5 <sup>x</sup> (0-3)	0 (0-3)	0 (0-3)	0 (0-3)



**Table II.xii.** (continued).

	Before challenge	Day of challenge	After day of challenge	Before challenge	Day of challenge	After day of challenge
<b>Subgroup B1</b>	<b>Inoculated side of the udder</b>			<b>Contralateral side of the udder</b>		
Mammary parenchyma grey-scale	66.9 <sup>a</sup> (53.8-94.5)	64.8 <sup>b,m,x</sup> (36.9-77.6)	79.4 <sup>a,b,z</sup> (51.7-105.6)	68.6 (59.5-76.0)	71.4 <sup>m</sup> (51.5-116.4)	76.2 (49.1-105.3)
External pudendal artery diametre (cm)	0.54 <sup>a</sup> (0.45-0.64)	0.58 <sup>a,b,m,x</sup> (0.50-0.79)	0.55 <sup>b</sup> (0.49-0.63)	0.54 (0.48-0.73)	0.56 <sup>m,x</sup> (0.47-0.62)	0.53 (0.47-0.70)
Resistance index	0.62 <sup>a,m</sup> (0.54-0.91)	0.54 <sup>a,b</sup> (0.47-0.66)	0.63 <sup>b,z</sup> (0.53-0.84)	0.61 <sup>a,b,m</sup> (0.51-0.68)	0.52 <sup>a,c,z</sup> (0.45-0.73)	0.64 <sup>b,c</sup> (0.54-0.89)
Pulsatility index	1.16 <sup>a,x,z</sup> (0.76- 1.41)	0.92 <sup>a,b,z</sup> (0.68-1.23)	1.06 <sup>b,z</sup> (0.69-2.01)	1.12 <sup>a,b</sup> (0.85-1.33)	0.94 <sup>a,c</sup> (0.58-1.46)	1.20 <sup>b,c</sup> (0.86-2.08)
Systolic:diastolic velocity ratio	2.61 <sup>z</sup> (2.10-3.26)	2.76 <sup>m,z</sup> (2.21-3.16)	2.60 <sup>z</sup> (2.22-2.35)	2.60 (2.47-3.12)	2.66 <sup>m</sup> (1.89-3.76)	2.67 (2.01-3.55)
General pressure (mm Hg)	3.50 <sup>a</sup> (2.50-7.50)	4.40 <sup>z</sup> (2.00-7.50)	3.40 <sup>a,x,z</sup> (2.30-5.90)	4.00 (2.00-7.50)	4.20 <sup>x,z</sup> (2.20-7.50)	3.80 <sup>z</sup> (2.10-7.50)
Mean pressure (mm Hg)	1.30 <sup>a,m,x,z</sup> (0.70-2.80)	1.60 <sup>a,b,z</sup> (0.90-3.60)	1.30 <sup>b,z</sup> (0.60-2.40)	1.40 <sup>m,x</sup> (0.80-4.20)	1.40 <sup>a,z</sup> (0.60-3.80)	1.30 <sup>a,z</sup> (0.60-3.20)
Mean velocity (m s <sup>-1</sup> )	0.51 <sup>a,z</sup> (0.42-0.58)	0.61 <sup>a,b,z</sup> (0.50-0.70)	0.54 <sup>b,z</sup> (0.36-0.79)	0.50 <sup>a,z</sup> (0.43-0.58)	0.55 <sup>a,x,z</sup> (0.45-0.85)	0.54 <sup>z</sup> (0.38-0.87)
Systolic acceleration (m s <sup>-2</sup> )	8.69 <sup>a</sup> (6.35-11.78)	12.12 <sup>a,b,x,z</sup> (7.14-15.47)	8.51 <sup>b,x,z</sup> (4.98-14.31)	9.05 <sup>a</sup> (4.78-11.78)	11.16 <sup>a,b,z</sup> (6.87-15.57)	8.68 <sup>b,z</sup> (6.48-11.27)
Blood input into parenchyma (mL min <sup>-1</sup> )	527 <sup>a,z</sup> (249-803)	709 <sup>a,b,m,x,z</sup> (333-998)	418 <sup>b</sup> (171-894)	523 <sup>a,z</sup> (271-888)	538 <sup>b,m,x,z</sup> (390-691)	321 <sup>a,b</sup> (168-868)
Blood input at teat (score 0-4)	0 <sup>a,b</sup> (0-1)	3 <sup>a,c,m,x</sup> (2-4)	1.5 <sup>b,c,m,x,z</sup> (0-2)	0 <sup>a,z</sup> (0-1)	0.5 <sup>a,b,m</sup> (0-2)	0 <sup>b,m,z</sup> (0-1)

**Table II.xii.** (continued).

	Before challenge	Day of challenge	After day of challenge	Before challenge	Day of challenge	After day of challenge
<b>Subgroup B2</b>	Ipsilateral to inoculated side of the udder			Contralateral side of the udder		
Mammary parenchyma grey-scale	65.8 <sup>a</sup> (56.9-73.6)	91.9 <sup>a,x</sup> (68.8-115.6)	79.2 (59.8-117.4)	66.2 (63.7-78.4)	85.5 <sup>a</sup> (52.5-85.6)	83.5 <sup>a</sup> (61.9-90.7)
External pudendal artery diametre (cm)	0.54 (0.51-0.59)	0.53 <sup>a,x</sup> (0.50-0.55)	0.56 <sup>a</sup> (0.51-0.64)	0.54 (0.49-0.63)	0.51 <sup>x</sup> (0.50-0.56)	0.55 (0.50-0.58)
Resistance index	0.60 (0.48-0.67)	0.64 (0.57-0.69)	0.62 (0.55-0.68)	0.60 (0.51-0.68)	0.64 (0.55-0.75)	0.63 (0.55-0.74)
Pulsatility index	1.13 <sup>x</sup> (0.95-1.18)	1.12 (0.87-1.27)	1.11 (0.82-1.34)	1.15 (0.94-1.29)	1.15 (0.95-1.55)	1.15 (0.92-1.64)
Systolic:diastolic velocity ratio	2.49 (1.91-3.00)	2.76 (2.33-3.19)	2.61 (2.21-3.12)	2.49 (2.10-3.12)	2.79 (2.21-3.93)	2.69 (2.23-3.09)
General pressure (mm Hg)	3.70 <sup>a</sup> (3.00-4.50)	5.00 (3.30-5.80)	5.00 <sup>a,x</sup> (3.40-5.80)	3.50 <sup>a,b</sup> (2.50-4.40)	5.40 <sup>a,x</sup> (4.50-6.00)	4.00 <sup>b</sup> (3.10-7.50)
Mean pressure (mm Hg)	1.40 <sup>a,b,x</sup> (1.10-1.60)	1.60 <sup>a</sup> (1.50-1.80)	1.30 <sup>b</sup> (1.00-1.70)	1.20 <sup>x</sup> (0.90-1.30)	1.30 (1.00-1.80)	1.30 (0.90-1.70)
Mean velocity (m s <sup>-1</sup> )	0.53 (0.47-0.59)	0.51 (0.50-0.60)	0.53 (0.47-0.62)	0.51 (0.43-0.54)	0.50 <sup>x</sup> (0.49-0.55)	0.51 (0.45-0.56)
Systolic acceleration (m s <sup>-2</sup> )	9.98 (6.23-15.71)	8.43 <sup>a,x</sup> (7.15-9.51)	11.17 <sup>a,x</sup> (5.71-15.57)	9.46 (6.40-13.44)	9.21 (8.03-15.47)	9.17 (7.50-13.28)
Blood input into parenchyma (mL min <sup>-1</sup> )	539 <sup>a,b</sup> (353-954)	383 <sup>a,x</sup> (367-506)	388 <sup>b</sup> (326-515)	519 <sup>a,b</sup> (444-878)	376 <sup>a,x</sup> (341-394)	326 (277-478)
Blood input at teat (score 0-4)	0 (0-2)	0 <sup>x</sup> (0-1)	0 <sup>x</sup> (0-3)	0 (0-1)	0 (0-1)	0 (0-2)

Before challenge: D-2 and D0; Day of challenge: D0+3 h, D0+6 h, D0+12 h (for subgroups A1 and B1) – D0+12 h (for subgroups A2 and B2); After day of challenge: D1, D2, D3 and D4 (D0: day of challenge).

A1, A2, B1, B2 = subgroups (A\*: ewes with increased  $\beta$ -hydroxybuturate blood concentration during the last stage of the preceding pregnancy, B\*: ewes with no increased  $\beta$ -hydroxybuturate blood concentration during the last stage of the preceding pregnancy; \*1: ewes challenged with deposition of *M. haemolytica* at the teat duct on the 5<sup>th</sup> day post-partum, \*2: uninfected control ewes).

<sup>a</sup>, <sup>b</sup>, <sup>c</sup>: significant difference between time period within parametre, subgroup and gland; <sup>m</sup>: significant difference between mammary glands within parametre, subgroup and time period; <sup>x</sup>: significant difference between \*1/\*2 subgroups within parametre, mammary gland and time period; <sup>z</sup>: significant difference between A\*/B\* subgroups within parametre, mammary gland and time period ( $P < 0.05$ ).

**Table II.xiii.** Quantitative results (median) of haemodynamic parameters in the external pudendal artery in the udder in ewes with one teat inoculated with *M. haemolytica*; results of the two sides of the udder are considered separately.

	D0	D0+3 h	D0+6 h	D0+12 h	D0	D0+3 h	D0+6 h	D0+12 h
<b>Subgroup A1</b>	<b>Inoculated glands</b>				<b>Contralateral glands</b>			
External pudendal artery diameter (cm)	0.55	0.55	0.58 <sup>z</sup>	0.57	0.55	0.57	0.59	0.56
Resistance index	0.61 <sup>m</sup>	0.57	0.52 <sup>m</sup>	0.55	0.58 <sup>m</sup>	0.59	0.56 <sup>m,z</sup>	0.56
Pulsatility index	1.02 <sup>z</sup>	0.93	0.80 <sup>m,z</sup>	0.86 <sup>m</sup>	1.00 <sup>z</sup>	0.95	0.88 <sup>m</sup>	0.91 <sup>m</sup>
Systolic:diastolic velocity ratio	2.43	2.51 <sup>z</sup>	2.55 <sup>m</sup>	2.48 <sup>z</sup>	2.32	2.40	2.27 <sup>m,z</sup>	2.32
General pressure (mm Hg)	4.4	4.7	7.1 <sup>z</sup>	7.5 <sup>z</sup>	4.7	4.7	7.2 <sup>z</sup>	7.5 <sup>z</sup>
Mean pressure (mm Hg)	1.9	2.1	3.3 <sup>z</sup>	3.1 <sup>z</sup>	1.8	2.2	3.0 <sup>z</sup>	3.4 <sup>z</sup>
Mean velocity (m s <sup>-1</sup> )	0.63	0.72 <sup>z</sup>	0.79 <sup>m,z</sup>	0.85 <sup>z</sup>	0.61	0.70 <sup>z</sup>	0.70 <sup>m,z</sup>	0.74 <sup>z</sup>
Systolic acceleration (m s <sup>-2</sup> )	7.75	7.94 <sup>m,z</sup>	10.61	10.76	9.54	9.16 <sup>m</sup>	9.54 <sup>z</sup>	9.41
Blood input into parenchyma (mL min <sup>-1</sup> )	357	435	568 <sup>z</sup>	536	338	441	467 <sup>z</sup>	454
Blood flow at teat (score 0-4)	0	2 <sup>m,z</sup>	3 <sup>m,z</sup>	2 <sup>m,z</sup>	0	1.5 <sup>m,z</sup>	1 <sup>m</sup>	0.5 <sup>m</sup>
<b>Subgroup B1</b>	<b>Inoculated glands</b>				<b>Contralateral glands</b>			
External pudendal artery diameter (cm)	0.55	0.58	0.65 <sup>m,z</sup>	0.53	0.54	0.55	0.58 <sup>m</sup>	0.55
Resistance index	0.62	0.59	0.49	0.56	0.62	0.57	0.50 <sup>z</sup>	0.52
Pulsatility index	1.03 <sup>z</sup>	0.96	0.90 <sup>z</sup>	0.93	1.16 <sup>z</sup>	0.96	0.77	0.96
Systolic:diastolic velocity ratio	2.54	2.85 <sup>z</sup>	2.75	2.75 <sup>z</sup>	2.65	2.62	2.67 <sup>z</sup>	2.57
General pressure (mm Hg)	3.6	4.0	5.4 <sup>z</sup>	4.2 <sup>z</sup>	4.1	4.5	4.1 <sup>z</sup>	4.1 <sup>z</sup>
Mean pressure (mm Hg)	1.3	1.6	1.8 <sup>z</sup>	1.6 <sup>z</sup>	1.4	1.6	1.4 <sup>z</sup>	1.4 <sup>z</sup>
Mean velocity (m s <sup>-1</sup> )	0.53	0.60 <sup>m,z</sup>	0.61 <sup>m,z</sup>	0.62 <sup>z</sup>	0.50	0.53 <sup>m,z</sup>	0.55 <sup>m,z</sup>	0.60 <sup>z</sup>
Systolic acceleration (m s <sup>-2</sup> )	8.84	10.95 <sup>z</sup>	13.20	10.98	9.43	11.59	12.16 <sup>z</sup>	10.51
Blood input into parenchyma (mL min <sup>-1</sup> )	476	596	737 <sup>m,z</sup>	759	477	500	545 <sup>m,z</sup>	544
Blood flow at teat (score 0-4)	0	2 <sup>m,z</sup>	3 <sup>m,z</sup>	3 <sup>m,z</sup>	0	0 <sup>m</sup>	1 <sup>m</sup>	0.5 <sup>m</sup>

Before challenge: D0; After challenge: D0+3 h, D0+6 h, D0+12 h.

A1 subgroup: ewes with increased  $\beta$ -hydroxybuturate blood concentration during the last stage of the preceding pregnancy, B1 subgroup: ewes with no increased  $\beta$ -hydroxybuturate blood concentration during the last stage of the preceding pregnancy.

<sup>m</sup>: significant difference between mammary glands within parametre, subgroup and sampling point; <sup>z</sup>: significant difference between subgroups within parametre, mammary gland and sampling point ( $P < 0.05$ ).

## Pathological findings after challenge of the experimental ewes

### *Inoculated side of the udder*

Measurement of the length of the internal teat structures after dissection of the teats of ewes in subgroups A1/B1 showed that the inoculum had always been deposited within the teat duct.

In subgroup A1 ewes, the inoculated teats had a thickened internal lining, often hyperaemic or petechiated; the teat duct and the teat cistern were distinguishable as two separate anatomical structures. Histologically, there was leucocytic infiltration (neutrophils, lymphocytes, plasma cells) observed subepithelially in the teat. The characteristic area of hyperplastic lymphoid tissue at the border between teat duct and teat cistern, with lymphocytes and plasma cells, was observed in 3/8 ewes (clinical mastitis developed in ewes in which these structures were not evident). In the remaining animals, accumulation of lymphocytes was evident subepithelially in the teat, but with no formation of lymphoid follicles (Fig. II.17.). In the mammary parenchyma, massive leucocytic infiltration was seen histologically; furthermore, intra-alveolar live and exhausted neutrophils, extravasation and destruction of epithelial cells and alveoli were evident. In ewes, in which clinical mastitis had developed, conspicuous haemorrhage was also noted; in ewes, from which samples were collected at later stages of the study (D3, D4), lymphocytes were also observed therein.

**Figure II.17.** Presence of high numbers of lymphocytes, subepithelially in the area between teat duct-teat cistern, but with no formation of lymphoid follicle (low magnification 10× objective, photograph taken on a Zeiss Axiostar photomicroscope III) (ewe with increased  $\beta$ -hydroxybuturate blood concentration during the last stage of the preceding pregnancy, subsequently challenged into the teat duct with *M. haemolytica* isolate VSM08L [subgroup A1] and euthanised on D3).




In subgroup A2, mild macroscopic lesions were evident in two teats. Histologically, leucocytic infiltration was recorded occasionally. The characteristic area of hyperplastic lymphoid tissue at the border between teat duct and teat cistern was observed in 1/2 ewes with bacterial isolation from tissue samples (in the other ewe, subclinical mastitis was noted). Neutrophils were seen in high numbers inter- and intra-alveolarly in one ewe, with destruction of epithelial cells also evident at some parts of the sample.

In subgroup B1, occasionally the inoculated teats of ewes had a hyperaemic internal lining, with folds, but no thickening, on the mucosa; the teat duct and the teat cistern were distinguishable as two separate anatomical structures. Histologically, there was marked leucocytic infiltration



(neutrophils, lymphocytes, plasma cells) in the teat. The characteristic area of hyperplastic lymphoid tissue at the border between teat duct and teat cistern, with lymphocytes and plasma cells (Fig. II.18.), was observed in 7/8 ewes ( $P=0.019$ , compared to respective proportion in subgroup A1). In the mammary parenchyma, leucocytic infiltration (neutrophils and lymphocytes), lysis of neutrophils, extravasation and destruction of epithelial cells were evident.

**Figure II.18.** Hyperplastic lymphoid follicle consisting of lymphocytes and plasma cells at the border between the teat duct and teat cistern (low magnification 10× objective, photograph taken on a Zeiss Axiostar photomicroscope III) (ewe with no increased  $\beta$ -hydroxybuturate blood concentration during the last stage of the preceding pregnancy, subsequently challenged into the teat duct with *M. haemolytica* isolate VSM08L [subgroup B1] and euthanised on D3); inset presents the follicle at larger magnification (magnification 100× objective).



In subgroup B2, the inoculated teats had no abnormal gross or histological findings. No abnormal findings were also seen in the mammary parenchyma.

Presence of increased  $\beta$ -hydroxybuturate blood concentration during the last stage of the preceding pregnancy was associated with increased bacteriological isolations from tissue samples (52% of samples from ewes with increased  $\beta$ -hydroxybuturate blood concentration compared to 28% of samples from ewes with no increased  $\beta$ -hydroxybuturate blood concentration,  $P=0.025$ ). More bacterial isolations were also achieved from samples from the inoculated side of challenged ewes than from uninoculated animals (60% and 17%, respectively,  $P<0.001$ ). When comparisons were performed between the various subgroups, bacteria were recovered more frequently from A1 than from B1 subgroup ( $P=0.008$ ) and from A2 than from B2 subgroup ( $P=0.058$ ). Details of isolation of *M. haemolytica* from tissue samples are in Table II.xiv.

In subgroup A1, cumulative score for macroscopic and histological pathological findings in the inoculated teat was 18 and 23, respectively; cumulative score for histopathological findings in the mammary parenchyma ipsilaterally to the inoculated teat was 24. Respective figures for subgroup A2 were 2, 9 and 5. Respective figures for subgroup B1 were 5, 31 and 16 ( $P=0.025$ ,  $P=0.05$  and  $P=0.01$ , respectively, when compared to results in subgroup A1) and for subgroup B2 were 0 in all cases ( $P>0.05$ , when compared to results in subgroup A2). In all above cases, maximum possible score was 32.

There was evidence that scores for teat macroscopic lesions and for parenchyma histological lesions in inoculated side of ewes with increased  $\beta$ -hydroxybuturate blood concentration during the preceding pregnancy (i.e., group A) were significantly different than scores in the inoculated side of ewes with no increased  $\beta$ -hydroxybuturate blood concentration (i.e., group B) ( $P=0.02$  and  $P<0.005$ , respectively), whilst scores for teat histological lesions did not differ significantly between the two groups ( $P>0.05$ ). Moreover, all scores (teat macroscopic or histological lesions, parenchyma histological lesions) differed significantly between inoculated ewes (i.e., subgroups A1 and B1) or not (i.e., subgroups A2 and B2) (in all cases,  $P<0.005$ ).

Detailed results of pathology scoring are in Table II.xv.

**Table II.xiv.** Isolation of *M. haemolytica* from udder samples from ewes with one teat inoculated or not with the organism.

	D1				D2				D3				D4			
	A1	A2	B1	B2	A1	A2	B1	B2	A1	A2	B1	B2	A1	A2	B1	B2
Site from where isolated																
Teat duct	2/2	0/2	2/2	0/1	2/2	0/2	2/2	0/1	2/2	1/2	2/2	0/1	1/2	1/2	1/2	0/1
Teat cistern	1/2	0/2	0/2	0/1	2/2	0/2	1/2	0/1	2/2	1/2	1/2	0/1	2/2	1/2	1/2	0/1
Mammary parenchyma	0/2	0/2	0/2	0/1	2/2	0/2	0/2	0/1	2/2	1/2	0/2	0/1	1/2	1/2	0/2	0/1

D1, D2, D3, D4 = days after inoculation

A1, A2, B1, B2 = subgroups (A\*: ewes with increased  $\beta$ -hydroxybuturate blood concentration during the last stage of the preceding pregnancy, B\*: ewes with no increased  $\beta$ -hydroxybuturate blood concentration during the last stage of the preceding pregnancy; \*1: ewes challenged with deposition of *M. haemolytica* at the teat duct on the 5<sup>th</sup> day post-partum, \*2: uninfected control ewes).

n/m=positive results out of total animals sampled



**Table II.xv.** Scores (median values) for pathology findings in the udder of ewes with one teat inoculated or not with *Mannheimia haemolytica*

	D1				D2				D3				D4			
	A1	A2	B1	B2	A1	A2	B1	B2	A1	A2	B1	B2	A1	A2	B1	B2
Score for macroscopic pathological findings in inoculated teat	0.5	0.0	0.5	0.0	3.0	0.0	1.0	0.0	3.0	1.0	1.0	0.0	2.5	0.0	0.0	0.0
Score for histopathological findings in inoculated teat	2.0	0.5	3.5	0.0	3.0	0.5	4.0	0.0	3.5	1.5	4.0	0.0	3.0	2.0	4.0	0.0
Score for histopathological findings in mammary parenchyma ipsilateral to inoculated teat	1.5	0.5	1.0	0.0	4.0	0.0	2.5	0.0	3.5	1.0	3.0	0.0	3.0	1.0	1.5	0.0

D1, D2, D3, D4 = days after inoculation

A1, A2, B1, B2 = subgroups (A\*: ewes with increased  $\beta$ -hydroxybuturate blood concentration during the last stage of the preceding pregnancy, B\*: ewes with no increased  $\beta$ -hydroxybuturate blood concentration during the last stage of the preceding pregnancy; \*1: ewes challenged with deposition of *M. haemolytica* at the teat duct on the 5<sup>th</sup> day post-partum, \*2: uninfected control ewes).

*Non-inoculated side of the udder*

In subgroup A1, *M. haemolytica* was isolated from the tissue sample from one ewe. Examination revealed diffuse neutrophilic infiltration in the mammary parenchyma (score 2). The sample was collected from the ewe from which bacteria had been isolated also from milk samples before she had been removed from the study.

All other mammary parenchyma tissue samples from the contralateral side of the experimental ewes did not yield bacteria, nor were evaluated with score higher than zero. All statistical comparisons did not reveal any significance ( $P>0.3$ ).

Scores for histopathological findings in mammary parenchyma ipsilaterally to inoculated teats (i.e., subgroups A1 and B1) were significantly higher than those for findings in contralateral side of these animals ( $P=0.02$ ). In contrast, respective scores in subgroups A2 and B2 were not significant ( $P>0.05$ ).

# GENERAL DISCUSSION

## Introduction

In sheep, the last stage of pregnancy is a metabolically demanding period, hence energy requirements of pregnant ewes increase, as end of gestation is approaching. Peri-parturient relaxation of immunity occurs in many mammalian species (Houdijk et al. 2001). In sheep, the relaxation of immunity around the lambing period (during the last month of gestation and the first fortnight of lactation period) has been studied initially in the 1980s, with a view to explain the peri-parturient rise in eggs of nematode helminthes in faeces of ewes (Armour 1980, Gibbs 1986, Barger 1993). This had been initially associated with increased prolactin concentrations. Fleming (1993, 1996) investigated the potential role of prolactin and prostaglandin  $F_{2\alpha}$  (two hormones, the concentrations of which increase at the end of pregnancy) and has found that total egg production by *Haemonchus contortus* in infected sheep increased after administration of prolactin, but not after administration of prostaglandin  $F_{2\alpha}$ , to those animals.

Prolactin is a peptide hormone, responsible for initiating and sustaining lactation (Castro et al. 2011). It acts in a cytokine-like manner and as an important regulator of the immune system (Rovensky et al. 1991). Blood concentrations of prolactin in pregnant ewes start to increase after the 115th to 135th day of gestation (Banchero et al. 2006). A theory had thus been developed regarding a potential role of the increased prolactin blood concentrations present at the end of pregnancy and the beginning of a lactation period to explain reduced immunocompetence during those periods. However, more recently, Beasley et al. (2010a, 2010b) have reported that, in ewes experimentally infected with *Trichostrongylus colubriformis*, the rise in faecal egg counts at the end of pregnancy had been preceded by a decrease in the immunological competence of the ewes, as shown by reduced numbers of circulating eosinophils and by decreased total antibody and IgG<sub>1</sub> titres; changes coincided with increased prolactin concentrations at the end of pregnancy, but, nevertheless, the authors conceded that they were unrelated to hormonal effects; this confirmed a similar earlier hypothesis by Coop et al. (1990).

In contrast, Coop and Kyriazakis (1999) have reviewed evidence supporting that the relaxation of immunity appeared to have a nutrition-based background and proposed a nutritional, rather than an endocrinological, involvement in the relaxation of immunity during that period, by presenting the following arguments: (i) grade of immunity expression in pregnant ewes was consistent with the reproductive effort, i.e., the number of fetuses borne, (ii) termination of pregnancy led to abrupt restoration of immunity and (iii) nutritional management of pregnant ewes could alter the time of first observation of the relaxation. Under conditions of high metabolic demand, which occur at the end of pregnancy, susceptibility of ewes to gastrointestinal parasites

was found to be increased (Kahn et al. 2003). Further, Beasley et al. (2012) have reported that ewes given a low energy diet had higher faecal epg counts than ewes fed a high quality diet.

There is little work to study potential effects of the reduced immunocompetence of ewes around the lambing period in microbial infections (Caroprese et al. 2015). Post-partum genital infections in ewes are of limited significance, due to their small incidence risk. Hence, there has been little scope to explore the idea of peri-parturient relaxation of immunity in association with metritis. In contrast, there has been shown an increased incidence risk of mastitis in the immediately post-partum period, which is followed by a subsequent reduction in the frequency of the disease (Jones and Lanyon 1987, Fthenakis and Jones 1990a). This indicates a potential contributory factor of the peri-parturient relaxation of immunity in the pathogenesis of the disease.

## **Definitions regarding pregnancy toxemia**

Subclinical disease is an illness that does not show clinical signs and cannot be recognised based on clinical evidence alone. A subclinical problem may be the early stage of a clinical disease or, alternatively, a very mild form of an established disease that does not produce clinical signs. Nevertheless, subclinical diseases are of importance, because they adversely influence the productivity of affected animals; for example, in sheep, gastrointestinal trichostrongylid parasitic infections cause a significant adverse effect in milk production or growth of parasitised animals (Fthenakis et al. 2015), although affected animals do not always show clinical signs relevant to those infections. Further, subclinical diseases may predispose animals to increased susceptibility to other diseases; for example, in sheep, *Mycoplasma ovipneumoniae* or *Parainfluenza virus* type 3 may cause subclinical respiratory infections, which predispose affected animals to severe *Mannheimia haemolytica*-associated pneumonia (Sharp and Nettleton 2007, Nicholas et al. 2008).

A subclinical infection can be identified by the detection of the causal organism (or its genetic material or its progeny) by using the appropriate laboratory technique, e.g., detection of *Bluetongue Virus* RNA in blood samples from sheep with no relevant clinical signs indicates subclinical disease. Nevertheless, infection may not always be equivalent to subclinical disease; for example, in sheep, diagnosis of subclinical mastitis is based upon simultaneous isolation of bacteria at increased numbers (usually >10 colony-forming-units in a standard 0.05 mL sample of milk) coupled with detection of increased somatic cell counts in milk, which are indicative of the animal's inflammatory response (Fragkou et al. 2014). It thus becomes evident that identification of a subclinical infection is based on tangible criteria, which include identification of the causative

organism (or its genetic material, e.g. viral RNA, or its progeny, e.g., eggs or larvae of parasites) and/or detection of the consequences of infection (e.g., inflammatory reaction).

In contrast, there may be some hinderance in the identification of subclinical problems of metabolic nature. In such cases, identification might be based in the results of biochemical tests in, for example, blood samples, which indicate that findings regarding specific parametres are outside the respective reference range. In such cases, there may be a question of validity of the findings, given that reference ranges of biochemical parametres can be influenced by a variety of factors, e.g., age, reproductive status or even breed of animals (Roger 2008). This limitation may be overcome by employing tests, which confirm changes in the function of the organism. For example, in sheep, the results of measurement of glutathione peroxidase enzymes GPx1, GPx2, GPx3 and GPx4 blood concentrations indicate the improper metabolism of selenium, hence this can be a more accurate test for diagnosis of selenium deficiency than measurement of selenium blood concentrations (Anderson et al. 1979, Jelinek et al. 1985); further, in dogs, the results of the glucose tolerance test indicate the inability of an animal to properly metabolise glucose, hence this can be a more accurate test for diagnosis of diabetes than measurement of glucose blood concentrations (Wang et al. 2013).

The term 'subclinical pregnancy toxemia' has been used infrequently, but increasingly in recent years (Ismail et al. 2008, Balikci et al. 2009, Gurdogan et al. 2014, Temizel et al. 2015). Nevertheless, not all those authors have provided adequate evidence of, additionally to increased  $\beta$ -hydroxybutyrate blood concentrations, presence of changes indicative of functional alterations in the affected animals. Concurrently with increased  $\beta$ -hydroxybutyrate blood concentrations, only Balikci et al. (2009) and Gurdogan et al. (2014) have indicated increased concentrations of the enzymes aspartate transaminase and alanine transaminase and of acute phase proteins (haptoglobin and serum amyloid A), respectively, in the blood of ewes with 'subclinical pregnancy toxemia'. These findings indicate a functional disorder (liver damage and stress), as the result of reduced energy intake during the final stage of gestation, and confirmed that animals were indeed affected, that way supporting use of the term 'subclinical pregnancy toxemia'. Possibly, other parametres might also be included in the definition of the disorder, e.g. blood concentration of non-esterified fatty-acids.

There may be however an entirely different approach, taken from a purely linguistic viewpoint. The word 'toxemia' derives from the Greek word '*τοξιναιμία*', meaning 'toxins in the blood', in this context ketone bodies (primarily  $\beta$ -hydroxybutyrate and in smaller quantities acetoacetate). Hence, any animal with increased blood concentration of  $\beta$ -hydroxybutyrate (independently of presence of clinical signs) would be considered with pregnancy toxemia and

should be further characterised 'with relevant clinical signs' or 'with no clinical signs'. This approach does not involve use of the term 'subclinical pregnancy toxemia', because the disorder is characterised based only on the increased blood concentration of  $\beta$ -hydroxybutyrate, as per its name.

Nevertheless, in both cases, the  $\beta$ -hydroxybutyrate blood concentration threshold defining the condition ('subclinical pregnancy toxemia' or 'pregnancy toxemia') remains undetermined. Currently, the value of  $1.2 \text{ mmol L}^{-1}$  is the threshold indicating ewes at risk to develop clinical signs associated with the disease; this has been defined some years ago and is still being used in establishing measures to prevent clinical cases of the disease in a flock (Sargison 2008, Brozos et al. 2011). However, it has not been validated as a threshold defining 'subclinical pregnancy toxemia', although many authors have used it as such to define that condition.

In their works, Balikci et al. (2009) and Gurdogan et al. (2014) have reported that the value of  $1.2 \text{ mmol L}^{-1}$  was the one above which concentrations of liver function-related enzymes and acute phase proteins, respectively, were found increased. This indicates that when values of  $\beta$ -hydroxybutyrate blood concentration are above that threshold, functional changes occur in the animal. It thus seems that the above value can be a possible measure for defining 'subclinical pregnancy toxemia' or 'pregnancy toxemia with no clinical signs'.

In relevant reports dealing with ketosis in cows, values in the range of  $1.1 \text{ mmol L}^{-1}$  to  $1.6 \text{ mmol L}^{-1}$  have been used to define 'subclinical ketosis', although results may be influenced by a variety of factors, e.g., the measurement technique, the vein from which sample was collected, the blood fraction (whole blood, serum, plasma) or even the ambient temperature (Kanz et al. 2015, McCarthy et al. 2015, Sun et al. 2015).

Although there are significant differences between sheep and cattle in genetic diversity, physiology and management practices, there is some scope in taking into account the findings from cows, given the scarcity of results from work performed specifically in ewes. The few results of work performed in ewes allied to the findings in cows support use of  $1.2 \text{ mmol L}^{-1}$  as threshold to define 'subclinical pregnancy toxemia' or 'pregnancy toxemia with no clinical signs'. However, given the importance of the issue and the significant variance that has been reported in reference values in sheep for blood biochemical parameters, further work would be necessary before a widely accepted threshold value could be fully established.

For the purposes of this discussion, the term 'pregnancy toxemia' is used to denote increased (i.e.,  $>1.2 \text{ mmol L}^{-1}$ )  $\beta$ -hydroxybutyrate blood concentration in ewes at the end of pregnancy. The terms 'subclinical pregnancy toxemia' and 'pregnancy toxemia with no clinical signs' are used interchangeably, as in reality there is no difference between them. The term

'clinical pregnancy toxaemia' is not employed, as the term 'pregnancy toxaemia with relevant clinical signs' has been considered more appropriate.

## **The experimental model used for induction of pregnancy toxaemia**

The model used for induction of pregnancy toxaemia was based in providing to animals a feed with reduced energy content during the last one-a-half months of pregnancy. The results indicate that the model was successful for induction of the disease. Its advantage is the fulfillment of satiation requirements of animals. Moreover, the adverse circumstances were introduced progressively, which resembles closely to development of the pathological condition in the field; that way, consequences of the problem would be similar to those occurring under clinical circumstances. In recent papers, Cal-Pereyra et al. (2015a, 2015b) have described a model for induction of pregnancy toxaemia, in which animals had been completely starved for a week; that approach would not be acceptable on welfare grounds. Moreover, parasitism of animals by gastrointestinal trichostrongylids, which reduce energy availability in sheep, has contributed to successful development of the model.

The model is characterised by repeatability, as in a subsequent study (not part of this thesis) the disease has been successfully reproduced (Ioannidi et al. 2015). The disease would be more severe in cases of heavy parasitic infections, as parasites increase the energy requirements of their hosts (Coop et al. 1977, Dakkak 1990), that way parasitism directly predisposing to pregnancy toxaemia (Papadopoulos et al. 2013).

Of the animals that were provided with the experimental diet, 57% were found with increased blood concentrations of  $\beta$ -hydroxybutyrate during the last 15 days of pregnancy. Repeat detection of blood concentrations of  $\beta$ -hydroxybutyrate  $>1.2$  mmol L<sup>-1</sup> was used to consider that animals had developed the disease, as  $\beta$ -hydroxybutyrate blood concentration may fluctuate during the last stage of gestation, with intermittent increases of the parametre as the result of transient effects (e.g., temporary anorexia of an animal). At the end, only 25% of ewes with pregnancy toxaemia (14% of all ewes into the study) developed relevant clinical signs.

There were differences between animals in their response to reduced energy administration during the last stage of pregnancy. Ewes bearing two or more foetuses were more likely to develop pregnancy toxaemia; increased number of foetuses is an established risk factor for the disease. From the viewpoint of development of the model, one may obtain a higher proportion of animals with the disease if one uses a more prolific breed of sheep. Alternatively,



administration of equine chorionic gonadotrophin at varying doses (200-500 IU, depending on season, latitude and breed of animals) will lead to increased ovulation rates (Abecia et al. 2011, 2012), therefore increased proportion of animals would develop the disease. Nevertheless, animals that would not develop disease may be used as potential controls in a planned experimental work.

The reverse correlation between  $\beta$ -hydroxybutyrate blood concentrations in pregnant females and birth bodyweights of lambs is in accord with the traditional view that ewes with pregnancy toxemia usually produce lambs with suboptimal bodyweight at birth (Andrews 1997). Moreover, the reverse correlation between faecal epg counts and birth bodyweight of lambs indicates the energy drain caused by the helminthes, which was reflected in reduced energy availability for foetuses, subsequently born with a suboptimal weight. In a previous study, Fthenakis et al. (2005) have highlighted the beneficial effects of anthelmintic administration during the last stage of pregnancy in the subsequent birth bodyweight of lambs of treated ewes. The results indicate the adverse effects of nematode helminthes in ewes during the last stage of pregnancy (i.e., pregnancy toxemia and reduced birth bodyweight of newborn lambs) and support the recommendation for anthelmintic treatment of ewes during the last stage of pregnancy (Fthenakis et al. 2012).

## **Doppler ultrasonographic examination of the foetus**

The results of ultrasonographic examination of the foetus indicated clearly that reduced volume of blood was entering into foetuses of ewes with pregnancy toxemia throughout the last stage of gestation. However, differences in other haemodynamic parameters were not significant, which does not support a hypothesis for a strong effect of pregnancy toxemia in the haemodynamic parameters (bar blood input) (unless results obtained on the last two examinations points would be considered together, i.e., the number of observations would be increased and also the statistical power of the study).

Nevertheless, findings in the present study are not in full accord with previous ones recorded in work performed in healthy individuals. Makikallio et al. (2006) have indicated that mean resistance index in the umbilical artery of sheep was 0.80 (on P130, results of present study: 0.68 and 0.68 for group A and B, respectively). Further, Galan et al. (2005) have reported that mean pulsatility rate was 0.94 (on P130, results of present study: 1.23 and 1.18 for group A and B, respectively) and mean systolic:diastolic velocity ratio was 2.28 (on P130, results of present study:

3.98 and 3.17 for group A and B, respectively). In this respect, it should be noted that, in general, there is little agreement between studies in results of haemodynamic parameters in sheep; for example, Locatelli et al. (2011), Dominguez et al. (2013), Poser et al. (2013) and Acorda and Pajas (2015) have all performed echocardiographic examination in sheep and have presented varying values for similar parameters; this has led researchers to suggest that in such studies comparisons should be made within the experiment (i.e., with control animals) and within the experimental animals (i.e., with values obtained before treatment) rather than with results of other studies. Different experimental conditions and/or methodological approaches may be decisive factors for these discrepancies between studies, although the animal species itself, given that over 400 breeds of domestic sheep (*Ovis aries*) are available worldwide, might also be of significance. To note that also proposed reference ranges for haematological values in healthy sheep differ among researchers (Kramer 2000, Martin and Aitken 2000, Roger 2008), indicating further discord in standardising results of parameters of the cardiovascular system in this species.

The present findings indicate the effort undertaken by ewes with pregnancy toxemia to increase blood input into the umbilical artery, in an attempt to provide fetuses with the necessary blood volume. Seemingly however, that compensatory mechanism was not fully successful. Although size of fetuses (as evidenced by results of measurement of abdominal diameter) was not found to be different between the two groups, birth bodyweights differed significantly between them, with lambs born from group B ewes being heavier than those from group A animals. The suboptimal birth bodyweight might reflect suboptimal growth of the heart (Han et al. 2004, Bubb et al. 2007), the lungs (Lipsett et al. 2006) and/or the pancreas (Dumortier et al. 2007). There is indirect evidence for a suboptimal growth of the heart, as changes in the pulsatility index or the resistance index in the umbilical artery have been found to be the result of significant changes in the heart of fetuses (Morrow et al. 1993, Olofsson et al. 2004), although vasoconstrictor drugs or placental embolisation (Adamson et al. 1990, Adamson 1999), as well as fetus-dependent factors (e.g., cardiac contractility, breathing movements, vessel elasticity) (Divon 1993), might have also accounted for changes in haemodynamic parameters. A potential suboptimal development of these organs might have led further to structural changes and impaired function in them, which would not have supported life or would have rendered newborns susceptible to infections (Gao et al. 2014). Further, hypoxia due to the reduced volume of blood transported into fetuses could have also been established. These factors would have contributed to increased mortality of fetuses at end of pregnancy and of newborn lambs soon after birth.

Embryos/fetuses are, in general, 'programmed' to live into the environment of their early life, to survive the neonatal period and to adopt a predictive adaptive response (Gluckman and

Hanson 2004, 2006). However, if pregnant females would be exposed to severe undernutrition, blood concentrations of growth factors might be affected and, consequently, intrauterine growth (Bloomfield et al. 2013); this adverse effect during intrauterine life may ensue various changes (often permanent) in the foetus and the offspring to be manifested after birth (Barker et al. 1989). Hea and Wua (2013) have indicated that effects of malnutrition of pregnant females during vital periods of intrauterine development might lead in reduction of growth of organs of embryos/foetuses borne by these females. Further, diet given to pregnant females might lead in epigenetic changes (DNA methylation) and modification of gene expression, resulting in changes in cell structure and function and, ultimately, in morphological and/or functional changes in tissues and/or organs of embryos/foetuses (Langley-Evans 2004). These may have effect(s) in various systems of newborns, including the immune system (Langley-Evans et al. 2003, Palmer 2011, Bloomfield et al. 2013). Merlot et al. (2008) have summarised the effects of stressful situations for pregnant females and indicated that such effects might have long-term consequences in the physiology and disease risk of their offspring; one may possibly include suboptimal nutrition of pregnant females as such a potentially stressful situation.

When a different allocation of ewes into groups was performed, with ewes into the study regrouped into animals that developed or did not develop dystocia, significant differences were evident in all haemodynamic parameters. By means of this transformation, it became evident that the statistical power of the study became strong enough and, even with a smaller number of observations, the findings indicated a clear difference between the two categories of ewes in the effect of pregnancy toxemia in the haemodynamic parameters. Obviously, animals which later developed dystocia, had increasing alterations in haemodynamics in the umbilical artery. In these animals, foetal and/or newborn death were also recorded, which indicates that in these ewes haemodynamic changes were so grave that could not be regulated by the pregnant animal.

In view of all the above, two points need further consideration. First, results of haemodynamic parameters proposed by previous workers (Galan et al. 2005, Makikallio et al. 2006) as reference values for haemodynamic parameters in the umbilical artery need to be reconsidered, especially in view of the general uncertainty in reference values in sheep (even for more commonly used parameters, e.g., haematological parameters), as pregnant ewes seem to develop compensatory mechanisms to counter-balance adverse effects. Second, results of Doppler examination of the umbilical artery in ewes with pregnancy toxemia may be of value in developing a predictive model for ewes to be considered at risk to developing dystocia and/or with a risk for death of their foetus/newborn lamb; perhaps, this technique may be used to establish

viability of foetuses, which would then guide towards the appropriate therapeutic strategy, i.e., caesarean section or induction of lambing (Brozos et al. 2011).

## **Ultrasonographic examination of the mammary glands during lactogenesis**

During the study, B-mode and Doppler ultrasonographic patterns of the mammary gland of ewes during lactogenesis have been described. The study presented relevant findings in healthy ewes, as well as in ewes with pregnancy toxemia. Changes in ultrasonographic appearance of the mammary parenchyma can reflect histological changes, which occur at the end of pregnancy, as the mammary gland prepares for the impending lactation.

Three fields were adequate to receive a representative ultrasonographic image of the mammary parenchyma. During the examination, anatomical structures of the mammary gland (as these did not change during the study) rather than dimensions (which changed as mammary development progressed) were used. Further, ultrasonographic examination by means of the Doppler technique was useful for measurement of the blood flow in the ovine mammary gland (Christensen et al. 1989, Piccione et al. 2004, Petridis et al. 2014).

The changes in intensity values of grey-scale appearance of ultrasonographic images of mammary parenchyma during the last stage of pregnancy indicated a progressively increasing echogenicity. This could be attributed to increasing numbers of mammary epithelial cells, compatible with the lactogenesis process taking place at that time. Remodelling of the extracellular matrix of the mammary gland takes place at that time, which results in increase of alveolar tissue and reduction of stromal tissue in the mammary parenchyma (Akers et al. 2006). These changes could be reflected in the intensity values of grey-scale imaging of the mammary tissue.

After lambing, a sharp reduction of echogenicity in the mammary parenchyma of ewes in both groups was observed. This, obviously, was the result of milk (a hypoechoic material) present in large volumes into the mammary alveoli, ductal system and cistern, as lactation started.

Presence of increased quantity of lacteal secretion in the mammary gland of healthy ewes could also be the reason for the reduced echogenicity of these mammary glands, compared to that in the mammary glands of ewes with pregnancy toxemia. This difference was significant and continued after lambing, when presence of increased quantity of milk in the mammary gland of healthy ewes (compared to that in ewes which had pregnancy toxemia) was further corroborated by the established greater milk quantity collected from healthy ewes. These findings are consistent with the documented smaller milk quantity of ewes with pregnancy toxemia at their subsequent

lactation period (Andrews 1997, Meyer et al. 2011), as efficient lactogenesis is dependent on appropriate nutrition during the last stage of pregnancy (Castro et al. 2011). Moreover, blood glucose concentrations have been associated with milk production (McNeill et al. 1998) and similar findings were evident in the present study, as group A ewes had significantly smaller blood glucose concentrations during the last stage of pregnancy and smaller milk yields immediately after lambing.

Along with the mammary parenchyma, the blood vessels of the gland also develop during the final stage of pregnancy. Increase of the diameter of the external pudendal artery further contributes to the greater amount of blood transferred to the mammary gland. Development of blood vessels in the udder is controlled by hormones, which stimulate growth of the mammary parenchyma, e.g. oestrogens and progesterone (Tucker 1981). This is further supported by the progressively decreasing blood concentrations of vasoconstrictor hormones (e.g., adrenaline, noradrenaline) during late pregnancy (Greiss 1972, Rosenfeld et al. 1976, Natrajan et al. 1982, Cvek 1997).

Doppler ultrasonographic indices and parameters also pointed out to the progressively increasing blood perfusion in the udder as lactogenesis advanced. During the last stage of pregnancy, blood is required for the synthesis of milk, as it carries precursors of milk constituents to the mammary gland and, hence, blood flow progressively increases as lactogenesis advances. Increasing blood volumes provide stronger signals for Doppler spectrum analysis, because of the greater number of moving erythrocytes (Widder and Goertler 2004).

The results have also indicated that blood input into the udder of ewes with pregnancy toxemia was significantly smaller than into the udder of healthy ewes. Increased blood input into the mammary gland has been found to lead to increased milk yield (Linzell 1960, 1974, Kronfeld et al. 1968, Davis and Collier 1985). Further, Lough et al. (1990) have identified increased energy provision with feed as a factor for increased mammary blood input. In the present study, the reverse aspect of the same phenomenon (i.e., reduced mammary blood flow consequently to reduced energy provision with feed) has been identified.

Smaller blood input (Doppler findings) in ewes, which had received reduced amount of energy during pregnancy, might have potentially resulted in smaller amount of epithelial cells in their mammary glands (B-mode findings) of these ewes. Further, smaller blood input may lead to reduced oxygen availability at alveolar level, with reduced oxygen transfer at mammary alveoli, as has been reported in lactating cows (Neary and Garry 2014); this may have further contributed to the reduced milk production by the experimental ewes.

Blood input appeared as the most accurate haemodynamic index for identification of differences between the two groups of animals (as was also the case in haemodynamic parameters in the umbilical artery). Low mean velocity of blood in large vessels has been correlated with increased blood input into the mammary gland of women (Madjar and Mendelson 2008), something also observed in healthy ewes at the end of pregnancy; it has been suggested that the finding indicates mammary glands with no potential abnormalities (Madjar and Mendelson 2008), which in the present study was confirmed by the lack of clinical abnormalities and bacterial isolations from milk samples. Pulsatility index also became of significance at the end of pregnancy, when differences in blood input into the udder further increased, indicating the increased cardiac output, hence increased blood flow to the udder of healthy ewes; low pulsatility index suggests compensatory distal vasodilatation following reduced volume of blood perfusion (Rajajee et al. 2012), as in the ewes with pregnancy toxemia. In fact, after lambing, approximately 15% of total cardiac output is directed to the udder (Linzell 1974, Olsson et al. 1998, Svennersten-Sjaunja and Olsson 2005). Increased blood demand in the udder during lactation (Braun and Forster 2012) puts a tremendous burden on and affects the entire circulatory system of the animal (Gürtler and Schweigert 2005). As start of lactation was imminent, in ewes with pregnancy toxemia, which had a smaller milk production, cardiac output (as shown in pulsatility rates just before lambing) was smaller than in healthy ewes. This becomes evident if the results of blood input into the mammary glands are allied to those of the blood input into the foetus. Possibly, reduction of antioxidant agents and increase of free radicals occurring in cases of hyperketonaemia (Jain et al. 1998, Sahoo et al. 2009) might lead to fatigue of the myocardial muscle (as shown by the increased blood concentration of cardiac troponin I; Tharwat et al. 2012), which in turn would result to decreased blood output. In this respect, Reksen et al. (2002) have found that in cows receiving high energy diets, blood flows were in general higher more increased in animals with smaller energy intakes. Finally, a case of cardiac failure has been described in a patient with ketosis (Diltoer et al. 2004).

Other haemodynamic indices take into account vascular effects, which are influenced by locally produced vasoregulators (Prosser et al. 1996). As there were differences in the amount of mammary tissue between ewes into group A or B, possibly there was also a different rate of production of these, which would have resulted in lack of significant differences in those indices between the two groups. In fact, some local (i.e., mammary) regulation of blood input into the udder has been reported by Cieslar et al. (2014).

## Peri-parturient problems in ewes with pregnancy toxaemia

The most common reasons for dystocia in ewes are incomplete dilatation of the cervix, malpresentation of the foetus(es) and foetal oversize (Hindson and Winter 2007). For example, Thomas (1990) reported that over 50% of dystocia cases in ewes were postural defects of the foetus, whilst Sobiraj (1994), Kloss et al. (2002) and Lopes Camara et al. (2009) diagnosed ringwomb as the main reason for dystocia. Uterine inertia is a rare cause of dystocia in ewes; the main stage of labour in that species is completed usually within 45 to 60 minutes and in any case within 2 hours (Noakes 1996).

In this study, appropriate treatment of the clinical signs relevant to pregnancy toxaemia had been initiated as soon as detected (Brozos et al. 2011). For treatment, propylene glycol was given *per os* and, occasionally, glucose was administered intravenously (Brozos et al. 2011). Hence, manifestations of the disease subsided and it has become possible to record clinical consequences of the disease at the peri-parturient period.

Ewes in group A (i.e., with pregnancy toxaemia) developed often expulsive deficiency, possibly as the consequence of uterine inertia. This might have been the effect of inadequate endocrinological mechanisms leading to parturition, due to the impaired stimulation of the foetal hypothalamo-pituitary-adrenocortical axis consequently to the undernutrition of the pregnant ewes (Edwards and McMillen 2002) and/or to the shorter duration of gestation recorded in the ewes with pregnancy toxaemia. Further, subclinical hypocalcaemia, which often occurs in association with pregnancy toxaemia (Brozos et al. 2011), as the result of feeding an incorrect ration, could have also led in uterine inertia;  $\text{Ca}^{2+}$  ions / calmodulin are important factors for regulating and promoting action of myosin light-chain kinase, which acts in the phosphorylation of myosine ATPase causing contraction of the myometrium (Noakes 1996). It is noteworthy that cows with fatty liver (also a feature in pregnancy toxaemia in ewes) have been found to have increased incidence of dystocia (Bobe et al. 2004).

Possibly, increased incidence of perinatal mortality in offspring of ewes with pregnancy toxaemia could have been the effect of reduced intrauterine growth, as the consequence of reduced energy availability in those ewes (group A), which was shown after lambing with the smaller birth bodyweights. Of course, reduced quantity of colostrum present in ewes which had developed pregnancy toxaemia (Andrews 1997), therefore inadequate intake by newborns, ultimately leading in inadequate protection of lambs, might have also contributed to neonatal mortality of lambs of these ewes.

One should also take into account that lambs born from ewes with pregnancy toxæmia had inadequate passive immunity, which depends on concentrations of immunoglobulins in colostrum (McGuirk and Collins 2004) and on absorptive capacity of the gut of newborns. Survival of newborns depends on quantity and quality of colostrum received during the first hours of life (Greenwood et al. 2000). In pregnant female animals, lactogenesis (which includes colostrogenesis) takes place for some weeks before parturition; hence, nutrition of females during the last stage of their pregnancy is important for quality and quantity of colostrum to be produced (Brandon et al. 1971a, 1971b, Miranda et al. 1983). Ewes that had received a correct nutritional regime during the final stage of pregnancy, have been found to produce greater quantities of colostrum (Mellor and Murray 1985, Swanson et al. 2008) with increased IgG content (Swanson et al. 2008) compared with malnourished animals. The ultrasonographic findings of the examination of the udder during pregnancy have supported a hypothesis that in ewes with subclinical pregnancy toxæmia smaller numbers of mammary epithelial cells had developed, which led in production of smaller amount of milk (including colostrum) during the lactation period.

Apart from immunoglobulins, colostrum also provides antioxidants (Przybylska et al. 2007), defence cells (including macrophages) and lipids and proteins that are easily oxidised and contribute to the function of various antibacterial systems. However, colostrum itself is a source of oxidation-reduction reactions (Kankofer and Lipko-Przybylska 2008) and the immunoglobulins contained therein are molecules with high susceptibility to peroxidation (Margiloff et al. 1998). In colostrum with increased IgG concentrations, antioxidants may be consumed in an attempt to protect the antibodies, which might explain the negative correlation between antioxidants and IgG found in colostrum (Abuelo et al. 2014). This could be allied to the findings of Kamada et al. (2007), who have indicated that colostrum supplementation with selenium (a confirmed antioxidant) increased IgG absorption. On the other hand, cortisol produced by the foetus, which plays a role in onset of parturition (Eliot et al. 1981), accounts for the increased absorptive capacity of the intestinal epithelium in the first hours of life of newborns (Bate et al. 1991). After birth, as blood concentrations of cortisol decrease, the absorptive capacity of the gut of newborns also decreases (Bate et al. 1991; Fowden et al. 1998). Hence, stressful situations before or during parturition might lead in irregular increases in glucocorticoid blood concentrations, which might play a role in modifying the duration of the absorptive capacity of the gut of newborns during early life, possibly resulting in reduced immunoglobulin absorption and consequent inadequate protection of the newborns, as might have occurred in this work.

Expulsive deficiency during the process of labour would have contributed to slow post-partum discharge of material from the uterus, leading to accumulation of fluids and debris into the



uterine cavity. This is a risk factor for development of metritis, as it would provide a substrate for significant bacterial growth (Noakes 1996). Further, the increased incidence of metritis could have been a result of reduced immunocompetence of ewes that had previously developed pregnancy toxemia (Lacetera et al. 2005), as the documented consequence of reduced energy availability, e.g., by causing decreased defensive activity of neutrophils (Sartorelli et al. 1999). Ketosis is also a confirmed risk factor for metritis in cows (Mee et al. 2014, Raboisson et al. 2014), whilst Suriyasathaporn et al. (2000) have presented an association between reduced energy balance and impaired leucocyte function in humans. Further, a direct role of  $\beta$ -hydroxybutyrate in leucocytic response in ketonaemic cows has been detected and can be responsible for the reduced defensive abilities of affected animals.

As leucocytes are also involved in the process of detachment of foetal membranes immediately after expulsion of the foetus, their reduced numbers or impaired function in ewes with pregnancy toxemia might have contributed to development of retention of foetal membranes in one ewe. The disorder has a really small incidence risk in ewes (Fthenakis et al. 2000). In cows, an association between reduced energy intake in late gestation and retention of foetal membranes has already been documented (Curtis et al. 1985, Gerloff et al. 1993, Kaneene et al. 1997). It is also noteworthy that Reid (1968) has mentioned that the metabolic disturbances associated with energy deficiency in pregnant ewes had effects in protein metabolism; this could in turn affect the enzymatic pathways leading to proteolysis of the cotyledons and consequently to retention of foetal membranes.

## **Association of pregnancy toxemia with mastitis during the subsequent lactation period**

The results confirmed that subclinical pregnancy toxemia could predispose ewes to mastitis in the immediately *post-partum* period. Previous field studies have described increased mastitis incidence in ewes in the immediately *post-partum* period (Jones and Lanyon 1987, Fthenakis and Jones 1990a), which had been attributed to increased stocking rates during that period. Although there may be some merit in that hypothesis, the possibility for a role of the peri-parturient reduction of immunity cannot be ruled out. To note that the hypothesis was proposed in the mid-1980's (Jones 1985), at which time the peri-parturient relaxation of immunity had not been adequately described. Increased sucking frequency by newborn lambs at that period (Gougoulis et

al. 2007) would contribute to increased chances of bacterial transmission during that period from the sucking lamb(s) to their dam(s) (Fragkou et al. 2011), which, coupled to the reduced defensive ability of the post-parturient ewe, would lead to increased mastitis incidence risk.

Deposition of *Mannheimia haemolytica* into the teat duct of healthy ewes has been found to result to subclinical mastitis or to expulsion of bacteria after a short period. Mavrogianni et al. (2005) and Fragkou et al. (2010) have documented the defensive ability of the teat and have attributed that to the lymphoid follicles present at the border between teat duct and teat cistern of ewes. In contrast, in cases with depleted local teat defences (e.g., *Orf virus* infection, chapped teats), clinical mastitis developed after deposition of bacteria at the same site (Mavrogianni et al. 2006b; Fragkou et al. 2011). Similar findings have been recorded in the present study, where deposition of *M. haemolytica* into the teat duct resulted in significantly more cases of mastitis (subgroup A1 ewes *versus* subgroup B1 ewes). Moreover, one case of spontaneous subclinical mastitis has been recorded in ewes with pregnancy toxemia that had not been challenged (subgroup A2); this indicates that the disease, likely, originated as the result of casual infection and depleted defensive abilities of the ewe occurring simultaneously.

In this study, the characteristic lymphoid structures at the border between teat duct – teat cistern have been observed in significantly less ewes with pregnancy toxemia (subgroup A1) than in ewes with no increased  $\beta$ -hydroxybutyrate blood concentrations (subgroup B1) after bacterial deposition into the teat. These structures have a confirmed protective role against pathogens invading the teat (Mavrogianni et al. 2005, Fragkou et al. 2010), whilst their absence or functional impairment may lead to development of mastitis (Mavrogianni et al. 2006b, Fragkou et al. 2007).

Pregnancy toxemia involves disturbances in liver metabolism and can affect hepatic function. In turn, this may affect defences of the ewes. Wentink et al. (1997) reported that cows with hepatic lesions had decreased ability of proliferation of lymphocytes, possibly as the result of changes in cell membranes based on the alterations of cellular metabolism. Further, Lacetera et al. (2005) found that such cows had a marked immunosuppression due to impaired lymphocyte function.

Increased blood concentrations of  $\beta$ -hydroxybutyrate have been found to impair the leucocytic response in the mammary gland, e.g., by causing an inhibitory effect on the respiratory burst activity of neutrophils, by impeding generation of superoxide anions (Hoeben et al. 1997) or by suppressing formation of extracellular traps, causing impaired pathogen killing (Grinberg et al. 2008). Further, increased ketone body blood concentrations have been found to reduce efficacy of lymphocytes in cows (Franklin et al. 1991), whilst lymphocytes of ketonaemic animals had reduced mitogenic responses than lymphocytes of healthy cows (Kandefer-Szerszen et al. 1992). These

findings can be allied to the impaired function of lymphoid follicles in the teat of the inoculated ewes.

Further studies have shown that increased concentrations of  $\beta$ -hydroxybutyrate might impede other facets of leucocytic response in cases of bacterial invasion into the mammary gland. Suriyasathaporn et al. (1999) reported that leucocytes from ketotic cows had lower chemotactic ability than those from healthy animals. Sartorelli et al. (1999), who worked specifically in ewes, have shown that increased concentrations of  $\beta$ -hydroxybutyrate could adversely affect particle uptake by neutrophils. This can support bacterial invasion into the mammary parenchyma and development of mastitic lesions as documented in our case.

Other mechanisms may also be implicated in the compromise of mammary defences. For example, work performed in samples from the experimental ewes (subgroup A1) has indicated modified concentrations of cytokines immediately after lambing (this work is not a part of the thesis). This can be another mechanism contributing in pregnancy toxaemia predisposing ewes to mastitis, as it has been well-documented that cytokine production is a significant defence mechanism of animals after intramammary infection (Sordillo and Babiuk 1991a, 1991b, Sordillo et al. 1991, Daley et al. 1993). Ketonaemic cows have been found with smaller amounts of interferons than normal animals (Kandefer-Szerszen et al. 1992), reduced quantities of which might inhibit animals to mount an effective defence response.

The observed increase of blood input into the mammary gland can be explained by the principles of inflammation. During pathogen invasion, there is a marked dilatation of arterioles, capillaries and venules, leading to transudation and hyperaemia (Meurer and Wolf 1999). Hence, an increased volume of blood moves into the mammary gland. This influx was first recorded 3 hours after deposition of bacteria into the teat duct and became maximum 3 hours later (i.e., 6 hours after initial challenge). In the literature, there are differences regarding timing of first increase of blood input into the mammary gland (Dhondt et al. 1977a, 1977b, Thivierge et al. 2000, Delamaire and Guinard-Flament 2006, Potapow 2010). However, the results are not always directly comparable, as differing antigenic stimuli (e.g., lipopolysaccharide, whole bacterial cells) or challenge sites (e.g., mammary parenchyma, mammary cistern, duct cistern) have been used by different workers. Further, varying ultrasonographic techniques have been used, which further increase difficulty in comparing results; for example, Potapow (2010) used transrectal ultrasonography to evaluate blood input into the pudendoepigastric trunk. Finally, objectives of previous works were directed more towards diagnosis of the disease, whilst interest in the present study was to elucidating its pathogenesis.

In ewes with pregnancy toxemia at the final stage of preceding pregnancy (subgroup A1), blood input after challenge was significantly less than in ewes in subgroup B1. It has been well documented that outcome of inflammation is dependent on blood influx into the organ where the process takes place (Granger and Senshenkova 2010, Kvietyš and Granger 2014). Specifically in the mammary gland, increase of blood input has been associated with pyrexia and local inflammation (Burvenich and Peeters 1980). Blood brings into the mammary gland inflammatory mediators, responsible for the initiation of the defensive response of the host (Sordillo and Streicher 2002). It also carries neutrophils, which play a prominent role in mammary defence during the initial stage of the infection. Therefore, the observed reduced blood input into the mammary gland of animals of subgroup A1 could have contributed to the impaired defence of the ewes and might have supported development of mastitis. Further, in animals of that subgroup, increased blood input in the teat was evident for a longer period than in ewes of subgroup B1; obviously, in the latter animals, as inflammation was under resolve, blood presence at the site of bacterial deposition decreased earlier than in animals of subgroup A1, where mastitis was evident in all ewes. Finally, increased blood input was also evident in the contralateral to the challenged side of the udder, albeit of smaller magnitude; the two external pudendal arteries branch off from the pudendoepigastric trunk, which in turn rises from the external iliac artery, therefore, blood from the external iliac artery and the pudendoepigastric trunk floods into the affected side, as well as, at the immediately post-challenge stage, also into the contralateral side.

In hypoglycaemic does (like in animals of subgroup A1), a reduced blood flow into the mammary gland has been recorded (Linzell 1967, 1974). Further, Perkins et al. (2002) have found that ketotic cows had slower heart rates than healthy animals after intramammary infusion of antigen; this led to a slower defence response and, ultimately, later and impaired recovery compared to healthy cows.

The smaller diameter of the external pudendal artery observed in ewes of subgroup A1 would have contributed to the reduced blood input. Diameter of arteries depends, among others, on contractile status of vascular smooth muscle, which, in turn, is influenced by the phosphorylation state of the regulatory myosin light-chain, ultimately responding to presence of increased amounts of  $Ca^{2+}$ /calmodulin (Aird 2007, Vanhoutte et al. 2009). Hence, the smaller increase in diameter immediately post-infection could have been the result of a subclinical hypocalcaemia, as discussed hereabove. There is a questionmark, as during lactogenesis no significant difference in diameter of the external pudendal artery was evident between group A and group B. However, during that period input of blood into the growing mammary parenchyma was

of much smaller amount than that observed during immediately post-challenge; hence, the vessel could not respond to the greatly increased blood requirements at that period.

As diameter of the pudendal artery was smaller, mean blood velocity had to increase, as observed in this study, in order accommodate as much as possible blood volume into the inflamed site. Rubin et al. (1995a, 1995b, 1997) have proposed a normalisation procedure to take account of that factor and have introduced a new measure, termed 'fractional moving blood volume', which estimates the volume of blood moving into a vessel, in dependence with the diameter of the vessel. Nevertheless, as discussed above, in this study, ewes that had pregnancy toxemia showed increased mean blood velocity, but not to the appropriate extent for blood input into the mammary gland.

The findings of the compromised teat immune defences and the results of Doppler ultrasonographic examination in blood input into the mammary gland indicate that the adverse effects of pregnancy toxemia in the general function of the affected ewes last longer than the end of pregnancy, as hitherto recognised.

## Epilogue

### Conclusions

The conclusions from the results of the present thesis are summarised herebelow.

(a) A model has been developed to induce pregnancy toxemia in ewes for use in experimental studies, at the same time covering satiation requirements of the animals. Parasitism might have further contributed to improving the efficacy of the model.

(b) Ultrasonographic examination of fetuses of ewes with pregnancy toxemia revealed reduced blood input in the umbilical artery.

➤ There was a particularly intense modification of haemodynamic properties in ewes which later developed dystocia.

(c) Ultrasonographic appearance of the mammary gland of ewes during lactogenesis has been described.

➤ Results of B-mode ultrasonographic examination indicated differences between healthy ewes and ewes with pregnancy toxemia in the development of mammary parenchyma.

➤ Smaller blood input into the udder of ewes with pregnancy toxemia was evident.

(d) Increased incidence of peri-parturient problems was recorded in ewes with pregnancy toxaemia; there was also increased perinatal mortality in their offspring.

(e) Pregnancy toxaemia can act as a potential predisposing factor for mastitis in the immediately *post-partum* period.

- Possibly, impairment of the lymphoid follicular structures present at the border between teat duct – teat cistern could have been the cause of reduced protection of the mammary gland.

## Prospects

Suggestions for further research, in continuation of the present work, are as below.

- The evaluation of Doppler ultrasonographic examination of the foetus(es) in ewes with pregnancy toxaemia to evaluate potential risk for peri-partum problems in the pregnant females, in order to establish a strategy for prevention.
- The elucidation of other mechanisms which may adversely affect the immune mechanisms of ewes affected with pregnancy toxaemia
- The extent (in terms of time after lambing) that the adverse effects of pregnancy toxaemia prevail in affected animals
- The formulation of management strategies to apply post-lambing in ewes affected with pregnancy toxaemia.

# REFERENCES

- Abecia JA, Forcada F, Gonzalez-Bulnes A (2011). Pharmaceutical control of reproduction in sheep and goats. *Veterinary Clinics of North America Food Animal Practice* 27:67-79.
- Abecia JA, Forcada F, Gonzalez-Bulnes A (2012). Hormonal control of reproduction in small ruminants. *Animal Reproduction Science* 130:173–179.
- Abuelo A, Perez-Santos M, Hernandez J, Castillo C (2014). Effect of colostrum redox balance on the oxidative status of calves during the first 3 months of life and their relationship with immune acquisition. *The veterinary Journal* 100:295-299.
- Acorda JA, Pajas AMGA (2015). M-mode echocardiographic values in male and female Philippine sheep (*Ovis aries*) (Artiodactyla: Bovidae) by age and status of lactation and pregnancy. *Philippine Journal of veterinary Medicine* 52:11-20.
- Adamson SL (1999). Arterial pressure, vascular input impedance, and resistance as determinants of pulsatile blood flow in the umbilical artery. *European Journal of Obstetrics and Gynecology* 84:119-125.
- Adamson SL, Morrow RJ, Langille BL, Bull SB, Ritchie JWK (1990). Site-dependent effects of increases in placental vascular resistance on the umbilical arterial velocity waveform in fetal sheep. *Ultrasound in Medicine & Biology* 16:19-27.
- Aird WC (2007). Phenotypic heterogeneity of the endothelium II. Representative vascular beds. *Circulation Research* 100:174-190.
- Akers RM, Capuco AV, Keys JE (2006). Mammary histology and alveolar cell differentiation during late gestation and early lactation in mammary tissue of beef and dairy heifers. *Livestock Science* 105:44-49.
- Ali A, Hayder M (2007). Ultrasonographic assessment of embryonic, fetal and placental development in Ossimi sheep. *Small Ruminant Research* 73:277-282.
- Anderson PH, Berrett S, Patterson DSP (1979). Biological selenium status of livestock in Britain as indicated by sheep erythrocyte glutathione-peroxidase activity. *The veterinary Record* 104:235-238.
- Anderson MS, He J, Flowers-Ziegler J, Devaskar SU, Hay WW Jr (2001). Effects of selective hyperglycemia and hyperinsulinemia on glucose transporters in fetal ovine skeletal muscle. *American Journal of Physiology* 281:R1256-R1263.
- Andrews A (1997). Pregnancy toxemia in the ewe. *In Practice* 19:306-312.
- Andrews AH (1998). Recombinant bovine somatotropin and propylene glycol following glucose injection in treating pregnancy toxemia. *Large Animal Practice* 19:31-33.
- Armour J (1980). Epidemiology of helminth disease in farm animals. *Veterinary Parasitology* 6:7-46.



- Association of Official Analytical Chemists (1995). *Official Methods of Analysis of AOAC International*, 16th edn. AOAC International, Arlington.
- Balikci E, Yildiz A, Gurdogan F (2009). Investigation of some biochemical and clinical parameters for pregnancy toxemia in Akkaraman ewes. *Journal of animal and veterinary Advances* 8:1268-1273.
- Banchero GE, Quintans G, Martin A, Lindsay DR, Milton JTB (2004). Nutrition and colostrum production in sheep. 1. Metabolic and hormonal responses to a high-energy supplement in the final stages of pregnancy. *Reproduction Fertility and Development* 16:633-643.
- Banchero GE, Quintans G, Milton JTB, Lindsay DR (2002). Supplementation of Corriedale ewes with maize during the last week of pregnancy increases production of colostrum. *Animal Production in Australia* 24:273.
- Banchero GE, Perez Clariget R, Bencini R, Lindsay DR, Milton JT, Martin GB (2006). Endocrine and metabolic factors involved in the effect of nutrition on the production of colostrum in female sheep. *Reproduction Nutrition Development* 46:447-460.
- Barger IA (1993). Influence of sex and reproductive status on susceptibility of ruminants to nematode parasitism. *International Journal of Parasitology* 23:463-469.
- Barker DJ, Winter PD, Osmond C, Margetts B, Simmonds SJ (1989). Weight in infancy and death from ischaemic heart disease. *Lancet* 2:577-580.
- Barrow GI, Feltham RKA (1993). *Manual for the Identification of Medical Bacteria*, 3rd edn. Cambridge University Press, Cambridge.
- Bassett, NS, Oliver MH, Breier BH, Gluckman PD (1990). The effect of maternal starvation on plasma insulin-like growth factor I concentrations in the late gestation ovine fetus. *Pediatric Research* 27:401-404.
- Bate LA, Ireland W, Connell BJ, Grimmelt B (1991). Development of the small intestine of piglets in response to prenatal elevation of glycocorticoids. *Histology and Histopathology* 6:207-216.
- Bath GK (1983). Differentiation between hypocalcaemia and pregnancy ketosis in sheep. *Proceedings of the XXII World Veterinary Congress*, Perth, Australia, p. 147.
- Beasley AM, Kahn LP, Windon RG (2010a). The periparturient relaxation of immunity in Merino ewes infected with *Trichostrongylus colubriformis*: endocrine and body compositional responses. *Veterinary Parasitology* 168:51-59.
- Beasley AM, Kahn LP, Windon RG (2010b). The periparturient relaxation of immunity in Merino ewes infected with *Trichostrongylus colubriformis*: parasitological and immunological responses. *Veterinary Parasitology* 168:60-67.

- Beasley AM, Kahn LP, Windon RG (2012). The influence of reproductive physiology and nutrient supply on the periparturient relaxation of immunity to the gastrointestinal nematode *Trichostrongylus colubriformis* in Merino ewes. *Veterinary Parasitology* 188:306-324.
- Bell AW, Bauman DE (1997). Adaptations of glucose metabolism during pregnancy and lactation. *Journal of mammary Gland Biology and Neoplasia* 2:265-278.
- Bergman EN (1973). Glucose metabolism in ruminants as related to hypoglycemia and ketosis. *American Journal of Physiology* 215:865-873.
- Bickhardt K, Henze P, Ganter M (1998). Clinical findings and differential diagnosis in ketosis and hypocalcemia in sheep. *Deutsche Tierärztliche Wochenschrift* 105:413-419.
- Bickhardt K, König G (1985). Blood values of healthy ewes of the Merino and Blackhead breed during parturition. *Deutsche Tierärztliche Wochenschrift* 92:319-322.
- Bispham J, Gopalakrishnan GS, Dandrea J, Wilson V, Budge H, Keisler DH, Broughton Pipkin F, Stephenson T, Symonds ME (2003). Maternal endocrine adaptation throughout pregnancy to nutritional manipulation: consequences for maternal plasma leptin and cortisol and the programming of fetal adipose tissue development. *Endocrinology* 144:3575-3585.
- Bloomfield FH, Spiroski AM, Harding JE (2013). Fetal growth factors and fetal nutrition. *Seminars in fetal and neonatal Medicine* 18:118-123.
- Bobe G, Young JW, Beitz DC (2004). Pathology, etiology, prevention, and treatment of fatty liver in dairy cows. *Journal of Dairy Science* 87:3105-3124.
- Boden G, Shulman GI (2002). Free fatty acids in obesity and type 2 diabetes: defining their role in the development of insulin resistance and beta-cell dysfunction. *European Journal of clinical Investigation* 32:14-23.
- Brameld JM, Mostyn A, Dandrea J, Stephenson TJ, Dawson JM, Buttery PJ, Symonds ME (2000). Maternal nutrition alters the expression of insulin-like growth factors in fetal sheep liver and skeletal muscle. *Journal of Endocrinology* 167:429-437.
- Brandon MR, Watson DL, Lascelle AK (1971a). Concentration of immunoglobulins in blood serum and whey of dairy cows. *Australian Journal of Dairy Technology* 26:143-148.
- Brandon MR, Watson DL, Lascelle AK (1971b). Mechanism of transfer of immunoglobulins into mammary secretion of cows. *Australian Journal of experimental Biology and medical Science* 49:613-619.
- Braun U, Forster E (2012). B-mode and colour Doppler sonographic examination of the milk vein and musculophrenic vein in dry cows and cows with a milk yield of 10 and 20 kg. *Acta veterinaria scandinavica* 54:15.

- Braun JP, Trumel C, Bézille P (2010). Clinical biochemistry in sheep: a selected review. *Small Ruminant Research* 92:10-18.
- Brozos C, Mavrogianni VS, Fthenakis GC (2011). Treatment and control of peri-parturient metabolic diseases: pregnancy toxemia, hypocalcemia, hypomagnesemia. *Veterinary Clinics of North America Food Animal Practice* 27:105-113.
- Bubb KJ, Cock ML, Black MJ, Dodic M, Boon WM, Parkington HC, Harding R, Tare M (2007). Intrauterine growth restriction delays cardiomyocyte maturation and alters coronary artery function in the fetal sheep. *Journal of Physiology* 578:871-881.
- Burvenich C, Peeters G (1980). Effect of intramammary infusion of colchicine on mammary blood flow in lactating goats. *Journal of animal Physiology and animal Nutrition* 44:211-217.
- Buswell JF, Haddy JP, Bywater RJ (1986). Treatment of pregnancy toxemia in sheep using a concentrated oral rehydration solution. *The veterinary Record* 118:208-209.
- Cal-Pereyra L, Benech A, Gonzalez-Montana JR, Acosta-Dibarrat J, Da Silva S, Martin A (2015a). Changes in the metabolic profile of pregnant ewes to an acute feed restriction in late gestation. *New Zealand veterinary Journal* 63:141-146.
- Cal-Pereyra L, Gonzalez-Montana JR, Benech A, Acosta-Dibarrat J, Martin MJ, Perini S, Abreu, MC, Da Silva S, Rodriguez P (2015b). Evaluation of three therapeutic alternatives for the early treatment of ovine pregnancy toxemia. *Irish veterinary Journal* 68:25.
- Caroprese M, Giannenas I, Fthenakis GC (2015). Interactions between nutritional approaches and defences against microbial diseases in small ruminants. *Veterinary Microbiology* 181:8-14.
- Caroprese M, Napolitano F, Mattiello S, Fthenakis GC, Ribó O, Sevi A (2016). On-farm welfare monitoring of small ruminants. *Small Ruminant Research* 135:20-25.
- Castro N, Capote J, Bruckmaier RM, Argüello A. (2011). Management effects on colostrumogenesis in small ruminants: a review. *Journal of applied Animal Research* 39:85-93.
- Chen X, Bai X, Mai P, Cai J, Liu Z, Wang H, Xiao H, Dong W, Wang S, Sun Z (2012). Angiotensin-like protein 3 expression is down-regulated in experimentally pregnant toxemic goats. *Journal of integrative Agriculture* 11:1181-1188.
- Chilliard Y, Ferlay A, Faulconnier Y, Bonnet M, Rouel J, Bocquier F (2000). Adipose tissue metabolism and its role in adaptations to under-nutrition in ruminants. *Proceedings of the Nutrition Society* 59:127-134.
- Christensen K, Nielsen MO, Bauer R, Hilden K (1989). Evaluation of mammary blood flow measurements in lactating goats using the ultrasound Doppler principle. *Comparative Biochemistry and Physiology* 92:385-392.

- Cieslar SRL, Madsen TG, Purdie NG, Trout DR, Osborne VR, Cant JP (2014). Mammary blood flow and metabolic activity are linked by a feedback mechanism involving nitric oxide synthesis. *Journal of Dairy Science* 97:2090-2100.
- Conklin D, Gilbertson D, Taft DW, Maurer MF, Whitmore TE, Smith DL, Walker KM, Chen LH, Wattler S, Nehls M, Lewis KB (1999). Identification of a mammalian angiopoietin-related protein expressed specifically in liver. *Genomics* 62:477-482.
- Coop RL, Kyriazakis I (1999). Nutrition-parasite Interaction. *Veterinary Parasitology* 84:187–2004.
- Coop RL, Mellor DJ, Jackson E, Jackson F, Flint DJ, Vernon RG (1990). Teladorsagia circumcincta egg output at the outset of natural and induced lactation in ewes. *Veterinary Parasitology* 35:295-305.
- Coop RL, Sykes AR, Angus KW (1977). The effect of a daily intake of Ostertagia circumcincta larvae on body weight, food intake and concentration of serum constituents in sheep. *Research in veterinary Science* 23:76-83.
- Curtis CR, Erb HN, Sniffen CJ, Smith RD, Kronfeld DS (1985). Path analysis of dry-period nutrition, postpartum metabolic and reproductive disorders, and mastitis in Holstein cows. *Journal of Dairy Science* 68:2347-2360.
- Cvek K (1997). *Mammary Gland Function with Special Reference to Vascularisation and Atrial Natriuretic Peptide*. MSc Dissertation, Swedish University of Agricultural Sciences, Uppsala, Sweden.
- Dakkak A (1990). Gastrointestinal strongylosis and malabsorption of nutrients. *Annales de Parasitologie humaine et comparee* 65:73-76.
- Daley M, Williams T, Coyle P, Furda G, Dougherty R, Hayes P (1993). Prevention and treatment of Staphylococcus aureus infections with recombinant cytokines. *Cytokine* 5:276-284.
- Davis SR, Collier RJ (1985). Mammary blood flow and regulation of substrate supply for milk synthesis. *Journal of Dairy Science* 68:1041-1058.
- Delamaire E, Guinard-Flament J (2006). Increasing milking intervals decreases the mammary blood flow and mammary uptake of nutrients in dairy cows. *Journal of Dairy Science* 89:3439-3446.
- Dhondt G, Burvenich C, Peeters G (1977a). Mammary blood flow during experimental Escherichia coli endotoxin induced mastitis in goats and cows. *Journal of Dairy Research* 44:433-440.
- Dhondt G, Houvenaghel A, Peeters G, Jöchle W (1977b). Effect of prostaglandins F2 $\alpha$  and E2 on milk ejection, blood pressure and blood flow through the mammary artery in the cow. *Prostaglandins* 13:1185-1199.

- Diltoer MW, Troubleyn J, Lauwers R, De Wijngaard S, Vercammen MJ, Hubloue I, Huyghens LP (2004). Ketosis and cardiac failure: common signs of a single condition. *European Journal of Emergency Medicine* 11:172-175.
- Divon MY (1993). Fetal respiratory sinus arrhythmia. In: Chervenak F, Isaacson G, Campbell S (Eds) *Textbook of Ultrasound in Obstetrics and Gynecology*. Little Brown, Boston, pp. 533-538.
- Dominguez E, del Alamo MMR, Novellas R, Espada Y, Santos L, Garcia E, Andaluz A, Fresno L (2013). Doppler evaluation of the effects of propofol, etomidate and alphaxalone on fetoplacental circulation hemodynamics in the pregnant ewe. *Placenta* 34:738-744.
- Duehlmeier R, Fluegge I, Schwert B, Parvizi N, Ganter M (2011). Metabolic adaptations to pregnancy and lactation in German Blackheaded Mutton and Finn sheep ewes with different susceptibilities to pregnancy toxemia. *Small Ruminant Research* 96:178-184.
- Duehlmeier R, Noldt S, Ganter M (2013). Pancreatic insulin release and peripheral insulin sensitivity in German black headed mutton and Finish Landrace ewes: evaluation of the role of insulin resistance in the susceptibility to ovine pregnancy toxemia. *Domestic Animal Endocrinology* 44:213-221.
- Dumortier O, Blondeau B, Duvillié B, Reusens B, Bréant B, Remacle C (2007). Different mechanisms operating during different critical timewindows reduce ratfetal beta cell mass due to a maternal low-protein or low-energy diet. *Diabetologia* 50:2495-2503.
- Dwyer CM (2003). Behavioural development in the neonatal lamb: effect of maternal and birth-related factors. *Theriogenology* 59:1027-1050.
- Edwards LJ, McMillen IC (2002). Impact of maternal undernutrition during the periconceptional period, fetal number, and fetal sex on the development of the hypothalamo-pituitary adrenal axis in sheep during late gestation. *Biology of Reproduction* 66:1562-1569.
- Eliot RJ, Klein AH, Glatz TH, Nathanielsz PW, Fisher DA (1981). Plasma norepinephrine, epinephrine, and dopamine concentrations in maternal and fetal sheep during spontaneous parturition and in premature sheep during cortisol-induced parturition. *Endocrinology* 108:1678-1682.
- European Food Safety Authority (2014). Scientific opinion on the welfare risks related to the farming of sheep for wool, meat and milk production. *EFSA Journal* 12:3933, 128 p.
- Euzeby JP (1997). List of bacterial names with standing in nomenclature: a folder available on the Internet. *International Journal of systematic Bacteriology* 47:590-592.

- Everett-Hincks JM, Blair HT, Stafford KJ, Lopez-Villalobos N, Kenyon PR, Morris ST (2005). The effect of pasture allowance fed to twin- and triplet-bearing ewes in late pregnancy on ewe and lamb behaviour and performance to weaning. *Livestock Production Science* 97:253-266.
- Everts H, Kuiper H 1983. Energy intake and pregnancy toxemia in prolific ewes. *Proceedings of the 5th International Conference on Production Diseases in Farm Animals*, Uppsala, Sweden, pp. 133–136.
- Faulconnier Y, Bonnet M, Bocquier F, Leroux C, Chilliard Y (2001). Effects of photoperiod and feeding level on adipose tissue and muscle lipoprotein lipase activity and mRNA level in dry non-pregnant sheep. *British Journal of Nutrition* 85:299-306.
- Ferris TF, Herdson PB, Dunnill MS, Radcliffe-Lee M (1969). Toxemia of pregnancy in sheep: a clinical, physiological and pathological study. *Journal of clinical Investigation* 48:1643-1655.
- Fleming MW (1993). Acute or chronic administration of prolactin alters ovine infections of *Haemonchus contortus*. *Veterinary Parasitology* 50:109-115.
- Fleming MW (1995). Effects of exogenous reproductive hormones on *Haemonchus contortus* populations in lambs. *Journal of the helminthological Society of Washington* 64:269-274.
- Foster LA, Ames NK, Emery RS (1991). Food intake and serum insulin responses to intraventricular infusions of insulin and IGF-I. *Physiology and Behavior* 50:745-749.
- Fowden AL (2003). The insulin-like growth factors and feto-placental growth. *Placenta* 24:803-812.
- Fowden AL, Li J, Forhead AJ (1998). Glucocorticoids and the preparation for life after birth: are there long-term consequences of the life insurance? *Proceedings of the Nutrition Society* 57:113-122.
- Fragkou IA, Boscós CM, Fthenakis GC (2014). Diagnosis of clinical or subclinical mastitis in ewes. *Small Ruminant Research* 188:86-92.
- Fragkou IA, Dagleish MP, Papaioannou N, Cripps PJ, Boscós CM, Ververidis HN, Orfanou DC, Solomakos N, Finlayson J, Govaris A, Kyriazakis I, Fthenakis GC (2010). The induction of lymphoid follicle-like structures in the ovine teat duct following experimental infection with *Mannheimia haemolytica*. *Veterinary Journal* 184:194-200.
- Fragkou IA, Gougoulis DA, Billinis C, Mavrogianni VS, Bushnell MJ, Cripps PJ, Tzora A, Fthenakis GC (2011). Transmission of *Mannheimia haemolytica* from the tonsils of lambs to the teat of ewes during sucking. *Veterinary Microbiology* 148:66-74.
- Fragkou IA, Mavrogianni VS, Papaioannou N, Boscós C, Cripps PJ, Skoufos J, Fthenakis GC (2007). Presence of sub-epithelial lymphoid nodules in the teat of ewe-lambs and adult ewes. *Small Ruminant Research* 70:286-291.

- Franklin ST, Young JW, Nonnecke BJ (1991). Effects of ketones, acetate, butyrate, and glucose on bovine lymphocyte proliferation. *Journal of Dairy Science* 74:2507-2514.
- Fthenakis GC (1994). Prevalence and aetiology of subclinical mastitis in ewes of southern Greece. *Small Ruminant Research* 13:293-300.
- Fthenakis GC (1995). California Mastitis Test and Whiteside Test in diagnosis of subclinical mastitis of dairy ewes. *Small Ruminant Research* 16:271-276.
- Fthenakis GC, Arsenos G, Brozos C, Fragkou IA, Gianidis ND, Giannenas I, Mavrogianni VS, Papadopoulos E, Valasi I (2012). Health management of ewes during pregnancy. *Animal Reproduction Science* 130:198-212.
- Fthenakis GC, Jones (1990a). Incidence and aetiology of clinical ovine mastitis in flocks in Central Macedonia, Greece. *Bulletin of the Hellenic veterinary medical Society* 41:133-141.
- Fthenakis GC, Jones JET (1990b). The effect of inoculation of coagulase-negative staphylococci into the ovine mammary gland. *Journal of comparative Pathology* 102:211-219.
- Fthenakis GC, Leontides LS, Amiridis GS, Saratsis P (2000). Incidencerisk and clinical features of retention of foetal membranes in ewes in 28 flock in southern Greece. *Preventive veterinary Medicine* 44:113-120.
- Fthenakis GC, Leontides L, Skoufos J, Taitzoglou IA, Tzora A (2004). Case report: high prevalence rate of ovine mastitis, caused by coagulase-negative staphylococci and predisposed by increased gossypol consumption. *Small Ruminant Research* 52:185-189.
- Fthenakis GC, Mavrogianni VS, Gallides E, Papadopoulos E (2015). Interactions between parasitic infections and reproductive efficiency in sheep. *Veterinary Parasitology* 208:56-66.
- Fthenakis GC, Papadopoulos E, Himonas C (2005). Effects of three anthelmintic regimes on milk yield of ewes and growth of lambs. *Journal of veterinary Medicine A* 52:78-82.
- Furll M, Deniz A, Westphal B, Illing C, Constable PD (2010). Effect of multiple intravenous injections of butaphosphan and cyanocobal-amin on the metabolism of periparturient dairy cows. *Journal of Dairy Science* 93:4155-4164.
- Galan HL, Anthony RV, Rigano S, Parker TA, de Vrijer B, Ferrazzi E, Wilkening RB, Regnault TRH (2005). Fetal hypertension and abnormal Doppler velocimetry in an ovine model of intrauterine growth restriction. *American Journal of Obstetrics and Gynecology* 192:272-279.
- Galatos AD (2011). Anesthesia and analgesia in sheep and goats. *Veterinary Clinics of North America Food Animal Practice* 27:47-59.

- Gao F, Liu Y, Li L, Li M, Zhang C, Ao C, Hou X (2014). Effects of maternal undernutrition during late pregnancy on the development and function of ovine fetal liver. *Animal Reproduction Science* 147:99-105.
- Gerloff BJ, Herdt TH, Emery RS (1993). The relationship of hepatic lipidosis to health and performance in dairy cattle. *Journal of the American Veterinary Medical Association* 188:845-850.
- Giadinis ND, Panousis N, Petridou EJ, Siarkou VI, Lafi SQ, Pourliotis K, Hatzopoulou E, Fthenakis GC (2011). Selenium, vitamin E and vitamin A blood concentrations in dairy sheep flocks with increased or low clinical mastitis incidence. *Small Ruminant Research* 95:193-196.
- Gibbs HC (1986). Hypobiosis and the periparturient rise in sheep. *Veterinary Clinics of North America Food Animal Practice* 2:345-353.
- Ginther OJ (2007). *Ultrasonic Imaging and Animal Reproduction: Color-Doppler Ultrasonography*. Equiservices Publishing, Wisconsin.
- Gluckman PD, Breier BH, Davis SR (1987). Physiology of the somatotrophic axis with particular reference to the ruminant. *Journal of Dairy Science* 70:442-466.
- Gluckman PD, Hanson MA (2004). Developmental origins of disease paradigm: a mechanistic and evolutionary perspective. *Pediatric Research* 56:311-317.
- Gluckman P, Hanson M (2006). *Mismatch: why our World no longer fits our Bodies*. Oxford University Press, Oxford.
- Gougoulis DA, Kyriazakis I, Mavrogianni VS, Fragkou IA, Skoufos J, Tzora A, Taitzoglou IA, Kokoli AN, Fthenakis GC (2007). Patterns of maternal-offspring behaviour of dairy sheep and potential association with mammary health. *Canadian Journal of Animal Science* 87:469-478.
- Granger DN, Senshenkova E (2010). *Inflammation and the Microcirculation*. Morgan & Claypool Life Sciences, San Rafael.
- Greenwood PL, Hunt AS, Hermanson JW, Bell AW (2000). Effects of birth weight and postnatal nutrition on neonatal sheep: II. Skeletal muscle growth and development. *Journal of Animal Science* 78: 50-61.
- Greiss FC (1972). Differential reactivity of the myoendometrial and placental vasculatures: adrenergic responses. *American Journal of Obstetrics and Gynaecology* 112:20-30.
- Grinberg N, Elazar S, Rosenshine I, Shpigel NY (2008). Beta-hydroxybutyrate abrogates formation of bovine neutrophil extracellular traps and bactericidal activity against mammary pathogenic *Escherichia coli*. *Infection and Immunity* 76:802-807.
- Guard C (1995). Metabolic diseases: a herd approach. In: Rebhun W (ed.) *Diseases of Dairy Cattle*, Williams & Wilkins, Baltimore, pp. 497-509.



- Gurdogan F, Balikci E, Yildiz A (2014). Some acute phase proteins, oxidative stress biomarkers and antioxidant enzyme activities in ewes with pregnancy toxemia. *Iranian Journal of veterinary Research* 15:297-299.
- Gürtler H, Schweigert FJ (2005). Physiologie der Laktation. In: Engelhardt WV, Breves G (eds) *Physiologie der Haustiere*, 2nd edn. Enke im Hippokrates Verlag, Stuttgart, pp. 572-593.
- Han HC, Austin KJ, Nathanielsz PW, Ford SP, Nijland MJ, Hansen TR (2004). Maternal nutrient restriction alters gene expression in the ovine fetal heart. *Journal of Physiology* 558:111-121.
- Harmeyer J, Schlumbohm C (2006). Pregnancy impairs ketone body disposal in late gestating ewes: implications for onset of pregnancy toxemia. *Research in veterinary Science* 81:254-264.
- Hart IC (1983). Endocrine control of nutrient partition in lactating ruminants. *Proceedings of the Nutrition Society* 42:181-194.
- Hay WW Jr, Sparks JW, Wilkening RB, Battaglia FC, Meschia G (1983). Partition of maternal glucose production between conceptus and maternal tissues in sheep. *American Journal of Physiology* 245:E347-E350.
- Hea ZX, Wua DQ (2013). Protein or energy restriction during late gestation alters fetal growth and visceral organ mass: an evidence of intrauterine programming in goats. *Animal Reproduction Science* 137:177-182.
- Henderson GD, Read LC, Sniswell AM (1982) Studies of liver lipids in normal, alloxan-diabetic and pregnancy-toxaemic sheep. *Biochimica et biophysica Acta* 710:236-241.
- Henry BA, Goding JW, Tilbrook AJ, Dunshea FR, Clarke IJ (2001). Intracerebroventricular infusion of leptin elevates the secretion of luteinising hormone without affecting food intake in long-term food-restricted sheep, but increases growth hormone irrespective of bodyweight. *Journal of Endocrinology* 168:67-77.
- Henze P, Bickhardt K, Fuhrmann H (1994). The influences of insulin, cortisol, growth hormone and total oestrogen on the pathogenesis of ketosis in sheep. *Deutsche Tierärztliche Wochenschrift* 101:61-65.
- Henze P, Bickhardt K, Fuhrmann H, Sallmann HP (1998). Spontaneous pregnancy toxemia (ketosis) in sheep and the role of insulin. *Journal of veterinary Medicine A* 45:255-266.
- Hindson JC, Winter AC (1990). *Outline of Clinical Diagnosis in Sheep*. Wright, London.
- Hindson JC, Winter AC (2007). Genital abnormalities, obstetrical problems and birth injuries. In: Aitken ID (ed.) *Diseases of Sheep*, 4th edn. Blackwell Publishing, Oxford, pp. 75-80.

- Hoeben D, Heyneman R, Burvenich C (1997). Elevated levels of beta-hydroxybutyric acid in periparturient cows and in vitro effect on respiratory burst activity of bovine neutrophils. *Veterinary Immunology and Immunopathology* 58:165-170.
- Houdijk JGM, Jessop NS, Kyriazakis I (2001). Nutrient partitioning between reproductive and immune functions in animals. *Proceedings of the Nutrition Society* 60:515-525.
- Ioannidi KS, Vasileiou NGC, Barbagianni MS, Orfanou DC, Houzouris TM, Dovolou E, Amiridis GS, Fthenakis GC, Mavrogianni VS (2015). Clinical, bacteriological and cytological findings during uterine involution in ewes that had developed pregnancy toxemia. *Proceedings of the 32nd World Veterinary Congress, Istanbul, Turkey*, p. 26.
- Ismail ZAB, Al-Majali AM, Amireh F, Al-Rawashdeh OF (2008). Metabolic profiles in goat does in late pregnancy with and without subclinical pregnancy toxemia. *Veterinary Clinical Pathology* 37:434-437.
- Jain SK, Kannan K, Lim G (1998). Ketosis (acetoacetate) can generate oxygen radicals and cause increased lipid peroxidation and growth inhibition in human endothelial cells. *Free Radical Biology and Medicine* 25:1083-1088.
- Jaquiere AL, Oliver MH, Bloomfield FH, Connor KL, Challis JRG, Harding JE (2006). Fetal exposure to excess glucocorticoid is unlikely to explain the effects of periconceptional undernutrition in sheep. *Journal of Physiology* 572:109-118.
- Jelinek PD, Steele P, Masters HG, Allen JG, Copland MD, Petterson DS (1985). Erythrocyte selenium-75 uptake as a measure of selenium status in weaner sheep, and its relationship to erythrocyte glutathione peroxidase activity. *Australian Veterinary Journal* 62:327-331.
- Jones JET (1985). An investigation of mastitis in sheep: preliminary phase. *Proceedings of Meetings of the Sheep Veterinary Society* 10:48-51.
- Jones JI, Clemmons DR (1995). Insulin-like growth factors and their binding proteins: biological actions. *Endocrine Reviews* 16:3-34.
- Jones JET, Lanyon ME (1987). An investigation of ovine mastitis in England and Wales: preliminary results. *Proceedings of Annual Meeting of Association of Veterinary Teachers and Research Workers, Scarborough, Great Britain*, pp. (1) 21-22.
- Kahn LP, Knox MR, Gray GD, Lea JM, Walkden-Brown SW (2003). Enhancing immunity to nematode parasites in single-bearing Merino ewes through nutrition and genetic selection. *Veterinary Parasitology* 112:211-225.

- Kamada H, Nomaka I, Ueda Y, Murai M (2007). Selenium addition to colostrums increases immunoglobulin G absorption by newborn calves. *Journal of Dairy Science* 90:5665-5670.
- Kandefer-Szerszen M, Filar J, Szuster-Ciesielska A, Rzeski W (1992). Suppression of interferon response of bovine leukocytes during clinical and subclinical ketosis in lactating cows. *Deutsche Tierärztliche Wochenschrift* 99:440-443.
- Kaneene JB, Miller R, Herdt TH, Gardiner JC (1997). The association of serum non-esterified fatty acids and cholesterol, management and feeding practices with peri-partum disease in dairy cows. *Preventive veterinary Medicine* 31:59-72.
- Kankofer M, Lipko-Przybylska J (2008). Physiological antioxidative/oxidative status in bovine colostrum and mature milk. *Acta veterinaria Beograd* 58:231-239.
- Kanz P, Drillich M, Klein-Joebstl D, Mair B, Borchardt S, Meyer L, Schwendenwein I, Iwersen M (2015). Suitability of capillary blood obtained by a minimally invasive lancet technique to detect subclinical ketosis in dairy cows by using three different electronic hand-held devices. *Journal of Dairy Science* 98:6108-6118.
- Kloss S, Wehrend A, Failing K, Bostedt H (2002). Investigations about kind and frequency of mechanical dystocia in ewes with special regard to the vaginal prolapse ante partum. *Berliner und Munchener Tierärztliche Wochenschrift* 115:247-251.
- Koutsoumpas AT, Giadinis ND, Petridou EJ, Konstantinou E, Brozos C, Lafi SQ, Fthenakis GC, Karatzias H (2013). Consequences of reduced vitamin A administration on mammary health of dairy ewes. *Small Ruminant Research* 110:120-123.
- Kramer JW (2000). Normal hematology of cattle, sheep and goats. In: Feldman BF, Zinkl JG, Jain NC (eds) *Schalm's Veterinary Hematology*, 5th edn. Lippincott Williams & Wilkins, Baltimore, pp. 1075-1084.
- Kronfeld DS, Raggi F, Ramberg CF Jr (1968). Mammary blood flow and ketone body metabolism in normal, fasted, and ketotic cows. *American Journal of Physiology* 215:218-227.
- Kronfeld DS, Simesen MG (1961). Glucose biokinetics in ovine pregnancy toxemia. *Cornell Veterinarian* 51:478-488.
- Kulcsar M, Danko G, Delavaud C, Faigl V, Kerestes M, Gaspard A, Chilliard Y, Cseh S, Huszenicza G (2006). Endocrine status and subsequent reproduction in subclinical form of ovine ketosis. *Reproduction in domestic Animals* 41:372.
- Kvietys PR, Granger DN (2014). The splanchnic circulation. In: Reinus JF, Simon D (eds) *Gastrointestinal Anatomy and Physiology: the Essentials*. Wiley-Blackwell, Chichester.

- Lacetera N, Bernabucci U, Ronchi B, Nardone A (2001). Effects of sub-clinical pregnancy toxemia on immune responses in sheep. *American Journal of veterinary Research* 62:1020-1024.
- Lacetera N, Scalia N, Bernabucci U, Ronchi B, Pirazzi D, Nardone A (2005). Lymphocyte functions in overconditioned cows around parturition. *Journal of Dairy Science* 88:2010-2016.
- Langley-Evans S.C. (2004). Fetal programming of adult disease: an overview. In: Langley-Evans SC (ed.) *Fetal Nutrition and Adult Disease: Programming of Chronic Disease through Fetal Exposure to Undernutrition*. CAB International, Wallingford, pp. 1-20.
- Langley-Evans SC, Wakelin D, Buttery PJ (2003). Undernutrition during fetal life programmes immune function in the rat. *Pediatric Research* 53: P104.
- Leal-Cerro A, Soto A, Martinez MA, Diequez C, Casanueva FF (2001). Influence of cortisol status on leptin secretion. *Pituitary* 4:111-116.
- Lean IJ (2002). Diseases of dairy animals, noninfectious/Pregnancy toxemia. In: Rojinski H (ed.) *Encyclopedia of Dairy Sciences*. Elsevier, Amsterdam, pp.830-836.
- LeBlanc S (2010). Monitoring metabolic health of dairy cattle in the transition period. *Journal of Reproduction and Development* 56:29-35.
- Lemley CO, Meyer AM, Neville TL, Hallford DM, Camacho LE, Maddock-Carlin KR, Wilmoth TA, Wilson ME, Perry GA, Redmer DA, Reynolds LP, Caton JS, Vonnahme KA (2014). Dietary selenium and nutritional plane alter specific aspects of maternal endocrine status during pregnancy and lactation. *Domestic Animal Endocrinology* 46:1-11.
- Lima MS., Pascoal RA, Stilwell GT (2012). Glycaemia as a sign of the viability of the foetuses in the last days of gestation in dairy goats with pregnancy toxemia. *Irish veterinary Journal* 65:1.
- Linzell JL (1960). Mammary gland blood flow and oxygen, glucose and volatile fatty acid uptake in the conscious goat. *Journal of Physiology* 153:492-509.
- Linzell JL (1967) The effect of very frequent milking and of oxytocin on the yield and composition in fed and fasted goats. *Journal of Physiology* 190:333-346.
- Linzell JL (1974). Mammary blood flow and methods of identifying and measuring precursors of milk. In: Larson BL, Smith VR (eds.) *Lactation. A Comprehensive Treatise*. Academic Press, New York, vol. 1, pp. 143-225.
- Lipsett J, Tamblyn M, Madigan K, Roberts P, Cool JC, Runciman SI, McMillen IC, Robinson J, Owens JA (2006). Restricted fetal growth and lung development: a morphometric analysis of pulmonary structure. *Pediatric Pulmonology* 41:1138-1145.
- Locatelli P, Olea F, De Lorenzi A, Salmo F, Janavel GLV, Hnatiuk AP, Guevara E, Crottogini A (2011). Reference values for echocardiographic parameters and indexes of left ventricular

function in healthy, young adult sheep used in translational research: comparison with standardized values in humans. *International Journal of clinical and experimental Medicine* 4:258-264.

Lopes Camara ACL, Afonso JAB, Dantas AC, Guimaraes JA, Costa NA, Souza MI, Mendonca CL (2009). Analysis of factors related to 60 dystocia cases in ewes in the Agreste and semiarid region of Pernambuco, Northeastern Brazil. *Ciencia Rural* 39:2458-2463.

Lough DS, Beede DL, Wilcox CJ (1990). Effects of feed intake and thermal stress on mammary blood flow and other physiological measurements in lactating dairy cows. *Journal of Dairy Science* 73:325-332.

Lowry R (2012). *Concepts and Applications of Inferential Statistics*. Vassar College, Poughkeepsie.

Lowry R (2015). VassarStats Website. <http://vassarstats.net>.

Luther J, Aitken R, Milne J, Matsuzaki M, Reynolds L, Redmer D, Wallace J (2007). Maternal and fetal growth, body composition, endocrinology, and metabolic status in undernourished adolescent sheep. *Biology of Reproduction* 77:343-350.

Lynch JJ, Leng RA, Hinch GN, Nolan J, Bindon BM, Piper LR (1990). Effects of cottonseed supplementation on birth weights and survival of lambs from a range of litter sizes. *Proceedings of the Australian Society of Animal Production* 18:516.

Madjar H, Mendelson EB (2008). *The Practice of Breast Ultrasound*, 2nd edn. Thieme, Stuttgart.

Makikallio K, Erkinaro T, Niemi N, Kavasmaa T, Acharya G, Pakkila M, Rasanen J (2006). Fetal oxygenation and Doppler ultrasonography of cardiovascular hemodynamics in a chronic near-term sheep model. *American Journal of Obstetrics and Gynecology* 194:542-550.

Margiloff L, Chaplia L, Chow A, Singhal PC, Mattana J (1998). Metal-catalyzed oxidation of immunoglobulin G impairs Fc receptor-mediated binding to macrophages. *Free Radical Biology and Medicine* 25:780-785.

Marteniuk JV, Herdt TH (1988). Pregnancy toxemia and ketosis of ewes and does. *Veterinary Clinics of North America Food Animal Practice* 4:307-315.

Martin WB, Aitken ID (2000). Haematological reference values for healthy sheep. In: Martin WB, Aitken ID (eds) *Diseases of Sheep*, 3rd edn. Blackwell, Oxford, p. 499.

Maulik D (2005). Spectral Doppler sonography: waveform analysis and hemodynamic information. In: Maulik D (ed.) *Doppler Ultrasound in Obstetrics and Gynecology*, 2nd edn. Springer, Berlin, pp. 35-56.

- Mavrogianni VS, Brozos C (2008). Reflections on the causes and the diagnosis of peri-parturient losses of ewes. *Small Ruminant Research* 76:77-82.
- Mavrogianni VS, Cripps PJ, Fthenakis GC (2006a). Description and validation of a novel technique to study the bacterial flora of the teat duct of ewes. *Small Ruminant Research* 37:89-105.
- Mavrogianni VS, Cripps PJ, Papaioannou N, Taitzoglou I, Fthenakis GC (2006b). Teat disorders predispose ewes to clinical mastitis after challenge with *Mannheimia haemolytica*. *Veterinary Research* 66:258-264.
- Mavrogianni VS, Fthenakis GC, Brooks H, Papaioannou N, Cripps PJ, Taitzoglou I, Brellou G, Saratsis P (2005). The effects of inoculation of *Mannheimia haemolytica* into the teat of lactating ewes. *Veterinary Research* 36:13-25.
- McCarthy MM, Mann S, Nydam DV, Overton TR, Mcart JAA (2015). Short communication: Concentrations of nonesterified fatty acids and beta-hydroxybutyrate in dairy cows are not well correlated during the transition period. *Journal of Dairy Science* 98:6284-6290.
- McCausland IP, O'Hara PJ, Hederson PB, Leet NG (1974). Spontaneous toxæmia of pregnancy in sheep: a study of renal function and glomerular fine structure. *Journal of comparative Pathology* 84:375-380.
- McGuirk SM, Collins M (2004). Managing the production, storage and delivery of colostrum. *Veterinary Clinics of North America Food Animal Practice* 20:593-607.
- McNeill DM, Murphy PM, Lindsay DR (1998). Blood lactose versus milk lactose as a monitor of lactogenesis and colostrum production in Merino ewes. *Australian veterinary Journal* 49:581-587.
- Mee JF (2014). Risk factors for reproductive disorders (metritis, endometritis, cystic ovaries, anovulation, low conception rate, embryonic mortality) in dairy cows. *Cattle Practice* 22:42-53.
- Mellor DJ, Murray L (1985). Effects of maternal nutrition on udder development during late pregnancy and on colostrum production in Scottish Blackface with twin lambs. *Research in veterinary Science* 39:230-234.
- Merlot E, Couret D, Otten W (2008). Prenatal stress, fetal imprinting and immunity. *Brain, Behavior and Immunity* 22:42-51.
- Meurer DG, Wolf S (1999) *Allgemeine Pathologie: Kompendium für die Veterinärmedizin*. Schattauer Verlagsgesellschaft, Stuttgart.
- Meyer AM, Reed JJ, Neville TL, Thorson JF, Maddock-Carlin KR, Taylor JB, Reynolds LP, Redmer DA, Luther JS, Hammer CJ, Vonnahme KA, Caton JS (2011). Nutritional plane and

- selenium supply during gestation affect yield and nutrient composition of colostrums and milk in primiparous ewes. *Journal of animal Science* 89:1627-1639.
- Miles AA, Misra JS (1938). The estimation of the bactericidal power of the blood. *Journal of Hygiene* 38:732-749.
- Ministry of Agriculture, Fisheries and Food (1986). *Manual of Veterinary Parasitological Laboratory Techniques*. Her Majesty's Stationery Office, London.
- Miranda R, Saravia NG, Ackerman R, Murphy N, Berman S, McMurray DN (1983). Effect of the maternal nutritional status on immunological substances in human colostrum and milk. *American Journal of clinical Nutrition* 37:632-640.
- Monin P (1994). Modifications of ventilator reflexes. An efficient therapy for apneas of prematurity. *Biology of the Neonate* 65:247-251.
- Morrison CD, Daniel JA, Holmberg BJ, Djiane J, Raver N, Gertler A, Keisler DH (2001). Central infusion of leptin into well-fed and undernourished ewe lambs: effects on feed intake and serum concentrations of growth hormone and luteinizing hormone. *Journal of Endocrinology* 168:317-324.
- Morrow RJ, Bull SB, Adamson SL (1993). Experimentally induced changes in heart rate alter umbilicoplacental hemodynamics in fetal sheep. *Ultrasound in Medicine and Biology* 19:309-318.
- Mukasa-Mugerwa E, Said AN, Lahlou-Kassi A, Sherington J, Mutiga ER (1994). Birth weight as a risk factor for perinatal lamb mortality, and the effects of stage of pregnant ewe supplementation and gestation weight gain in Ethiopian Menz sheep. *Preventive veterinary Medicine* 19:45-56.
- National Institutes of Health 2013 *Image J: Image Processing and Analysis in Java*.  
<http://rsbweb.nih.gov/ij/>
- Natrajan PG, McGarrigle HH, Lawrence DM, Lachelin GC (1982). Plasma noradrenaline and adrenaline levels in normal pregnancy and in pregnancy-induced hypertension. *British Journal of Obstetrics and Gynaecology* 89:1041-1045.
- Neary JM, Garry FB (2014). Impaired alveolar-arterial oxygen transfer is associated with reduced milk yield in primiparous post-partum dairy heifers at moderate altitude. *Journal of Dairy Research* 81:434-439.
- Nicholas R, Ayling R, McAuliffe L (2008). *Mycoplasma Diseases of Ruminants*. CAB International, Wallingford.

- Noakes DE (1996). Parturition and the care of parturient animals. In: Arthur GH, Noakes DE, Pearson H, Parkinson TJ (eds) *Veterinary Reproduction & Obstetrics*, 7th edn. Saunders, London, pp. 141-170.
- Oike Y, Akao M, Kubota Y, Suda T (2005). Angiopoietin-like proteins: potential new targets for metabolic syndrome therapy. *Trends in molecular Medicine* 11:473-479.
- Ojala T, Pietikaeinen M, Maenpaa T (2002). Multiresolution gray-scale and rotation invariant texture classification with local binary patterns. *IEEE Transactions on Pattern Analysis and Machine Intelligence* 24:971-987.
- Olofsson P, Olofsson H, Molin J, Marsal K(2004). Low umbilical artery vascular flow resistance and fetal outcome. *Acta obstetrica et gynecologica scandinavica* 83:440-442.
- Olsson K, Hydbring E, Cvek K (1998). Cardiovascular and fluid regulatory mechanisms during pregnancy and lactation in small ruminants. *Trends in comparative Biochemistry and Physiology* 5:283-296.
- Otranto D, Traversa D (2002). A review of dicrocoeliosis of ruminants including recent advances in the diagnosis and treatment. *Veterinary Parasitology* 107:317-335.
- Palmer AC (2011). Nutritionally mediated programming of the developing immune system. *Advances in Nutrition* 2:377-395.
- Panousis N, Brozos C, Fthenakis GC, Karatzias C (2001). Pregnancy toxemia of ewes. A literature review. *Journal of the hellenic veterinary medical Society* 52:89-96.
- Panousis N, Brozos C, Karagiannis I, Giadinis ND, Lafi S, Kritsepi-Konstantinou M (2012). Evaluation of Precision Xceed® meter for on-site monitoring of blood beta-hydroxybutyric acid and glucose concentrations in dairy sheep. *Research in veterinary Science* 93:435-439.
- Papadopoulos E, Mavrogianni VS, Mitsoura A, Ptochos S, Spanos SA, Fthenakis GC (2013). Potential association between trematode infections and development of pregnancy toxemia in sheep. *Helminthologia* 50:161-166.
- Pereira RA, Fensterseifer S, Barcelos VB, Martins CF, Schneider A, Schmitt E, Pfeifer LFM, Pino FABD, Correa MN (2013). Metabolic parameters and dry matter intake of ewes treated butaphosphan andcyanocobalamin in the early postpartum period. *Small Ruminant Research* 114:140-145.
- Perkins KH, VandeHaar MJ, Burton JL, Liesman JS, Erskine RJ, Elsasser TH (2002). Clinical responses to intramammary endotoxin infusion in dairy cows subjected to feed restriction. *Journal of Dairy Science* 85:1724-1731.



- Pethick DW, Lindsay DB (1982). Metabolism of ketone bodies in pregnant sheep. *British Journal of Nutrition* 48:549-563.
- Petridis IG, Gouletsou PG, Barbagianni MS, Amiridis GS, Brozos C, Valasi I, Fthenakis GC (2014). Ultrasonographic findings in the ovine udder during involution. *Journal of Dairy Research* 81:288-296.
- Petterson JA, Dunshea FR, Ehrhardt RA, Bell AW (1993). Pregnancy and undernutrition alter glucose metabolic responses to insulin in sheep. *Journal of Nutrition* 123:1286-1295.
- Piccione G, Arcigli A, Fazio F, Giudice E, Caola G (2004). Pulsed wave-doppler ultrasonographic evaluation of mammary blood flow speed in cows during different productive periods. *Acta scientiae veterinariae* 32:171-175.
- Pichler M, Damberger A, Schwendenwein I, Gasteiner J, Drillich M, Iwersen M (2014). Thresholds of whole-blood beta-hydroxybutyrate and glucose concentrations measured with an electronic hand-held device to identify ovine hyperketonemia. *Journal of Dairy Science* 97:1388-1399.
- Poser H, Semplicini L, De Benedictis GM, Gerardi G, Contiero B, Maschietto N, Valerio E, Milanese O, Semplicini A, Bernardini D (2013). Two-dimensional, M-mode and Doppler-derived echocardiographic parameters in sedated healthy growing female sheep. *Laboratory Animals* 47:194-202.
- Potapow A. (2010). *Investigation of Mammary Blood Flow Changes by Transrectal Colour Doppler Sonography in an Escherichia coli Mastitis Model*. PhD thesis, Ludwig-Maximilians-University Munich, Munich.
- Prosser CG, Davis SR, Farr VC, Lacasse P (1996). Regulation of blood flow in the mammary microvasculature. *Journal of Dairy Science* 79:1184-1197.
- Przybylska J, Albera E, Kankofer M (2007). Antioxidants in bovine colostrum. *Reproduction in domestic Animals* 42:402-409.
- Putu IG, Poindron P, Lindsay DR (1988). A high level of nutrition during late pregnancy improves subsequent maternal behaviour in Merino ewes. *Proceedings of the New Zealand Society of Animal Production* 17:294-297.
- Raboisson D, Mounie M, Maigne E (2014). Diseases, reproductive performance, and changes in milk production associated with subclinical ketosis in dairy cows: a meta-analysis and review. *Journal of Dairy Science* 97:7547-7563.
- Radostits OM, Gay CC, Hinchcliff KW, Constable PD (2007). *Veterinary Medicine*, 10th edn. Saunders, Edinburgh.

- Rajajee V, Fletcher JJ, Pandey AS, Gemmete JJ, Chaudhary N, Jacobs TL, Thompson BG (2012). Low pulsatility index on transcranial Doppler predicts symptomatic large-vessel vasospasm after aneurysmal subarachnoid hemorrhage. *Neurosurgery* 70:1195-1206.
- Regnault TR, Oddy HV, Nancarrow C, Sriskandarajah N, Scaramuzzi RJ (2004). Glucose-stimulated insulin response in pregnant sheep following acute suppression of plasma non-esterified fatty acid concentrations. *Reproductive Biology and Endocrinology* 2:64.
- Rehbein S, Kokott S, Lindner T (1999). Evaluation of techniques for the enumeration of *Dicrocoelium* eggs in sheep faeces. *Journal of Veterinary Medicine A* 46:133-139.
- Reid RL (1960). Studies on the carbohydrate metabolism of sheep. XI. The role of the adrenals in ovine pregnancy toxemia. *Australian Journal of Agricultural Research* 11:364-382.
- Reid RL (1968). The pathophysiology of undernourishment in pregnant sheep with particular reference to pregnancy toxemia. *Advances in Veterinary Science* 72:163-238.
- Reksen O, Grohn YT, Havrevoll O, Bolstad T, Waldmann A, Ropstad E (2002). Relationships among milk progesterone, concentrate allocation, energy balance, milk yield and conception rate in Norwegian cattle. *Animal Reproduction Science* 73:169-184.
- Robciuc MR, Tahvanainen E, Jauhiainen M, Ehnholm C (2010). Quantitation of serum angiopoietin-like proteins 3 and 4 in a Finnish population sample. *Journal of Lipid Research* 51:824-831.
- Robinson JJ (1986). Changes in body composition during pregnancy and lactation. *Proceedings of the Nutrition Society* 45:71-80.
- Roger PA (2008). The impact of disease and disease prevention on sheep welfare. *Small Ruminant Research* 76:104-111.
- Rollin E, Berghaus RD, Rapnicki P, Godden SM, Overton MW (2010). The effect of injectable butaphosphan and cyanocobalamin on postpartum serum beta-hydroxybutyrate, calcium, and phosphorus concentrations in dairy cattle. *Journal of Dairy Science* 93:978-987.
- Rook JS (2000). Pregnancy toxemia of ewes, does, and beef cows. *Veterinary Clinics of North America Food Animal Practice* 16:293-317.
- Rosenfeld CR, Barton MD, Meschia G (1976). Effects of epinephrine on distribution of blood flow in the pregnant ewe. *American Journal of Obstetrics and Gynaecology* 124:156-163.
- Rossi R, Dörig S, Del Prete E, Scharrer E (2000). Suppression of feed intake after parenteral administration of D-b-hydroxybutyrate in pygmy goats. *Journal of Veterinary Medicine A* 47:9-16.

- Rovensky J, Vidas M, Marek J, Blazickova S, Korcakova L, Vyletelkova L, Takac A (1991). Evidence for immunomodulatory properties of prolactin in selected in vitro and in vivo situations. *International Journal of Immunopharmacology* 13:267-272.
- Rubin J, Adler R, Fowkles J, Spratt R, Pallister J, Chen JF, Carson P (1995a). Fractional moving blood volume: estimation with power Doppler US. *Radiology* 197:183-190.
- Rubin J, Bude R, Carson P, Bree R, Adler R (1995b). Power Doppler US: a potentially useful alternative to mean frequency-based color Doppler US. *Radiology* 190:853-856.
- Rubin J, Bude R, Fowkles J, Spratt R, Carson P, Adler R (1997). Normalizing fractional moving blood volume estimates with power Doppler US: defining a stable intravascular point with the cumulative power distribution function. *Radiology* 205:757-765.
- Russel AJF (1984). Means of assessing the adequacy of nutrition of pregnant ewes. *Livestock Production Science* 11:429-436.
- Sahoo SS, Patra RC, Behera PC, Swarup D (2009). Oxidative stress indices in the erythrocytes from lactating cows after treatment for subclinical ketosis with antioxidant incorporated in the therapeutic regime. *Veterinary Research Communications* 33:281-290.
- Saratsis P, Leontides L, Tzora A, Alexopoulos C, Fthenakis GC (1998). Incidence risk and aetiology of mammary abnormalities in dry ewes in 10 flocks in Southern Greece. *Preventive veterinary Medicine* 37:173-183.
- Sargison ND (1995). Recent advances in the diagnosis, prognosis and treatment of ovine pregnancy toxemia. *Proceedings of Meetings of the Sheep Veterinary Society* 19:27-32.
- Sargison ND (2007). Pregnancy toxemia. In: Aitken ID (ed.) *Diseases of Sheep*, 4th edn. Blackwell Publishing, Oxford, pp. 359–363.
- Sargison N (2008). *Sheep Flock Health. A Planned Approach*. Blackwell, Oxford.
- Sargison ND, Scott PR, Penny CD, Pirie RS, Kelly JM (1994). Plasma enzymes and metabolites as potential prognostic indices of ovine pregnancy toxemia – a preliminary study. *British veterinary Journal* 150:271-277.
- Sartorelli P, Paltrinieri S, Agnes F (1999). Non-specific immunity and ketone bodies. I: in vitro studies on chemotaxis and phagocytosis in ovine neutrophils. *Zentralblatt fur Veterinarmedizin* 46:613-619.
- Sartorelli P, Paltrinieri S, Comazzi S (2000). Non-specific immunity and ketone bodies. II: in vitro studies on adherence and superoxide anion production in ovine neutrophils. *Zentralblatt fur Veterinarmedizin* 47:1-8.
- Schalm OW, Carrol EJ, Jain NC (1971). *Bovine Mastitis*. Lea & Febiger, Philadelphia, USA.

- Scharrer E (1999). Control of food intake by fatty acid oxidation and ketogenesis. *Nutrition* 15:704-714.
- Schlumbohm C, Harmeyer J (1990). Hypocalcemia reduces rate of disappearance of glucose from plasma. *Journal of veterinary Medicine A* 37:285-293.
- Schlumbohm C, Harmeyer J (1999). Effect of hypocalcemia on glucose metabolism in hyperketonemic piglets. *Experimental Physiology* 84:707-723.
- Schlumbohm C, Harmeyer J (2003). Hypocalcemia reduces endogenous glucose production in hyperketonemic sheep. *Journal of Dairy Science* 86:1953-1962.
- Schlumbohm C, Harmeyer J (2004). Hyperketonemia impairs glucose metabolism in pregnant and nonpregnant ewes. *Journal of Dairy Science* 87:350-358.
- Schwartz MW, Woods SC, Porte D, Seeley RJ, Baskin DG (2000). Central nervous system control of food intake. *Nature* 404:661-671.
- Scott PR (1992). Analysis of cerebrospinal fluid from field cases of some common ovine neurological diseases. *British veterinary Journal* 148:15-22.
- Scott PR, Sargison ND, Penny CD, Strachen WD (1995). Aqueous humor and cerebrospinal fluid collected at necropsy as indicators of antemortem serum 3-OH butyrate concentration in pregnant sheep. *British veterinary Journal* 151:459-461.
- Sharp JM, Nettleton PF (2007). Acute respiratory virus infections. In: Martin WB, Aitken ID (eds) *Diseases of Sheep*, 4th edn. Blackwell, Oxford, pp. 177-180.
- Shimamura M, Matsuda M, Kobayashi S, Ando Y, Ono M, Koishi R, Furukawa H, Makishima M, Shimomura I (2003). Angiotensin-like protein 3, a hepatic secretory factor, activates lipolysis in adipocytes. *Biochemical and Biophysical Research Communications* 301:604-609.
- Sigurdsson H (1988). The effects of pregnancy and feeding on the insulin and glucose concentrations in blood of ewes in late pregnancy. *Acta veterinaria scandinavica* 29:401-405.
- Simensen MG (1971). Investigations on pregnancy toxæmia in sheep. *Nordisk veterinær Medicin* 23:99-113.
- Smith MC, Sherman DM (2009). *Goat Medicine*, 2nd edn. Wiley-Blackwell, Iowa.
- Sobiraj A (1994). Dystocia in sheep and goats. A survey of seven lambing periods at a clinic for obstetrics. *Deutsche Tierärztliche Wochenschrift* 101:471-476.
- Sordillo LM, Babiuk LA (1991a). Controlling acute *Escherichia coli* mastitis during the periparturient period with recombinant bovine interferon gamma. *Veterinary Microbiology* 28:189-198.

- Sordillo LM, Babiuk LA (1991b). Modulation of bovine neutrophil function during the periparturient period following in vitro exposure to recombinant bovine interferon gamma. *Veterinary Immunology and Immunopathology* 27:393-402.
- Sordillo LM, Snider M, Hughes H, Afseth G, Campos M, Babiuk LA (1991). Pathological changes in bovine mammary glands following intramammary infusion of recombinant interleukin-2. *Journal of Dairy Science* 74:4164-4174.
- Sordillo LM, Streicher KL (2002). Mammary gland immunity and mastitis susceptibility. *Journal of mammary Gland Biology and Neoplasia* 7:135-146.
- Sun YH, Wang B, Shu S, Zhang H, Xu C, Wu L, Xia C (2015). Critical thresholds of liver function parameters for ketosis prediction in dairy cows using receiver operating characteristic (ROC) analysis. *Veterinary Quarterly* 35:159-164.
- Suriyasathaporn W, Daemen AJ, Noordhuizen- Stassen EN, Dieleman SJ, Nielen M, Schukken YH (1999). Beta-hydroxybutyrate levels in peripheral blood and ketone bodies supplemented in culture media affect the in vitro chemotaxis of bovine leukocytes. *Veterinary Immunology and Immunopathology* 68:177-186.
- Suriyasathaporn W, Schukken YH, Nielen M, Brand A (2000). Low somatic cell count: a risk factor for subsequent clinical mastitis in a dairy herd. *Journal of Dairy Science* 83:1248-1255.
- Svennersten-Sjaunja K, Olsson K (2005). Endocrinology of milk production. *Domestic Animal Endocrinology* 29:241-258.
- Swanson TJ, Hammer CJ, Luther JS, Carlson DB, Taylor JB, Redmer DA, Neville TL, Reed JJ, Reynolds LP, Caton JS, Vonnahme KA (2008). Effect of gestational plane of nutrition and selenium supplementation on mammary development and colostrum quality in pregnant ewe lambs. *Journal of Animal Science* 86:2415-2423.
- Taylor MA (2010). Parasitological examinations in sheep health management. *Small Ruminant Research* 92:120-125.
- Temizel EM, Batmaz H, Keskin A, Orman A, Gencoglu H, Catik S, Topal O (2015). Butaphosphan and cyanocobalamin treatment of pregnant ewes: metabolic effects and potential prophylactic effect for pregnancy toxemia. *Small Ruminant Research* 125:163-172.
- Tharwat F, Al-Sobayil (2014). Cord and jugular blood acid–base and electrolyte status and haematobiochemical profiles in goats with naturally occurring pregnancy toxemia. *Small Ruminant Research* 117:73-77.

- Tharwat M, Al-Sobayil F, Al-Sobayil K (2012). The cardiac biomarkers troponin I and CK-MB in nonpregnant and pregnant goats, goats with normal birth, goats with prolonged birth, and goats with pregnancy toxemia. *Theriogenology* 78:1500-1507.
- Thivierge MC, Petitclerc D, Bernier JF, Couture Y, Lapierre H (2000). External pudic venous reflux into the mammary vein in lactating dairy cows. *Journal of Dairy Science* 83:2230-2238.
- Thomas GB, Mercer JE, Karalis T, Rao A, Cummins JT, Clarke IJ (1990). Effect of restricted feeding on the concentrations of growth hormone (GH), gonadotropins, and prolactin (PRL) in plasma, and on the amounts of messenger ribonucleic acid for GH, gonadotropin subunits, and PRL in the pituitary glands of adult ovariectomized ewes. *Endocrinology* 126:1361-1367.
- Thomas L, Wallace JM, Aitken RP, MercerJG, Trayhurn P, Hoggard N (2001). Circulating leptin during ovine pregnancy in relation to maternal nutrition, body composition and pregnancy outcome. *Journal of Endocrinology* 169:465-476.
- Tucker HA (1981). Physiological control of mammary growth, lactogenesis and lactation. *Journal of Dairy Science* 64:1403-1407.
- Twardock AR, Symonds HW, Sansom BF, Rowlands GJ (1973). The effect of litter size upon foetal growth rate and the placental transfer of calcium and phosphorus in superovulated Scottish half bred ewes. *British Journal of Nutrition* 29:437-446.
- Van Saun RJ (2000). Pregnancy toxemia in a flock of sheep. *Journal of the american veterinary medical Association* 217:1536-1539.
- Vanhoutte PM, Shimokawa H, Tang EHC, Feletou M (2009). Endothelial dysfunction and vascular disease. *Acta physiologica* 196:193-222.
- Varnam GC, Jeacock MK, Shepherd DA (1978). Hepatic ketonebody metabolism in developing sheep and pregnant ewes. *British Journal of Nutrition* 40:359-367.
- Verbeek E, Oliver MH, Waas JR, McLeay LM, Blache D, Matthews LR (2012). Reduced cortisol and metabolic responses of thin ewes to an acute cold challenge in mid-pregnancy: implications for animal physiology and welfare. *PLOS One* 7:e37315.
- Wallace JM, Bourke DA, Aitken RP, Cruickshank MA (1999). Switching maternal dietary intake at the end of the first trimester has profound effects on placental development and fetal growth in adolescent ewes carrying singleton fetuses. *Biology of Reproduction* 61:101-110.

- Wang Y, Wu Y, He E, Zeng D, Shen J, Lu Y, Zhang G, Zhu Q, Li S, Tian X, Gong X, Miao X, Li X, Schumacher J, Moldenhauer M, Tate M, Lin L (2013). Glucose tolerance test and clinical chemistry in male Beagle dogs. *International Journal of Toxicology* 32:62-63.
- Wathes DC, Reynolds TS, Robinson RS, Stevenson KR (1998). Role of the insulin-like growth factor system in uterine function and placental development in ruminants. *Journal of Dairy Science* 81:1778-1789.
- Wentink GH, Rutten VP, van den Ingh TS, Hoek A., Muller KE, Wensing T (1997). Impaired specific immunoreactivity in cows with hepatic lipidosis. *Veterinary Immunology and Immunopathology* 56:77-83.
- Widder B, Goertler M (2004). Grundlagen. In: Widder B, Goertler M (eds) *Doppler- und Duplexsonographie der hirnersorgenden Arterien*, 6th edn. Springer, Berlin, pp. 3-86.
- Wood M., Romine LE, Lee YK, Richman KM, O'Boyle NMK, Paz DA., Chu PK, Pretorius DH (2010). Spectral Doppler signature waveforms in ultrasonography: a review of normal and abnormal waveforms. *Ultrasound Quarterly* 26:83-99.
- Woods SC (2005). Signals that influence food intake and body weight. *Physiology and Behavior* 86:709-716.
- Woods SC, Seeley RJ (2000). Adiposity signals and the control of energy homeostasis. *Nutrition* 16:894-902.
- Zamir S, Rozov A, Gootwin E (2009). Treatment of pregnancy toxemia in sheep with flunixin meglumine. *The veterinary Record* 165:265-266.