



**ΠΑΝΕΠΙΣΤΗΜΙΟ ΘΕΣΣΑΛΙΑΣ**  
**ΣΧΟΛΗ ΕΠΙΣΤΗΜΩΝ ΥΓΕΙΑΣ**  
**ΤΜΗΜΑ ΙΑΤΡΙΚΗΣ**



Εργαστήριο Βιομαθηματικών

Διευθυντής: Ζιντζαράς Ηλίας, Καθηγητής Τμήματος Ιατρικής ΠΘ

---

**Διδακτορική Διατριβή**

**«Δίκτυα θεραπειών από τυχαιοποιημένες κλινικές μελέτες:**

**μεθοδολογική προσέγγιση»**

υπό

**Γεώργιου Μπάκαλου**

MD, MSc

Υπεβλήθη για την εκπλήρωση μέρους των

απαιτήσεων για την απόκτηση του

Διδακτορικού Διπλώματος

Λάρισα, 2016

© 2016 Georgios Bakalos

Η έγκριση της διδακτορικής διατριβής από το Τμήμα Ιατρικής της Σχολής Επιστημών Υγείας του Πανεπιστημίου Θεσσαλίας δεν υποδηλώνει αποδοχή των απόψεων του συγγραφέα (σύμφωνα με τις διατάξεις του άρθρου 202, παράγραφος 2 του Ν.5343/1932).

**Εγκρίθηκε από τα Μέλη της Επταμελούς Εξεταστικής Επιτροπής (6<sup>η</sup>/20-01-2016 ΓΣΕΣ):**

<b>1<sup>ος</sup> Εξεταστής (Επιβλέπων)</b>	<b>Ζιντζαράς Ηλίας</b> Καθηγητής Τμήματος Ιατρικής ΠΘ
<b>2<sup>ος</sup> Εξεταστής</b>	<b>Στεφανίδης Ιωάννης</b> Καθηγητής Τμήματος Ιατρικής ΠΘ
<b>3<sup>ος</sup> Εξεταστής</b>	<b>Παπανδρέου Χρήστος</b> Καθηγητής Τμήματος Ιατρικής ΠΘ
<b>4<sup>ος</sup> Εξεταστής</b>	<b>Νταφόπουλος Κων/νος</b> Αν. Καθηγητής Τμήματος Ιατρικής ΠΘ
<b>5<sup>ος</sup> Εξεταστής</b>	<b>Κυριάκου Δέσποινα</b> Καθηγήτρια Τμήματος Ιατρικής ΠΘ
<b>6<sup>ος</sup> Εξεταστής</b>	<b>Χατζηγεωργίου Γεώργιος</b> Καθηγητής Τμήματος Ιατρικής ΠΘ
<b>7<sup>ος</sup> Εξεταστής</b>	<b>Ζαχαρούλης Δημήτριος</b> Καθηγητής Τμήματος Ιατρικής ΠΘ

## **ACKNOWLEDGMENTS**

I would like to express my sincere gratitude to my advisor, Prof.Elias Zintzaras, for his continuous guidance, mentorship, and support in overcoming numerous obstacles.

Besides my advisor, I would like to thank the rest of my thesis committee, for their encouragement and insightful comments.

My sincere thanks also go to Mr. Theodoros Mprotsis, for his support on the statistical analyses performed for this exercise.

Last, but not least, I would like to thank Mrs. Katerina Kalogianni for the valuable administrative support provided during my research.

## **CURRICULUM VITAE**

George Bakalos is a Medical Doctor with postgraduate studies in Health Crisis Management and Pharmaco-Epidemiology. He has more than 10 years of experience in pharmaceutical drug development, including senior positions in clinical research and pharmacovigilance. He joined Roche Greece in 2009 as Medical Manager in charge of the Hepatitis group. Since 2013, he moved with his family to Basel, Switzerland, where he works as International Medical Director at Global Medical Affairs Department, responsible for late stage drug development in virology, and most recently in oncology.

**«Δίκτυα θεραπειών από τυχαιοποιημένες κλινικές μελέτες:  
μεθοδολογική προσέγγιση»**

**Γεώργιος Μπάκαλος**

Πανεπιστήμιο Θεσσαλίας, Τμήμα Ιατρικής, 2016

**ΤΡΙΜΕΛΗΣ ΣΥΜΒΟΥΛΕΥΤΙΚΗ ΕΠΙΤΡΟΠΗ**

1. **Ζιντζαράς Ηλίας**, Καθηγητής Τμήματος Ιατρικής ΠΘ (**Επιβλέπων**)
2. **Στεφανίδης Ιωάννης** Καθηγητής Τμήματος Ιατρικής ΠΘ
3. **Παπανδρέου Χρήστος** Καθηγητής Τμήματος Ιατρικής ΠΘ

## **Abstract**

### **Network Meta-analysis of randomized control trials - A methodological perspective**

**Objective** The aim of this thesis was to explore methodological aspects related to network meta-analysis, especially in oncology field. For the purposes of this analysis, Small Cell Lung Cancer (SCLC) was used as an example, where the combination of Cisplatin plus Etoposide (EP) is currently the standard treatment.

**Methods** PubMed, EMBASE and Cochrane Central Register of Controlled Trials were systematically searched to identify all RCTs that compared treatments for SCLC. Then, effectiveness of the treatments relative to the combination of Cisplatin plus Etoposide, reference treatment) was estimated by performing a network of treatments analysis, using both Bayesian and frequentist approaches.

**Results** We identified 71 articles eligible for inclusion, involving 91 different treatments. In total, 16,026 patients were included in the analysis. Frequentist analysis (direct) revealed combination of Cisplatin plus Cyclophosphamide plus Etoposide plus Epirubicin showed better response than EP for the ORR outcome, but with worse tolerability. Indirect analysis revealed that the combination of Cisplatin plus Doxorubicin plus Etoposide (plus Vincristine) showed better response than EP for the ORR outcome. Bayesian analysis revealed that the combination of carboplatin or cisplatin plus etoposide with granulocyte-colony stimulating factor (GCSF) provides higher probability of achieving ORR compared to other treatments.

**Conclusions** The results should be interpreted with caution because the network was dominated by indirect comparisons. Large scale head-to-head RCTs are needed to confirm the present findings. Bayesian and frequentist approaches should be considered complementary tools in the clinical evaluator's toolkit.

## ***Table of Contents***

Chapter 1 - Overview of Network Meta-Analysis .....	17
1.1 Introduction .....	17
1.2 Attractiveness of NMA .....	19
1.3 Rationale for the research.....	21
Chapter 2 – Methodological consideration for network meta-analyses.....	22
2.1 Assumptions and validity considerations: transitivity and consistency .....	22
2.1.2 Transitivity and Similarity .....	22
2.1.2 Consistency.....	24
2.2 Network Geometry .....	24
2.3 Statistical Methodology.....	27
2.4 Internal and External Validity of NMA .....	29
Chapter 3 – Overview of treatment options in Small Cell Lung Cancer .....	30
3.1 Overview .....	30
3.2 LD-SCLC .....	31
3.3 ED-SCLC .....	32
3.4 Current issues in pharmacotherapy .....	33
3.4.1 Cisplatin versus carboplatin.....	33
3.4.2 First-line regimens other than platin plus etoposide.....	34
3.4.3 New medicinal products .....	38
3.4.4 Targeted Therapies .....	40
Chapter 4 – Selection of an appropriate endpoint to assess relative effectiveness of treatments in SCLC .....	43
Chapter 5 – Assessing the relative effectiveness of treatments in SCLC - Frequentist Approach.....	48
<i>Bakalos G et al. Assessing the relative effectiveness and tolerability of treatments in small cell lung cancer: a network meta-analysis. Cancer Epidemiol. 2013 Oct;37(5):675-82.....</i>	<i>48</i>
5.1 Introduction .....	48
5.2 Materials & Methods.....	48
5.2.1 Search strategy-Selection of RCTs.....	48
5.2.2 Eligibility criteria.....	50
5.2.3 Data extraction and outcomes definition .....	51
5.2.4 Treatment definition .....	51
5.2.5 Statistical methods – Frequentist Analysis.....	52



5.3 Results .....	53
5.3.1 Eligible studies and summary characteristics .....	53
5.3.2 The networks .....	55
5.3.3 Direct analysis for comparing treatments with EP .....	57
5.3.4 Sub-analysis of most common interventions in SCLC.....	61
Chapter 6 – Assessing the relative effectiveness of treatments in SCLC - Bayesian Approach.....	64
6.1 Introduction .....	64
6.2 Results .....	65
6.2.1 Sub analysis of most common interventions in SCLC .....	65
6.2.2 Platin-pooled data sub-analysis in ED-SCLC .....	70
6.2.3 Sub-analysis – New treatments .....	77
Charter 7 – Discussion .....	79
Charter 8 – Conclusions.....	84
References.....	85
Appendices.....	103
Supplementary Tables .....	103
Supplementary Figures.....	135
Manuscripts .....	138

## ***List of Tables***

Table 1: Networks published between 2005 and 2014 .....	19
Table 2: Networks in Oncology published between 2005 and 2014 .....	20
Table 3: Phase III trials comparing platinum-based regimens: carboplatin or cisplatin.....	34
Table 4: Randomized trials of cisplatin or carboplatin plus irinotecan or topotecan in ED-SCLC .....	38
Table 5: Targeted therapies in first line treatment of small cell lung cancer.....	40
Table 6: Targeted therapies in treatment of relapsed small cell lung cancer.....	41
Table 7: ED-SCLC studies used to estimate correlation coefficient and equation of the regression line.....	45
Table 8: Characteristics of 71 included randomized controlled trials (RCTs) in the network of treatments meta-analysis.....	53
Table 9: Direct analysis results for comparing treatments for small cell lung cancer (SCLC) with reference treatment (Cisplatin + Etoposide, EP) by outcome. The treatments were sorted according to their significance and magnitude of effect size.....	58
Table 10: Indirect analysis results for comparing treatments for small cell lung cancer (SCLC) with reference treatment (Cisplatin + Etoposide) by outcome, for treatments that produced significantly ( $P < 0.05$ ) different response than reference treatment .....	59
Table 11: Network Characteristics.....	62
Table 12: Intervention Characteristics .....	62
Table 13: Direct Comparison Characteristics .....	63
Table 14: Rankogram of most common interventions in SCLC, FE.....	67
Table 15: Rankogram of most common interventions in SCLC, RE .....	68
Table 16: SUCRA of most common interventions in SCLC.....	69
Table 17: Network characteristics of platin-pooled data sub-analysis in ED-SCLC.....	72
Table 18: Intervention characteristics of platin-pooled data sub-analysis in ED-SCLC .....	72
Table 19: Direct Comparison Characteristics of platin-pooled data sub-analysis in ED-SCLC .....	72
Table 20: League table of platin-pooled data sub-analysis in ED-SCLC - FE.....	74
Table 21: League table of platin-pooled data sub-analysis in ED-SCLC - RE .....	74
Table 22: SUCRA of platin-pooled data sub-analysis in ED-SCLC .....	75
Table 23: Network characteristics of platin-pooled data sub-analysis compared to investigational treatments in ED-SCLC .....	77
Table 24: SUCRA of platin-pooled data sub-analysis compared to investigational treatments in ED-SCLC .....	78

## ***List of Figures***

Figure 1: Direct evidence comes from the synthesis of trials of A versus B. Indirect evidence comes through an intermediate/common comparator C (many intermediate comparators are possible) by combining trials of A versus C and of C versus B (prior to combination the trials were synthesized).....	18
Figure 2: Methods used to synthesize the data in relation to the shape of the network [18] ...	20
Figure 3: Examples of treatment networks. ....	26
Figure 4: Surrogate endpoints for long-term effectiveness in oncology/hematology.....	43
Figure 5: Equation of the regression line .....	46
Figure 6: Flow diagram of the screening process and RCTs selection for multiple-treatments meta-analysis of treatments for SCLC.....	49
Figure 7: Network of direct comparisons for the outcome “Objective Response Rate”. ....	56
Figure 8: Network Diagram of most common interventions .....	61
Figure 9: Forrest Plot of most common interventions in SCLC .....	66
Figure 10: Stacked Bar Chart of most common interventions in SCLC.....	68
Figure 11: Inconsistency results of most common interventions in SCLC .....	69
Figure 12: Network diagram of platin-pooled data sub-analysis in ED-SCLC .....	71
Figure 13: Forest plot of platin-pooled data sub-analysis in ED-SCLC .....	73
Figure 14: Rankogram of platin-pooled data sub-analysis in ED-SCLC .....	75
Figure 15: Inconsistency results of platin-pooled data sub-analysis in ED-SCLC.....	76
Figure 16: Rankogram of platin-pooled data sub-analysis compared to investigational treatments in ED-SCLC.....	78

### ***List of Supplementary Tables***

Supplementary Table 1: List of network meta-analyses in oncology reported 2006-2014 ...	103
Supplementary Table 2: The characteristics of the individual RCTs included in network meta-analysis for assessing the relative effectiveness of treatments in SCLC.....	109
Supplementary Table 3: Quality assessment of of the individual RCTs included in network meta-analysis for assessing the relative effectiveness of treatments in SCLC.....	118
Supplementary Table 4: Proportion of reporting of 24 data items in a total of 81 randomized clinical trials in small cell lung cancer by publication period (pre- and post-CONSORT and combined)* .....	122
Supplementary Table 5: Definition of treatments.....	125
Supplementary Table 6: Results of all direct and indirect comparisons.....	128
Supplementary Table 7: Common grade 3-4 hematological toxicities described by the included RCTs in the network meta-analysis, according to the grouping of chemotherapy treatments.....	131
Supplementary Table 8: Additional studies identified after the initial literature research ....	133

### ***List of Supplementary Figures***

Supplementary Figure 1: Network of treatments for the outcome “patients with complete response”. .....	135
Supplementary Figure 2: Network of treatments for the outcome “patients with neutropenia”. .....	136
Supplementary Figure 3: Network of treatments for the outcome “patients with febrile neutropenia”.....	137

### ***List of Publications***

<i>Bakalos G, Miligos M, Doxani C et al. Assessing the relative effectiveness and tolerability of treatments in small cell lung cancer: a network meta-analysis. Cancer Epidemiol. 2013 Oct;37(5):675-82.....</i>	138
---	-----

## ***Glossary***

Adjusted indirect comparison	A statistical technique that permits comparison between two interventions that have not been compared directly (head-to-head) but have both been compared to the same third comparator. This method preserves the principle of randomization.
Bayesian analysis	A statistical method that employs prior knowledge combined with data.
Closed loop	A network of 3 comparisons, each of which has been compared directly with the others.
Consistency or coherence:	The manifestation of transitivity in the data from a network of interventions. It exists when treatment effects from direct and indirect evidence are in agreement (subject to the usual variation due to heterogeneity in the direct evidence). Unlike transitivity, consistency can be evaluated statistically.
(In)coherence	The (dis)agreement in treatment effect estimates between direct and indirect evidence.
Co-occurrence	The over-representation of RCTs comparing specific interventions rather than other available interventions.
Credible intervals	Bayesian analogy to confidence intervals
Direct (head-to-head) evidence	Data from RCTs that have compared interventions against each other.
Diversity of a network	A measure of how many treatments are available and whether they are equally represented or not across the network
Fixed effects analysis	A method of analysis that assumes that treatment effects are the same across all included trials
Frequentist analysis	A statistical approach that places the emphasis on available data (conventional approach to statistical analysis, contrast with Bayesian).
Geometry of a network	A graphical representation of the distribution of treatments and their comparisons across the network
Heterogeneity	The extent of inconsistency of treatment effects in a pairwise meta-analysis.
Homogeneity	The inverse of heterogeneity
Inconsistent loop	A closed loop in which treatment effect estimates from direct and indirect evidence are in statistically significant disagreement.
Indirect evidence	Evidence bearing on the relative effect of treatments that have not been compared directly against each other but have a common comparator. Indirect evidence may be evaluated using accepted statistical approaches, including adjusted indirect comparisons and MTCs.

Lumley model of network meta-analysis	A frequentist method for MTCs.
Meta-regression	A regression in which the dependent variable is the magnitude of treatment effect in individual studies and the independent variable are study characteristics. It is used to see if study characteristics can explain differences in magnitude of treatment effect across studies.
Network meta-analysis	Synthesis of information over a network of comparisons to assess the comparative effectiveness of more than 2 alternative treatment options for the same condition. The method relies on mixed comparison and synthesizes direct and indirect evidence over the entire network to obtain the relative treatment effects for all comparisons and a ranking of the treatments.
ORR	ORR is defined as the proportion of patients with tumor size reduction of a predefined amount and for a minimum time period. Response duration usually is measured from the time of initial response until documented tumor progression. According to FDA, ORR is defined as the sum of partial responses plus complete responses. When defined in this manner, ORR is a direct measure of drug antitumor activity.
OS	Overall survival is defined as the time from randomization until death from any cause, and is measured in the intent-to-treat population. Survival is considered the most reliable cancer endpoint, and when studies can be conducted to adequately assess survival, it is usually the preferred endpoint.
PFS	Progression-free survival (PFS) is defined as the time from random assignment in a clinical trial to disease progression or death from any cause, , and is measured in the intent-to-treat population
Priors (informed)	The representation of external (prior) knowledge about the intervention effects or degree of heterogeneity that is incorporated in Bayesian analysis.
Priors (non-informative)	In Bayesian analysis, the assumption that nothing is known about the intervention effect or degree of heterogeneity prior to looking at the available data
Posterior distribution	In Bayesian analysis, the probability distribution obtained by mixing prior knowledge with data
Random effects analysis	A method of analysis that incorporates variation between trials and allows the treatment effect to vary across the included trials.

Similarity	A principle that all the trials are broadly similar with respect to populations, trial design and outcomes. Used interchangeably with the more statistical term “exchangeability.”
Star Network	A network where all treatments have been compared with a common comparator treatment but not between themselves. Statistical evaluation of consistency is impossible in a star network.
Treatment rankings	Ordering of treatments according to decreasing probability that they can produce better outcomes than competing interventions

## *Abbreviations*

ASCO	American Society of Clinical Oncology
CI	Confidence Interval
CR	Complete Response
ECOG	Eastern Cooperative Oncology Group
ED	Extensive-Stage Disease
ESMO	European Society for Medical Oncology
FDA	Food and Drug Administration
FE	Fixed Effects
HR	Hazard Ratio
LD	Limited-Stage Disease
MA	Meta-Analyses
MR	Meta Regression
mo(s).	Month(s)
N	Number
NCI	National Cancer Institute
NMA	Network Meta-Analyses
OR	Odds Ratio
ORR	Overall Response Rate
OS	Overall Survival
PFS	Progression-Free Survival
RCT	Randomized Controlled Trials
Q	Heterogeneity Statistic
QoL	Quality of Life
PR	Partial Response
RCT	Randomized Controlled Trial
RD	Risk Difference
RE	Random Effects
RR	Risk Ratio
SCLC	Small Cell Lung Cancer
SD	Stable Disease
SE	Standard Error
SUCRA	Surface Under the Cumulative Ranking
WHO	World Health Organization
wk(s)	Week(s)
yr(s)	Year(s)



# Chapter 1 - Overview of Network Meta-Analysis


## 1.1 Introduction

Meta-analysis is a statistical technique to synthesize information from a collection of all relevant studies comparing the same intervention for a medical condition of interest to address a focused research question in the context of a systematic review [1, 2]. Systematic reviews and meta-analyses of randomized controlled trials (RCTs) are fundamental tools for the clinical practice and are placed at the peak of evidence based medicine pyramid, influencing significantly the decision making of clinicians, scientists and policy makers. When conducted well, and transparently reported, systematic reviews and meta-analyses produce information that can be helpful for the evaluation of healthcare interventions.

However, a potential drawback of traditional meta-analyses is their ability to compare only two interventions a time, a significant obstacle for decision-making when the medical condition under study has many relevant treatment options to be considered in clinical practice [3–8].

Moreover, in situations where there are no studies directly comparing two or more interventions, traditional meta-analysis cannot estimate their comparative benefits and harms. A simple example of this scenario is when information from RCTs is available regarding the effectiveness of two active treatments, generically “A” and “B”, in comparison to a common comparator “C” (commonly placebo or standard of care practice); an indirect treatment comparison may be used to estimate a comparison of the relative effectiveness of “A” compared with “B” (Figure 1).

**Figure 1: Direct evidence comes from the synthesis of trials of A versus B. Indirect evidence comes through an intermediate/common comparator C (many intermediate comparators are possible) by combining trials of A versus C and of C versus B (prior to combination the trials were synthesized).**



In that case, an indirect comparison of “A” versus “B” can be obtained by synthesizing the results of the trials providing information on the direct comparisons of “A” versus “C” and “C” versus “B”, by means of so-called network meta-analysis (NMA).

NMA also allows for the combination of direct and indirect information in the estimation of a single treatment effect, which has come to be known as a mixed comparison. In this example, the mixed comparison between “A” and “B” incorporates the results of the direct comparison of the two treatments (i.e., outcome data from RCTs of “A” vs. “B”) with indirect comparison results obtained from the information of other related direct comparisons (i.e., outcome data from RCTs of “A” vs. “C” and “C” vs. “A”).

In practice, mixed comparisons can be estimated as a weighted average of the direct and indirect estimates of treatment effects, and they can complement information for those comparisons in which there is scarce direct information. Both direct and indirect evidence contribute to the total body of evidence. Statistical methods for comparing multiple interventions (using a Bayesian or frequentist framework) have been described extensively in the literature [9–12].

## 1.2 Attractiveness of NMA


Network meta-analysis (NMA) is becoming increasingly popular for evidence synthesis [13–16] and is evolving to the ‘new norm’ for comparative effectiveness research [17, 18]. Most recently, Petropoulou et al identified 456 NMAs being published between 1999 and 2015 by searching three bibliographic databases and assessed the characteristics of their statistical analysis and reporting of results [18]. Between 1999 and 2004 only 6 NMAs were published (1 in 1999, 2 in 2000, 1 in 2003 and 2 in 2004). The number of NMAs published per year after 2004 is presented in Table 1. It is evident that the number of published studies applying NMA methods to clinical research questions has been increasing significantly over the last two decades ( $p=0.04$ ). [18]

**Table 1: Networks published between 2005 and 2014**

2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
6	12	9	12	27	30	53	59	96	103

The quality and transparency of reporting also increased: in recent years around 90% of articles clearly reported whether a random-effects or fixed-effect model was used, and in 2015 all reports included a description of the statistical methods used. It should be noted that the Bayesian hierarchical model remained the most popular approach for NMA during the study period: only five articles reported the use of frequentist multivariate meta-analysis or meta-regression [Figure 2].

**Figure 2: Methods used to synthesize the data in relation to the shape of the network [18]**



We performed a similar analysis focusing only on oncology studies [Appendix Table 1] and a similar effect was observed [Table 2].

Specifically there is a significant increase of NMAs in oncology/hematology after 2010, reaching up to 25% of the total NMA reported in 2014.

**Table 2: Networks in Oncology published between 2005 and 2014**

2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
0/6	1/12	1/9	2/12	4/27	0/30	5/53	8/59	14/96	24/103

*Note: Literature Research String: (network OR mixed treatment\* OR multiple treatment\* OR mixed comparison\* OR indirect comparison\* OR umbrella OR simultaneous comparison\*) AND (meta-analysis) AND (oncology); Literature Research Timelines: up to April 14, 2015*

In the lack of many direct comparisons, NMA is anticipated to play a key role in evaluating established treatments but also comparing innovative treatments to standard of care.

For the purposes of this exercise, Small Cell Lung Cancer was selected as a setting to explore methodological perspective for NMAs

### **1.3 Rationale for the research**

The aim of this thesis is to explore methodological aspects related to network meta-analysis, especially in oncology field. For the purposes of this analysis, SCLC will be used as an example.

# **Chapter 2 – Methodological consideration for network meta-analyses**

## **2.1 Assumptions and validity considerations: transitivity and consistency**

The validity of a NMA depends on a set of assumptions. The main assumption, which supports the validity of indirect and mixed comparison, is that there are no important differences between the trials making different comparisons other than the treatments being compared.

This assumption has been described alongside several terms in the literature, including similarity (19, 20), transitivity (4, 21), consistency (22), and coherence (11). These core assumptions for NMAs can be verified conceptually and epidemiologically but are, however, subject to substantial uncertainty.

### **2.1.2 Transitivity and Similarity**

The synthesis of studies making a direct comparison of two treatments is meaningful only when the studies are sufficiently similar in terms of key clinical and methodological characteristics (which are known as effect modifiers). The effect modifiers are not necessary to be identical; however heterogeneity of effects across studies should be acceptable. A valid indirect comparison (such as A versus B) requires that the sets of A versus C and B versus C studies are similar in their distributions of effect modifiers (for example, population characteristics, disease stage at baseline, treatment dose, sample size, and study quality). In that case we assume that the intervention effects are transitive. Transitivity can be viewed as the extension of clinical and methodological homogeneity to comparisons across groups of

studies that compare treatments. In complex network structures, the transitivity assumption should hold for all cases where indirect or mixed estimates are derived.

For example, let's assume that all A versus C studies include patients with Extended-Stage Disease and all BC studies include patients with Limited-Stage Disease. Each study set is similar within itself (at least according to this particular characteristic), but the two study sets deal with clinically different populations. So, if severity is an effect modifier, the transitivity assumption would not hold, and synthesis of these two meta-analyses would not give a valid A versus B estimate (21).

A special case of an effect modifier that can vary across comparisons violating the transitivity assumption is the nature of the common comparator. If comparator C is systematically different in A versus C and B versus C studies (for example, treatment C is administered as an oral tablet in A versus C studies but as a different formulation in B versus C studies), then the transitivity assumption probably might not hold, and the indirect comparison between treatments A and B might not be valid.

The plausibility of the transitivity assumption requires clinical judgment to decide whether differences in the distributions of the effect modifiers across studies are significant enough to make network meta-analysis invalid. If an imbalanced distribution of effect modifiers is identified, adjustment can be used to improve transitivity through network meta-regression (23, 24). Adjustment should take place only for study or patient characteristics that are effect modifiers (such as severity of illness at baseline, number of previous episodes, age, or gender) (25).

### **2.1.2 Consistency**

Consistency (or coherence) is the statistical manifestation of transitivity and occurs when the subtraction equation is supported by the data. It can be evaluated only when a loop in the evidence network exists, that is, when there is direct and indirect evidence for a particular comparison of interventions. The distinction between transitivity and consistency is analogous to the distinction between clinical or methodological heterogeneity and statistical heterogeneity seen in standard meta-analysis:

- Heterogeneity refers to the degree of disagreement between study-specific treatment effects and is measured by differences in estimates of study treatment effect beyond what chance can explain.
- Inconsistency refers to the degree of disagreement between source-specific (and not study-specific) treatment effects and is measured by differences between direct and indirect estimates beyond what chance can explain (4).

Heterogeneity is usually evaluated by the Cochran Q test or the I<sup>2</sup> statistic (26). Consistency in a network meta-analysis can be evaluated statistically by comparing the direct and indirect summary effects in specific loops (10, 11) or across a network by fitting models that allow and do not allow for inconsistency (22, 27, 28).

## **2.2 Network Geometry**

In principle, the graphical representation of a network, showing the multiple competing treatments, supports the understanding and the assessment of the strength of the clinical evidence for each of the various comparisons under study. It also improves the transparency of the results of a NMA from the perspective of determining the degree of confidence one may place in interpreting particular comparisons. Network geometry addresses what the shape of the treatment network looks like in terms of the number of included interventions




(i.e., “treatment nodes”), the extent to which there are trials comparing different pairs of these interventions (i.e., the adjoining lines or “edges”), and the numbers of patients associated with different comparisons [5].

By studying and presenting the network geometry, we could understand how strong the evidence is for some treatment comparisons and whether specific comparisons are over- or under-represented, or even avoided (comparator preference bias) [5, 7].

Generally, the edges between treatment nodes in the network indicate the comparisons made within eligible randomized trials identified during the process of study identification. The widths of these edges are commonly sized to proportionally reflect the numbers of studies evaluating each pair of treatments, and the sizes of each treatment node are typically sized to proportionally reflect the numbers of subjects randomized to each treatment.

Circles represent treatment nodes in the network; lines represent direct comparisons for which data are available from RCTs. Line thickness is proportionally weighted according to the number of studies evaluating each comparison, while nodes are proportionally weighted according to the number of patients that have received each treatment relative to the total number of participants across all studies. Examples of possible network geometries are presented in Fig. 3; If all of the treatments have been compared against a common comparator (e.g., placebo), but not among active treatment options themselves, the network geometry looks like a star. If all of the active treatments have been compared with each other, the network plot can be represented as a complex polygon with all treatment nodes connected to each other [Figure 3].

**Figure 3: Examples of treatment networks.**



*Note: Nodes represent a treatment or an intervention; lines show where direct comparisons exist from 1 or more RCTs:*

- (i) *“Star network”: all interventions have a single mutual comparator.*
- (ii) *“Single closed loop” involves 3 interventions and can provide data to calculate direct comparisons and indirect comparisons (mixed evidence).*
- (iii) *“Connected network”: all interventions have been compared with each other in several trials.*
- (iv) *“Complex network”: combination of star and connected networks*

## 2.3 Statistical Methodology

Network meta-analysis can be performed either with “Frequentist” or “Bayesian” methodologies. Both approaches were developed at the beginning of 20th century. The progress of Bayesian techniques was somewhat delayed because they usually require much more processing power than frequentist ones

**Frequentist** inference has been associated with the frequentist interpretation of probability, specifically that any given experiment can be considered as one of an infinite sequence of possible repetitions of the same experiment, each capable of producing statistically independent results. In this view, the frequentist inference approach to drawing conclusions from data is effectively to require that the correct conclusion should be drawn with a given (high) probability, among this notional set of repetitions. However, exactly the same procedures can be developed under a subtly different formulation. This is one where a pre-experiment point of view is taken. It can be argued that the design of an experiment should include, before undertaking the experiment, decisions about exactly what steps will be taken to reach a conclusion from the data yet to be obtained. These steps can be specified by the scientist so that there is a high probability of reaching a correct decision where, in this case, the probability relates to a yet to occur set of random events and hence does not rely on the frequency interpretation of probability.

**Bayesian** inference has often been thought of as almost equivalent to the Bayesian interpretation of probability and thus that the essential difference between frequentist inference and Bayesian inference is the same as the difference between the two interpretations of what a "probability" means. However, where appropriate, Bayesian inference (meaning in this case an application of Bayes' theorem) is used by those employing a frequentist interpretation of probabilities.

There are two major differences in the frequentist and Bayesian approaches to inference that is not included in the above consideration of the interpretation of probability:

- In a frequentist approach to inference, unknown parameters are often, but not always, treated as having fixed but unknown values that are not capable of being treated as random variants in any sense, and hence there is no way that probabilities can be associated with them. In contrast, a Bayesian approach to inference does allow probabilities to be associated with unknown parameters, where these probabilities can sometimes have a frequency probability interpretation as well as a Bayesian one. The Bayesian approach allows these probabilities to have an interpretation as representing the scientist's belief that given values of the parameter are true.
- While "probabilities" are involved in both approaches to inference, the probabilities are associated with different types of things. The result of a Bayesian approach can be a probability distribution for what is known about the parameters given the results of the experiment or study. The result of a frequentist approach is either a "true or false" conclusion from a significance test or a conclusion in the form that a given sample-derived confidence interval covers the true value: either of these conclusions has a given probability of being correct, where this probability has either a frequency probability interpretation or a pre-experiment interpretation.

## **2.4 Internal and External Validity of NMA**

The external validity of the network meta-analysis will naturally be limited by the external validity of the RCTs included in the evidence network. Quality assessment of RCTs included in the analysis is crucial.

The internal validity is based on the appropriate identification of studies that form the evidence network, the quality of the individual RCTs, and the extent of confounding bias due to similarity and consistency violations.

Another parameter which is critical for the applicability of the outcomes in clinical practice is the selection of an appropriate endpoint in order to perform the comparisons among treatments. [Charter 4]

## **Chapter 3 – Overview of treatment options in Small Cell**

### **Lung Cancer**

#### **3.1 Overview**

The management of SCLC is complicated by the aggressiveness of the disease. Most patients present with symptoms of bulky intra-thoracic disease and/or wide-spread metastases that cause significant debility. Due to the high prevalence of tobacco use, many patients also have substantial co-morbidities that contribute to their impaired performance status and limit the delivery of optimal treatment. These factors also make it challenging to enroll patients with SCLC onto appropriate clinical trials.

Platinum-based chemotherapy is the treatment of choice in patients with both limited (LD-SCLC) and extended (ED-SCLC) disease that has good Performance Status (PS) and organ function. Several prospective trials have compared platinum-based (cisplatin plus etoposide) versus non-platinum-based (anthracycline-based) regimens in both LD and ED patients.

Combination chemotherapy is clearly superior to single-agent treatment in SCLC, and during the 1970s, the cyclophosphamide, anthracycline and vincristine (CAV) regimen became the standard treatment [29].

Only in the early 1980s, the combination of cisplatin and etoposide emerged as first-line treatment in SCLC. Although randomized Phase III studies failed to prove a definitive survival benefit compared with CAV [30-32], an overview of US National Cancer Institute sponsored trial (conducted between 1972 and 1990) confirmed an improvement in median survival for patients treated with cisplatin-based regimens [33]. The EP regimen was better

tolerated in combination with thoracic radiotherapy and soon became the most frequently used chemotherapy regimen for SCLC [34].

### **3.2 LD-SCLC**

LD-SCLC is a potentially “curable” disease in which recent progress has mainly been made through advances in the use of radiotherapy. Two meta-analyses have demonstrated that the addition of definitive thoracic radiation to chemotherapy significantly improves overall survival in patients with LD-SCLC [35, 36]. Further studies have shown that early thoracic radiotherapy resulted in a greater overall survival benefit than late radiotherapy [37].

Although a large, randomized trial reported an added improvement in survival with hyper-fractionated, twice daily, thoracic radio-therapy, this strategy remains controversial and confirmatory studies are on-going [38].

Up to 60 % of patients with SCLC are going to develop brain metastases during the course of their illness. A meta-analysis of randomized trials evaluating prophylactic cranial irradiation (PCI) reported a significant decrease in the incidence of brain metastases and a 5.4 % increase in 3-year overall survival [39].

At present, the standard-of-care for patients with LD-SCLC consists of 4-6 cycles of cisplatin and etoposide plus early, concurrent thoracic radiotherapy. PCI is recommended for those achieving a good response to initial therapy. With such treatment, objective response is noted in 90 % of patients with long-term survival in 25 %

### **3.3 ED-SCLC**

ED-SCLC remains an incurable disease in which the mainstay of treatment is platinum-based, two-drug chemotherapy, such as cisplatin or carboplatin plus etoposide, with the goal of palliating symptoms and prolonging survival. This treatment yields an objective response in 60-70 % of patients with up to 10 % having a complete radiographic response. Patients who attain a good response are considered for PCI based on the demonstration of improved survival even in those with extensive-stage disease [40].

Although chemotherapy does significantly improve quality-of-life and prolong survival for patients with ED-SCLC, relapse is inevitable, and only 5 % of patients remain alive 2 years after the initial diagnosis. Numerous chemotherapy-based strategies, including dose intensification, weekly administration, three- or four-drug regimens, high-dose consolidation, alternating or sequential non-cross-resistant regimens, and maintenance therapy, have failed to improve survival, and several of these approaches have resulted in excessive toxicity [41].

Single-agent chemotherapy is the standard treatment for patients with relapsed SCLC. While response rates are generally higher with combination therapy, over-all survival is not improved, and the toxicity of combination regimens is problematic [42].

The benefits of subsequent therapy are strongly impacted by the duration of response to initial treatment, with lower response rates noted in patients who relapse within 2-3 months of initial therapy. Despite the relatively poor responses and short survival associated with second-line chemotherapy, a randomized trial comparing oral topotecan to best supportive care did demonstrate significantly better overall survival in patients receiving chemotherapy (median, 26 vs. 14 weeks;  $p = 0.01$ ) [43].



It is unlikely that empiric chemotherapy will lead to further significant improvements in outcome in patients with ED-SCLC. The overall survival of patients with ED-SCLC has changed little since the advent of active chemotherapy regimens in the 1970s [44].

Future advances will rely on efforts to better understand the underlying biology of SCLC and to identify molecular targets that drive survival, proliferation and metastasis. In addition, we must improve and broaden our clinical research infrastructure to optimize enrollment onto rational clinical trials.

### **3.4 Current issues in pharmacotherapy**

#### **3.4.1 Cisplatin versus carboplatin**

In clinical practice, EP is the most commonly used initial combination chemotherapy regimen; however, significant symptomatic non-hematological toxicities are related to cisplatin administration. Carboplatin is frequently substituted for cisplatin, but its use carries a greater risk of myelosuppression [45]. Although the mechanisms of action are similar, it is unclear whether carboplatin and cisplatin have the same clinical efficacy.

Four RCTs have suggested similar efficacy of cisplatin and carboplatin in patients with SCLC, indicating carboplatin-etoposide regimen as a valid alternative in elderly or poor-risk patients with ED-SCLC, in consideration of the risk-benefit balance (Table 3) [49-49].

On these studies, a meta-analysis of individual patients data, involving 663 LD (32%) and ED-patients (68%), was conducted: there was no evidence of treatment difference between the cisplatin and carboplatin arms in terms of OS (median OS of 9.6 vs 9.4 months, respectively; HR 1.08,  $p = 0.37$ ), progression-free survival (median PFS of 5.5 vs 5.3 months, respectively; HR 1.10,  $p = 0.25$ ), and ORR (67.1 vs 66.0%, respectively; RR 0.98,  $p = 0.83$ ) [50].

The range of toxicity of the two platinum agents was different. Carboplatin-based regimens were associated with a significantly higher incidence of Grade 3 hematologic toxicities compared with cisplatin-based. On the contrary, non-hematological side effects resulted higher with cisplatin-based treatment. Heterogeneity among studies was found for some adverse effects, probably due to the different drugs and doses used.

Overall, the choice of the platinum compound for first-line treatment of patients with SCLC in clinical practice is based on the expected toxicity profile, age, the patient's organ function and the patient's comorbidities

**Table 3: Phase III trials comparing platinum-based regimens: carboplatin or cisplatin.**

Author and year	Disease Stage	Treatment arms	Number of patients	Performance status	ORR (%)	PFS (months)	Median OS (months)
Skarlos, 1994 [47]	LD/ED	EP	71	0-2	LD:76; ED:60	8.4	12.5
		EC	72		LD:86; ED:67	8.6	11.8
Joss, 1995 [46]	ED	PAV CyMOC	27	0-2	65	NA	8.6
		CV	32		29	NA	4.8
Okamoto, 2007 [48]	ED	EP	110	0-2	73	4.7	9.9
		EC	110		73	5.2	10.6
Lee, 2009 [49]	LD/ED	EP	120	0-2 (ED)	63	6.3	8.1
		GC	121	2 (LD)	63	5.9	8.0

*Note: EC: Etoposide + carboplatin; EP: Etoposide + cisplatin*

### 3.4.2 First-line regimens other than platin plus etoposide

The most recent challenge to EP has come from the combination of **irinotecan plus cisplatin** (IP). In the first Phase III trial of this regimen, Noda et al. from the Japanese Cooperative

Oncology Group (JCOG) randomized 154 patients with previously untreated ED-SCLC to either EP or IP and reported that IP resulted in significantly better response rate, progression-free survival and overall survival (Table 4) [51].

As expected, patients receiving IP had significantly more severe diarrhea, while those receiving EP had greater hematologic toxicity. However, randomized trials in Western patients have failed to confirm the superiority of IP over EP. Hanna et al. in North America and Australia compared a modified IP regimen to EP in 331 patients with previously untreated ED-SCLC and reported no difference in overall efficacy. As in the JCOG trial, there was more myelosuppression and febrile neutropenia in patients receiving EP and more diarrhea in those treated with IP (Table 4) [52].

Lara et al. in the US, randomized 651 patients with previously untreated ED-SCLC to receive IP or EP using the same regimens and schedules as reported by Noda et al. [53]; there was no significant difference in response rate or survival between the two arms (Table 4). A European study by Zatloukal et al. randomized 405 patients with previously untreated ED-SCLC to receive IP or EP and reported non-inferiority of IP, with response rate and TTP non-significantly favoring EP, but overall survival non-significantly favoring IP (Table 4) [54].

Recently, the JCOG investigators, led by Kubota et al. reported a Phase III, randomized trial of IP versus EP consolidation therapy after induction with EP plus thoracic RT in patients with LD-SCLC [55].

In contrast to the prior JCOG study in ED-SCLC [51], the current study failed to demonstrate a significant difference in PFS (1.0 vs 1.1 years,  $p = 0.74$ ) or overall survival (2.8 vs 3.2 years,  $p = 0.70$ ) between IP and EP. The authors do note that the overall survival rates achieved in this trial are the best ever reported in LD-SCLC (5-year, EP 36 vs IP 34%), a finding likely due to improvements in staging and more stringent patient selection in terms of age and performance status.

The more favorable toxicity profile of carboplatin has led to the evaluation of irinotecan plus carboplatin (IC) in patients with SCLC. In a Phase III study, Hermes et al. randomized 209 patients with untreated ED-SCLC to receive IC or oral etoposide plus carboplatin (EC) and reported a significant improvement in overall survival with IC (Table 4) [56].

Quality-of-life measures were similar in both arms. A small, randomized Phase II study of IC versus EC also reported a significantly better response rate and PFS with IC, although overall survival was not reported (Table 4) [57].

Both irinotecan and topotecan are topoisomerase 1 inhibitors derived from camptothecin, although topotecan is generally considered to have a more favorable toxicity profile. Eckardt et al. randomized 784 patients with previously untreated ED-SCLC to receive either oral **topotecan plus cisplatin** (TP) or EP. Once again, efficacy was similar in both arms (Table 4) [58].

As in the prior trials of IP versus EP, there was more neutropenia and febrile neutropenia in patients receiving EP and more diarrhea in those treated with TP. Quality-of-life analysis slightly favored EP ( $p = 0.049$ ) [58]. ED-SCLC is a terminal disease, so many patients and oncologists consider quality of life to be at least as important as duration of survival. A similar study by Fink and collaborators, comparing TP to EP in 703 patients with untreated ED-SCLC, reported non-inferiority of TP, with a significant improvement in response rate and TTP, but only a non-significant trend in overall survival favoring TP (Table 4) [59].

The question of topoisomerase 1 inhibitors versus etoposide has been kept alive by meta-analyses reporting modest improvements in overall survival with platinum plus irinotecan or topotecan combinations [60, 61].

However, the poor long-term survival rates achieved with all these regimens are a clear sign that they are not the final answer for patients with ED-SCLC. These newer combinations (IP, EC and TP) do not appear to be significant steps forward and, for now, EP or EC remain the

standard of care for non-Japanese patients with SCLC. In Japan, IP is commonly used as a first-line regimen for patients with ED-SCLC.

Encouraging data have also been reported from Phase II trials in patients with ED-SCLC with a variety of newer regimens, such as **carboplatin plus paclitaxel** and **paclitaxel plus topotecan** [62, 63].

However, it is highly unlikely that any of these empiric regimens will result in clinically relevant improvements in long-term survival. Numerous chemotherapy-based strategies, including dose-intensification [64], dose-dense regimens [65], weekly administration [66], triplet therapy [67], high-dose consolidation [68], alternating or sequential non-cross-resistant regimens [69], maintenance therapy [70] and consolidation therapy [71], have failed to yield consistent or convincing improvements in survival, and several of these approaches have resulted in unacceptable toxicity [72].

**Table 4: Randomized trials of cisplatin or carboplatin plus irinotecan or topotecan in ED-SCLC**

Author and year	Disease Stage	Treatment arms	Number of patients	ORR (%)	PFS (months)	Median OS (months)
Noda et al. [51]	ED	IP	77	84	4.8	12.8
		EP	77	68	6.9	9.4
Hanna et al. [52]	ED	IP	221	48	4.1	9.3
		EP	110	44	4.6	10.2
Lara et al. [53]	ED	IP	324	60	5.7	9.9
		EP	327	57	5.2	9.1
Zatloukal et al. [54]	ED	IP	202	39	5.4	10.2
		EP	203	47	6.2	9.7
Hermes et al. [56]	ED	IC	105	17		8.5
		EC	104	7		7.1
Schmittl et al. [57]	ED	IC	35	67		
		EC	35	59		
Eckardt et al. [58]	ED	TP	389	63		9.0
		EP	395	69		9.2
Fink et al. [59]	ED	TP	358	56	6.9	11.2
		EP	345	46	6.1	10.2

*Note: EC: Etoposide + carboplatin; EP: Etoposide + cisplatin; IC: Irinotecan + carboplatin; IP: Irinotecan + cisplatin; TP: Topotecan + cisplatin.*

### 3.4.3 New medicinal products

Recent studies have demonstrated that **amrubicin**, a fully synthetic anthracycline that has been approved for clinical use in Japan, has promising activity in patients with SCLC. As first-line therapy for patients with ED-SCLC, Phase II trials have reported response rates of 79% for single-agent amrubicin and 88% for the combination of amrubicin plus cisplatin (AP) [73, 74].

A recently reported Phase III study compared AP to IP in 284 previously untreated patients with ED-SCLC [75]. Disappointingly, the study was stopped early due to futility on interim analysis with similar response rates for AP and IP (78 vs 72%,  $p = 0.33$ ), but with PFS (5.6 vs 5.1 months; hazard ratio 1.42, 95% CI 1.16-1.73) and over-all survival (17.7 vs 15.0 months;

hazard ratio 1.43, 95% CI 1.10 -- 1.85) significantly favoring IP over AP [75]. In patients with recurrent disease, single-agent Phase II studies of amrubicin have yielded response rates of 21-52% [76-78]. Interestingly, response rates and survival were similar in patients with relapsed/sensitive and refractory/resistant disease. However, severe toxicity, mainly hematologic, occurred in > 90% of patients [78, 79].

Three randomized trials have compared amrubicin to topotecan as second-line therapy in patients with SCLC [80-82].

A randomized Phase II trial from Japan compared amrubicin to topotecan in 59 patients with recurrent SCLC and reported significant improvements in response rate (38 vs 13%,  $p = 0.04$ ) and disease control rate (79 vs 46%,  $p = 0.02$ ) with amrubicin [80]. Response rates favoring amrubicin were 53 versus 21% in patients with relapsed/sensitive disease and 17 versus 0% in patients with refractory/resistant disease.

Similarly, a randomized Phase II trial from the US that compared amrubicin to topotecan in 76 patients with relapsed/sensitive SCLC demonstrated a significant improvement in response rate (44 vs 15%,  $p = 0.02$ ) with amrubicin [81].

Jotte et al. recently reported a Phase III trial in which 637 patients with recurrent SCLC were randomized in a 2:1 manner to receive either amrubicin or topotecan [82]. There was a significant improvement in response rate (31 vs 17%,  $p = 0.0002$ ) with amrubicin, but no difference in median PFS (4.1 vs 4.0 months,  $p = 0.98$ ) or overall survival (7.5 vs 7.8 months,  $p = 0.17$ ). Interestingly, in the subgroup of patients with refractory/resistant disease, the 1-year overall survival rate was significantly better with amrubicin (17 vs 8%,  $p = 0.019$ ). While topotecan resulted in more high-grade myelosuppression, amrubicin led to a significantly greater rate of febrile neutropenia and infection [82].

### 3.4.4 Targeted Therapies

A number of molecularly targeted therapies have been evaluated in SCLC either as monotherapy or in combination with other anti-tumor agents. These include clinical trials in the first-line setting, as maintenance therapy and in relapsed SCLC. Sharp et al. recently published a review compiling all investigational therapies for SCLC [83]. These studies are reported at Tables 5 and 6 [84-121].

**Table 5: Targeted therapies in first line treatment of small cell lung cancer**

Putative target	Agent	Author	Phase	Therapy	Outcome
Targeting angiogenesis					
VEGF-A	Bevacizumab	Pujol, 2015 [115]	II/III	Combination	Negative
		Patton, 2006 [89]	II	Monotherapy	15 m OS
		Spigel 2008, [97]	II	Combination	Stopped
		Ready, 2011 [90]	II	Combination	7 m PFS, 11.6 m OS
		Spigel, 2009 [96]	II	Combination	9.1 m TTP, 12.1m OS
		Spigel, 2011 [99]	II	Combination	Negative
RAF-1, VEGFR-2, VEGFR-3 and PDGFRb	Sorafenib	Sharma, 2014 [94]	II	Combination	7.4 m OS
VEGFR-1, VEGFR-2, VEGFR-3, PDGFR, c- KIT, FLT-3 and RET	Sunitinib	Spigel, 2012 [95]	II	Monotherapy	7.6 m PFS
		Ready, 2015 [91]	II	Monotherapy	Improved PFS but not OS
Targeting cell signaling					
BCR-Abl, c-KIT and PDGFR	Imatinib	Johnson, 2003 [85]	II	Monotherapy	0.8 m TTP
		Spigel, 2007 [98]	II	Monotherapy	5.4 m PFS, 8.4 m OS
		Schneider, 2010 [93]	II	Monotherapy	4.3 m PFS, 7.8 m OS
mTOR	Temsirolimus	Pandya, 2007 [88]	II	Monotherapy	2.5 m PFS
IGF1R	Cixutumumab	Belani, 2013 [84]	II	Combination	Negative
Smoothened	Vismodegib				
Targeting apoptosis					
BCL-2	Oblimersen	Rudin, 2008 [116]	II	Combination	Negative
BCL-2, MCL-1, BCL- W, BCL-XL	Obatoclax	Langer, 2014 [117]	II	Combination	Negative



Targeting DNA repair defects					
PARP	Veliparib	Owonikoko 2014 [87]	I	Combination	Negative
	Olaparib	Ongoing	II	Monotherapy	ISRCTN73164486
Targeting the immune system					
CTLA-4	Ipilimumab	Reck, 2013 [118]	II	Combination	Improved iRPFS
		Ongoing	II	Combination	NCT01331525
		Ongoing	II	Combination	NCT02046733
		Ongoing	III	Combination	NCT01450761

*Note: VEGF: Vascular endothelial growth factor; VEGFR: Vascular endothelial growth factor receptor; PDGFR: Platelet derived growth factor receptor; c-KIT: Stem cell factor receptor; FLT3: FMS-like tyrosine kinase 3; RET: Rearranged during transfection tyrosine kinase; mTOR - mammalian target of rapamycin; IGFR: Insulin like growth factor receptor; BCL-2: B-cell lymphoma; MCL-1: myeloid cell leukaemia 1; HDAC: Histone deacetylase; PARP: Poly-ADP ribose polymerase; EGFR: epidermal growth factor receptor; CTLA: Cytotoxic T-lymphocyte-associated protein 4; iRPFS: Immune-related progression-free survival.*

**Table 6: Targeted therapies in treatment of relapsed small cell lung cancer**

Putative target	Agent	Author	Phase	Therapy	Outcome
Targeting angiogenesis					
VEGF-A	Bevacizumab	Jalal, 2010 [106]	II	Combination	14.7 w PFS, 30 w OS
		Waterhous, 2010 [114]	II	Combination	17.4 w PFS, 31.6 w OS
		Mountzios, 2012 [110]	II	Combination	2.7 m PFS, 6.3 m OS
RAF-1, VEGFR-2, VEGFR-3 and PDGFRb	Sorafenib	Gitlitz, 2010 [103]	II	Monotherapy	6.7 m OS, 5.3 m OS
VEGFR-1, VEGFR-2, VEGFR-3, PDGFR, c- KIT, FLT-3 and RET	Sunitinib	Han, 2014, [104]	II	Monotherapy	1.4 m PFS, 5.6 m OS
VEGFR-1 and VEGFR- 2	Aflibercept	Allen, 2014 [100]	II	Combination	Negative
VEGFR-1, VEGFR-2 and VEGFR-3, PDGF and c-Kit	Pazopanib	Gandhi, 2012 [102]	II	Monotherapy	14.1 w PFS
Targeting cell signaling					
EGFR	Gefitinib	Moore, 2006 [109]	II	Monotherapy	50 d TTP
BCR-Abl, c-KIT and	Imatinib	Johnson, 2003 [85]	II	Monotherapy	1.2 m TTP

PDGFR		Krug, 2006 [107]	II	Monotherapy	All PD by 4 w
mTOR	Everolimus	Tarhini, 2010 [112]	II	Monotherapy	1.3 m PFS, 6.7 m OS
BCL-2, BCL-W, BCL-XL	Navitoclax	Rudin, 2012 [111]	II	Monotherapy	1.5 m PFS, 3.2 m OS
BCL-2, BCL-XL, BCL-W, MCL-1	Gossypol	Heist, 2010 [105]	I/II	Combination	17.4 w PFS, 11.7 w
Proteasome	Bortezomib	Lara, 2006 [108]	II	Monotherapy	1 m PFS, 3 m OS
HDAC	Panobinostat	De Marinis, 2013 [101]	II	Monotherapy	Negative
Targeting DNA repair defects					
PARP	BMN673	Wainberg, 2014 [113]	I	Monotherapy	18% RR
Targeting the immune system					
PD-1/CTLA-4	Nivolumab	Antonia, 2016 [119]	I/II	Combination	Nivolumab 18% ORR and 4.4 m OS, Combination 17% ORR and 8.2 m OS
	Ipilimumab				
Antibody-drug conjugates					
CD56	Lorvotuzumab mertansine	Beck, 2012 [120]	I	Monotherapy	25% PR/SD
DLL3	Rovalpituzumab tesirine	Rudin, 2015 [121]	I	Monotherapy	22% ORR


*Note: VEGF: Vascular endothelial growth factor; VEGFR: Vascular endothelial growth factor receptor; PDGFR: Platelet derived growth factor receptor; c-KIT: Stem cell factor receptor; FLT3: FMS-like tyrosine kinase 3; RET: Rearranged during transfection tyrosine kinase; mTOR - mammalian target of rapamycin; BCL-2: B-cell lymphoma; MCL-1: Myeloid cell leukaemia 1; HDAC: Histone deacetylase; CD5: Neural cell adhesion molecule; DLL3: delta-like 3; PARP: Poly-ADP ribose polymerase; CTLA: Cytotoxic T-lymphocyte-associated protein 4; EGFR: epidermal growth factor receptor; PD-1: programmed cell death protein-1.*

The majority of these trials have not found a benefit probably due to both disease and trial-related issues. This may be reflected by the relatively small number of patients enrolled into clinical trials of targeted therapies in SCLC.

## Chapter 4 – Selection of an appropriate endpoint to assess relative effectiveness of treatments in SCLC

In oncology/hematology setting, long term outcomes are considered of high importance for treatment selection, both by clinicians and patients. For current standard of care, long-term efficacy is well characterized, at least compared to new medicinal entities, where limited data on long-term outcome as well as safety (including immunogenicity) are usually available at the time of submission. In order the outcome of a NMA to be applicable in clinical practice, it is of great importance to use the most appropriate endpoint, closest to clinical practice. If it not feasible, it is crucial to identify an alternative sensitive surrogate endpoint (Figure 4)

**Figure 4: Surrogate endpoints for long-term effectiveness in oncology/hematology**



The last few years have seen an increase in the number of randomized controlled trials (RCTs) of new agents in metastatic solid tumors using progression-free survival (PFS) as the primary end point. Disease progression is one of the original four categorical outcomes to

describe change in tumor burden developed first by the WHO [122] and updated recently by the RECIST (Response Evaluation Criteria in Solid Tumors) working group. [123] In general, these criteria were intended for use in clinical trials that used response rate as a primary objective, such as phase II screening trials of new drugs. It is important to note that these measures were intended simply to describe what happens to tumors during therapy—not to infer a meaningful benefit from those changes.

However, there can be practical limitations to using OS as a primary trial end point, including the need for larger sample sizes and longer follow-up. An end point that is a surrogate for OS would be helpful in addressing these limitations but must first be validated by satisfying statistical criteria. [124-128]

Foster et al [129, 130] investigated the putative surrogate endpoints of best response, complete response, confirmed response, and progression-free survival for associations with overall survival and as possible surrogate endpoints for OS, by analyzing individual 2855 patients' data in ten ED-SCLC first-line therapy trials. PFS demonstrated strong surrogacy for OS ( $R = 0.81$ ) in first line ED-SCLC based on this external validation study of individual patient data. PFS is a good alternative end point to OS and should be considered when resource constraints (time or patient) might make it useful or desirable in place of OS

Surrogacy analyses of PFS versus OS have been performed across many disease sites with mixed results. [131] PFS has been shown to be a valid surrogate end point for OS in advanced ovarian and advanced colorectal cancer.[132-135] Other disease sites, including advanced breast cancer,[136-139] advanced prostate cancer,[140, 141] advanced gastric cancer,[142] and advanced NSCLC,[143] have not supported PFS as a surrogate end point for OS.

Tumor response may be may be another surrogate for long term outcomes in ED-SCLC. Based on ORR and PFS reported in the ED-SCLC studies (Table 8) identified for this analysis [Chapter 5], the end points were well correlated.

**Table 7: ED-SCLC studies used to estimate correlation coefficient and equation of the regression line**

Treatment	Study	X	Y	X·Y	X·X	Y·Y
Cisplatin + Etoposide versus Cisplatin + Irinotecan	Noda 2002 [51]	0.38	0.43	0.1634	0.1444	0.1849
	Hanna 2006 [52]	0.83	0.99	0.8217	0.6889	0.9801
	Lara 2009 [53]	0.89	0.74	0.6586	0.7921	0.5476
	Zatloukal 2010 [54]	1.34	0.88	1.1792	1.7956	0.7744
Cisplatin + Etoposide versus Carboplatin + Etoposide	Okamoto 2007 [48]	1.03	0.77	0.7931	1.0609	0.5929
Cisplatin + Etoposide versus Cisplatin + Topotecan	Eckardt 2006 [58]	1.30	1.00	1.3	1.69	1
	Fink 2012 [59]	0.67	0.84	0.5628	0.4489	0.7056
Carboplatin + Etoposide versus Carboplatin + Irinotecan	Schmittel 2011 [144]	0.92	0.74	0.6808	0.8464	0.5476
Carboplatin + Etoposide versus Cisplatin + Irinotecan	Socinski 2009 [145]	2.36	1.37	3.2332	5.5696	1.8769
Cisplatin + Irinotecan versus Cisplatin + Pemetrexed	Socinski 2006 [147]	1.21	1.35	1.6335	1.4641	1.8225

*X: Odd Ration for Objective Response Rate, Y: Hazard Ration for Overall Survival 1Y post treatment*

Specifically a using a weighted regression model with and without taking into account measurement error in the independent variable, revealed a strong uphill linear relationship (r=0.7811)


$$\sum X = 10.93, \sum Y = 9.11, \sum X \cdot Y = 11.0263, \sum X^2 = 14.5009, \sum Y^2 = 9.0325$$

$$r = \frac{n \cdot \sum XY - \sum X \cdot \sum Y}{\sqrt{[n \sum X^2 - (\sum X)^2] \cdot [n \sum Y^2 - (\sum Y)^2]}}$$

$$r = \frac{10 \cdot 11.0263 - 10.93 \cdot 9.11}{\sqrt{[10 \cdot 14.5009 - 10.93^2] \cdot [10 \cdot 9.0325 - 9.11^2]}} \approx 0.7811$$

The estimated regression line of the adjusted model was  $Y = 0.454 + 0.419 x$ , which predicts an approximately 40% increase in log HR of PFS for every unit increase in the log OR of ORR. (Figure 5)

**Figure 5: Equation of the regression line**



$$\sum X = 10.93, \sum Y = 9.11, \sum X \cdot Y = 11.0263, \sum X^2 = 14.5009$$

$$a = \frac{\sum Y \cdot \sum X^2 - \sum X \cdot \sum XY}{n \cdot \sum X^2 - (\sum X)^2} = \frac{9.11 \cdot 14.5009 - 10.93 \cdot 11.0263}{10 \cdot 14.5009 - 10.93^2} \approx 0.454$$

$$b = \frac{n \cdot \sum XY - \sum X \cdot \sum Y}{n \cdot \sum X^2 - (\sum X)^2} = \frac{10 \cdot 11.0263 - 10.93 \cdot 9.11}{10 \cdot 14.5009 - (10.93)^2} \approx 0.419$$

$$y = a + b \cdot x$$

$$y = 0.454 + 0.419 \cdot x$$

# **Chapter 5 – Assessing the relative effectiveness of treatments in SCLC - Frequentist Approach**

*Bakalos G et al. Assessing the relative effectiveness and tolerability of treatments in small cell lung cancer: a network meta-analysis. Cancer Epidemiol. 2013 Oct;37(5):675-82*

## **5.1 Introduction**

In order to evaluate the relative merits of the different treatments for SCLC based on the mode of action of each chemotherapy agent (or combination of individual chemotherapy agents), we systematically searched and catalogued all available published RCTs in SCLC. Then, we performed a network of multiple treatments analysis (network meta-analysis), involving direct analysis (synthesis of RCTs with the same treatment comparisons) and indirect analysis (comparison between treatments using an intermediate comparator). In the absence of direct comparison between treatments, the effect size can only be estimated only using an indirect comparison approach.

## **5.2 Materials & Methods**

### **5.2.1 Search strategy-Selection of RCTs**


We searched PubMed, EMBASE, and the Central Registry of Controlled Trials of the Cochrane Library to identify all RCTs that investigated chemotherapy regimens in adult patients with histologically proven SCLC. The search was limited to English language, RCTs, adults, and concerned the time period from 1980 until May 1<sup>st</sup>, 2011. The articles were



identified using as search criterion the terms: “small cell lung cancer” and “chemotherapy”.

The reference lists of the retrieved articles were also reviewed to identify additional publications. The search strategy for the selection of the eligible RCTs is shown in Figure 6.

**Figure 6: Flow diagram of the screening process and RCTs selection for multiple-treatments meta-analysis of treatments for SCLC**



### 5.2.2 Eligibility criteria

RCTs that compared at least two arms of different chemotherapy regimens in chemotherapy naïve patients with histologically proven SCLC were included in the network analysis. Only studies that provided sufficient data to calculate odds ratios (ORs) for estimating the magnitude of difference between treatments, and the corresponding precision were considered.

The following studies were excluded:

- Studies comparing second line chemotherapy treatments,
- Studies reporting radiotherapy interventions, i.e. radical radiotherapy in combination with chemotherapy or chemotherapy administration for sensitization to radiation,
- Studies reporting surgical interventions,
- Studies reporting adjuvant chemotherapy (i.e. chemotherapy following radical surgical intervention) or neo-adjuvant chemotherapy (i.e. chemotherapy prior to radical surgical interventions),
- Studies reporting supportive care interventions or comparison of chemotherapy with chemotherapy plus conventional supportive care and
- Follow-up and extension studies. In addition, studies with a crossover design, meeting abstracts and conference proceedings were excluded.

In RCTs involving more than two treatment arms, each pair-wise treatment comparison was considered as different study. Also, RCTs providing data for different SCLC stages were considered as separate studies in the analysis. In order to avoid the inclusion of duplicated data, the retrieved studies were appraised by geographic location, author names and period of study. Then, in studies with overlapping patients, the largest one was included in the analysis. Only studies conducted after approval from national ethical committees were considered.

### **5.2.3 Data extraction and outcomes definition**

The following information was extracted from each eligible article: name of first author, year of publication, country of origin, reported stage of SCLC, sample size (randomized patients, totally and per arm), types and intensity (dose and duration) of chemotherapies, effect size of each outcome of interest and chemotherapy regimen. Data extraction was undertaken by 2 investigators, independently. The overall agreement rate was 89%. Any disagreement was resolved by a third independent investigator.

Two primary outcomes were considered to assess relative effectiveness (CR and ORR).

Complete Response (CR) is achieved when all tumor lesions are disappeared after treatment initiation. Objective Response Rate (ORR) is the portion of patients with a predefined amount of tumor size reduction; ORR is defined as the sum of CR and partial response and it is a direct measure of drug antitumor activity. However, this exercise was based only on ORR (CR based analyses are presented as supplementary materials)

Among the many adverse events after treatment with chemotherapy, we chose to record the neutropenia (NP) and febrile neutropenia (FNP) because they are considered the most important ones.

### **5.2.4 Treatment definition**

Chemotherapy regimens containing the same chemotherapy agents, irrespective of dosage scheme and maximum duration of each chemotherapy cycle, were defined as the same treatment since we are interested in the assessment of the relative effectiveness of the different agent-based therapies. In addition, the effect of different dosage schemes and chemotherapy cycle intensity remains unresolved [147].

Furthermore, the current grouping allows the definition of a less complicated and analysable network. The combination of cisplatin and etoposide (EP) was set as the reference treatment in the subsequent treatment comparisons since it is the standard first line treatment and the most commonly investigated chemotherapy regimen.

### **5.2.5 Statistical methods – Frequentist Analysis**

Treatments were compared using odds ratios (ORs) with their respective 95% confidence intervals (CI). When more than two studies compared the same treatments, random effects (RE) pooled OR was calculated [148]. The RE model incorporates the between study variability and it is more conservative than the fixed effects model [149].

Indirect comparison was performed for treatments not compared directly [10]. Then, in comparing two treatments, A and B, where each treatment was compared directly with treatment C, the OR for comparing A and B was calculated using the following principle:  $\ln(\text{OR}_{A\text{vs}B}) = \ln(\text{OR}_{A\text{vs}C}) - \ln(\text{OR}_{B\text{vs}C})$ , and the respective 95% (CI) was estimated assuming asymptotic normality and lack of covariance [3, 10, 13, 150, 151] (Figure 1).

The network of treatments was constructed based on all investigated comparisons between treatments and the indirect analysis was performed utilizing all the possible pathways provided by the network. The OR was considered significant when the 95% CI included the one ("1").

The network graph was built using yFiles.NET (yWorks GmbH, Tübingen, Germany, <http://www.yWorks.com>) [152] and the network analysis was carried out using NET-MS (<http://netms.med.uth.gr>). MetaAnalyst (Evidence-Based Practice Center, Tufts Medical Center, Boston, MA, USA, [http://tuftscaes.org/meta\\_analyst](http://tuftscaes.org/meta_analyst)) [153] was used to validate the findings of data syntheses.

## 5.3 Results

### 5.3.1 Eligible studies and summary characteristics

The literature search in PubMed, EMBASE and Cochrane Central Register of Controlled Trials identified 243 articles that met the search criterion. After title selection and abstract reviewing of the articles, 98 articles from all databases were judged to be potentially relevant and they were reviewed in their entirety. Finally, 71 articles were selected for inclusion the network analysis: 71 articles for the outcome CR, 69 for ORR, 23 for NP and 5 for FNP. The articles were published between 1980 and 2011. The summary characteristics of included RCTs in the multiple treatments meta-analysis is shown in Table 9.

**Table 8: Characteristics of 71 included randomized controlled trials (RCTs) in the network of treatments meta-analysis**

Characteristic	Number
Number of Interventions	91
Number of Studies	71
Total Number of Patients in Network	16026
Number of Two-arm Studies	63
Number of Multi-Arms Studies	8

The characteristics of the individual RCTs including efficacy/tolerability results and their quality assessment are shown in **Supplementary Tables 2-4** and the definition of the treatments is given in **Supplementary Table 5**.

In total, the eligible studies involved 16,026 randomly assigned patients with SCLC and the majority of them were male (72%) while the median age was 61 (55-74) years. Most of the articles involved studies carried out in the US (21.1%) and 18.3% of them were multicenter

trials, involving several countries. 83.7% (63/71) of the studies included two comparing treatment arms and only 11.3% of them compared more than two arms.

More than half of the studies (37/71) included patients with extended disease, while two studies included patients with limited disease and 32 patients with ED and LD. The median sample size was 230 (12-455) patients.

Overall survival was reported in 67 articles. Overall, the median ORR was 65.2% (10.0%-96.9%) while the median overall survival was 10.3 (1.0-27.7) months. Adverse events of grade 3-4 were reported for 58.5% (9,371) of the patients. Almost half patients experienced grade 3-4 neutropenia (53.1%) and leucopenia (44.3%). Thrombocytopenia and anemia was reported in 22.0% and 19.8% of the patients, respectively.

In assessing the quality of reporting, seven items were considered:

- Precise details of the interventions in each arm,
- Description of study end-points,
- Description of sample size estimation,
- Method of randomization (sequence generation),
- Implementation of randomization,
- Blinding, and
- Participant flow.


The majority of studies were open-label and only two studies were blinded. The precise details of the interventions in each arm were reported in all studies, while the study end-points and the sample size estimation were reported in 50 (70.4%) and 54 (76%) studies respectively. Despite the fact that the method of randomization was described in 32 (45%)

reports, only two reports provided information about the implementation of randomization (2.8%). The participant flowchart was described in 66 studies (93%).

### **5.3.2 The networks**

The geometry of the network of comparisons for ORR is depicted in Figure 7; the other outcomes are represented in **Supplementary Figures 1 - 3**.

**Figure 7: Network of direct comparisons for the outcome “Objective Response Rate”.**



In the network figure, the size of the circles was directly related to the number of RCTs investigated each treatment, while the thickness of connecting lines was directly related to the number of available direct comparisons. More specifically, common treatments [e.g. EP] that were compared by more RCTs were drawn with larger circles whereas infrequently investigated regimens (eg. Cisplatin plus Doxorubicin) were represented by smaller circles. However, most of the treatments were compared against EP, which represented the most



commonly used treatment in the RCTs (26 direct comparisons). Carboplatin plus Etoposide involved the biggest sample size of randomized patients (455 patients). All regimens are listed in Supplementary Table 5.

### **5.3.3 Direct analysis for comparing treatments with EP**

Sixteen treatments were compared directly with EP in 18 trials [32, 48, 49, 51, 54, 58, 67, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, and 164]:

- Cyclophosphamide plus Doxorubicin (CAV),
- Cisplatin plus Etoposide plus Ifosfamide (VIP),
- Cyclophosphamide plus Doxorubicin plus Etoposide plus GCSF [ACE (intensified)],
- Cisplatin plus Epirubicin (PEP),
- Cisplatin plus Topotecan (TC),
- Cisplatin plus Etoposide/Cyclophosphamide plus Doxorubicin plus Vincristine (CAV/EP),
- Cisplatin plus Etoposide plus GCSF [EP (intensified)],
- Carboplatin plus Etoposide (EC),
- Carboplatin plus Gemcitabin (GEMCAR),
- Cisplatin plus Irinotecan (IP),
- Cisplatin plus Cyclophosphamide plus Etoposide plus Epirubicin (CCEE),
- Cisplatin plus Etoposide plus Megestrol acetate (EP+Ma),
- Cisplatin plus Etoposide plus interferon alpha (EP + IFNA-a),
- Etoposide plus Ifosfamide (IE) and
- Cisplatin plus Etoposide plus Paclitaxel (PET).

The numbers of direct comparisons with EP for the outcomes CR, ORR, NP and FNP were 18, 17, 9 and 1, respectively. None of the treatments showed better response compared to EP

for both efficacy outcomes. The significant results derived from the direct analysis are shown in Table 9; the treatments are sorted according to their effect sizes. The results of all direct comparisons are shown in Supplementary Table 6.

**Table 9: Direct analysis results for comparing treatments for small cell lung cancer (SCLC) with reference treatment (Cisplatin + Etoposide, EP) by outcome. The treatments were sorted according to their significance and magnitude of effect size**

Treatment	Stage	Patients, No	OR (95% CI)	P-value
<b>Patients with objective response</b>				
Cisplatin + Cyclophosphamide + Etoposide + Epirubicin	ED	226	2.07 (1.17- 3.67)	0.01
Cisplatin + Etoposide + Megestrol acetate	LD, ED	243	0.55 (0.31-0.99)	0.05
<b>Patients with neutropenia</b>				
Cyclophosphamide + Doxorubicin + Etoposide + GCSF	ED	280	7.37 (3.72-14.56)	<0.01
Cisplatin + Cyclophosphamide + Etoposide + Epirubicin	ED	226	5.59 (1.83-17.1)	<0.01
Cisplatin + Epirubicin	LD, ED	402	0.54 (0.37-0.81)	<0.01
Cisplatin + Topotecan	LD, ED	784	0.27 (0.2-0.39)	<0.01

**Patients with complete response:** No regimen was significantly different than EP ( $P \geq 0.05$ ).

**Patients with objective response:** The treatment combination Cisplatin plus

Cyclophosphamide plus Etoposide plus Epirubicin (CCEE) produced better response

[OR=2.07 (1.17-3.67)], whereas Cisplatin plus Etoposide plus Megestrol acetate (EP+Ma)

derived worse response [OR=0.55 (0.31-0.99)].

**Tolerability:** Two treatment combinations produced worse tolerability (in terms of NP) than

EP (Cyclophosphamide plus Doxorubicin plus Etoposide plus GCSF and Cisplatin plus

Cyclophosphamide plus Etoposide plus Epirubicin) and two treatments shown better

tolerability ( $P < 0.01$ ) (Table 9).

Regarding FNP, only one treatment (Cisplatin plus Etoposide plus Ifosfamide) was compared directly to EP, producing non-significant result [OR=1.81 (0.97, 3.40)] [158].

### Indirect analysis for comparing treatments with EP

Table 10 shows the indirect analysis significant results ( $P < 0.05$ ); the treatments are sorted according to their effect size. None of the treatments derived a better response than EP for both efficacy outcomes.

**Table 10: Indirect analysis results for comparing treatments for small cell lung cancer (SCLC) with reference treatment (Cisplatin + Etoposide) by outcome, for treatments that produced significantly ( $P < 0.05$ ) different response than reference treatment**

Treatment	OR (95% CI)	p value
<b>Patients with complete response</b>		
Etoposide	0.36 (0.14-0.88)	0.03
Carboplatin + Ifosfamide	0.31 (0.11-0.88)	0.03
Etoposide (intensified)	0.13 (0.02-0.65)	0.01
<b>Patients with objective response</b>		
Cisplatin + Doxorubicin + Etoposide + Vincristine (intensified)	3.79 (1.77-8.12)	<0.01
Ifosfamide + Mesna	0.43 (0.26-0.70)	<0.01
Carboplatin + Pemetrexeb	0.41 (0.21-0.79)	<0.01
Etoposide	0.40 (0.24-0.68)	<0.01
Doxorubicin + Etoposide + Vincristine	0.40 (0.17-0.94)	0.04
Cyclophosphamide + Doxorubicin + Etoposide	0.38 (0.16-0.93)	0.03
Teniposide	0.35 (0.21-0.57)	<0.01
Cisplatin	0.33 (0.18-0.61)	<0.01
Carboplatin + Ifosfamide	0.25 (0.066-0.94)	0.04
Etoposide (intensified)	0.006 (0.00-0.46)	0.02
<b>Patients with neutropenia</b>		
Carboplatin + Pemetrexeb	0.26 (0.09-0.76)	0.01

However, one treatment [Cisplatin plus Doxorubicin plus Etoposide plus Vincristine (intensified)] showed better response for the outcome ORR but, this treatment showed worse

tolerability in the direct analysis. The results of all indirect comparisons are shown in Supplementary Table 6.

**Patients with complete response:** None treatment showed better outcome than EP ( $P \geq 0.05$ ). However, the analysis indicated that monotherapy with etoposide (either standard or intensified) and combination therapy with Carboplatin plus Ifosfamide have less comparative effectiveness [OR= 0.36 (0.14-0.88), OR=0.13 (0.02-0.65) and OR=0.31 (0.11-0.88), respectively].

**Patients with objective response:** Only one treatment combination yielded better response to Cisplatin plus Etoposide: Cisplatin plus Doxorubicin plus Etoposide plus Vincristine (intensified) [OR=3.79 (1.77-8.12)]. However, nine treatments (table 10) revealed worse response:


- Ifosfamide plus Mesna,
- Carboplatin plus Pemetrexeb,
- Etoposide,
- Doxorubicin plus Etoposide plus Vincristine,
- Cyclophosphamide plus Doxorubicin plus Etoposide,
- Teniposide,
- Cisplatin,
- Carboplatin plus Ifosfamide and viii) Etoposide (intensified)

**Tolerability:** Only the treatment Carboplatin plus Pemetrexeb indicated a better tolerability for the outcome NP [OR=0.26 (0.09-0.76)]. For the outcome FNP, there is only one study [154] reporting this outcome (see direct analysis section) and thus, the comparative tolerability of treatments was not evaluated further.

5.3.4 Sub-analysis of most common interventions in SCLC

A sub-analysis performed including only the most commonly used interventions in ED-SCLC. Specifically we have isolated studies reflecting to current standard of care (48, 51, 54, 56, 59, and 157) reporting outcomes from various comparisons of the following interventions: Cisplatin+Etoposide, Carboplatin+Etoposide, Cisplatin+Irinotecan, and Cisplatin+Topotecan. The sub-network is reflected at figure 8

Figure 8: Network Diagram of most common interventions



Six studies included, where 1151 out of 1864 achieved objective response rate following chemotherapy treatment. The network characteristics, the intervention characteristics as well as the direct comparison characteristics are summarized at tables 11, 12 and 13, respectively.

**Table 11: Network Characteristics**

Characteristic	Number
Number of Interventions	6
Number of Studies	6
Total Number of Patients in Network	1'864
Total Number of Events in Network	1'151
Total Possible Pairwise Comparisons	15
Total Number Pairwise Comparisons With Direct Data	5
Number of Two-arm Studies	6
Number of Multi-Arms Studies	0
Number of Studies With No Zero Events	6
Number of Studies With At Least One Zero Event	0
Number of Studies with All Zero Events	1

**Table 12: Intervention Characteristics**

Treatment	# Studies	# Events	# Patients	Aggregate Rate	Min. Rate	Max. Rate
Cisplatin+Etoposide	5	562	900	0.6244	0.4631	0.7273
Cisplatin+Topotecan	1	245	389	0.6298	0.6298	0.6298
Cisplatin+Etoposide+GCSF	1	65	109	0.5963	0.5963	0.5963
Cisplatin+Irinotecan	2	144	279	0.5161	0.3911	0.8442
Carboplatin+Etoposide	2	103	147	0.7007	0.6216	0.7273
Carboplatin+Etoposide+GCSF	1	32	40	0.8000	0.8000	0.8000

**Table 13: Direct Comparison Characteristics**

<b>Comparison</b>	<b># Studies</b>	<b># Patients</b>	<b># Events</b>
<b>Cisplatin+Etoposide vs. Cisplatin+Topotecan</b>	1	784	517
<b>Cisplatin+Etoposide vs. Cisplatin+Etoposide+GCSF</b>	1	224	129
<b>Cisplatin+Etoposide vs. Cisplatin+Irinotecan</b>	2	559	290
<b>Cisplatin+Etoposide vs. Carboplatin+Etoposide</b>	1	220	160
<b>Carboplatin+Etoposide vs. Carboplatin+Etoposide+GCSF</b>	1	77	55

As it was expected from previous analyses, none of the treatments was significantly different compared to cisplatin plus etoposide, in terms of achieving ORR.

# **Chapter 6 – Assessing the relative effectiveness of treatments in SCLC - Bayesian Approach**

## **6.1 Introduction**

Additional analyses were performed by applying this time Bayesian statistics. For the purposes of this analysis we used WinBUGS (the MS Windows operating system version of BUGS: Bayesian Analysis Using Gibbs Sampling), which is a versatile package that has been designed to carry out Markov chain Monte Carlo (MCMC) computations for a wide variety of Bayesian models [165]. The software is currently distributed electronically from the BUGS Project website (<http://www.mrc-bsu.cam.ac.uk/bugs/overview/contents.shtml>).

We used a freely available Microsoft-Excel-based tool called NetMetaXL, programmed in Visual Basic for Applications, which provides an interface for conducting a Bayesian network meta-analysis using WinBUGS from within Microsoft Excel [166]. This tool allows the user to easily prepare and enter data, set model assumptions, and run the network meta-analysis, with results being automatically displayed in an Excel spreadsheet. It also contains macros that use NetMetaXL's interface to generate evidence network diagrams, forest plots, league tables of pairwise comparisons, probability plots (rankograms), and inconsistency plots within Microsoft Excel. This tool was developed to simplify running and reporting network meta-analyses and to highlight how NetMetaXL can be used to facilitate consistent reporting and more efficient and transparent critical appraisal of network meta-analyses submitted to HTA organizations such as the Canadian Agency for Drugs and Technologies in Health (CADTH) and the National Institute for Health and Care Excellence (NICE), as well as to journals which publish network meta-analyses.




## **6.2 Results**

### **6.2.1 Sub analysis of most common interventions in SCLC**

As it was presented at 5.3.4, a sub-analysis performed on a subnetwork of the most common interventions in SCLC. The network diagram (figure 8), network characteristics (table 11), intervention characteristics (table 12), and direct comparison characteristics (table 13) were presented in Chapter 5.

New analyses with Bayesian statistics reveals that carboplatin+etoposide combined with GCSF had the bigger treatment effect size (Figure 9)

**Figure 9: Forrest Plot of most common interventions in SCLC**



The summary league table is shown in tables 14 (Fixed Effects) and 15 (Random Effects). The league table arranges the presentation of summary estimates by ranking the treatments in order of most pronounced impact on the outcome under consideration, based on SUCRA [165]. SUCRA (table 16), the surface under the cumulative ranking, is a simple numerical summary of the probabilities. It is 100% when a treatment is certain to be the best and 0% when a treatment is certain to be the worst. SUCRA values enable the ranking of treatments overall for a particular outcome. In this particular analysis Carboplatin+Etoposide +GCSF is listed in the top left of the diagonal of the league table because it was associated with the most favorable SUCRA for ORR, while Cisplatin+Topotecan is listed in the bottom right of the diagonal of the league table because it was associated with the least favorable results. For interpretation purposes, the results are read from top to bottom and left to right.

Probability bars (or rankograms) were developed within NetMetaXL, to visualize the probability that each treatment is ranked first, second, and so on for a particular outcome. These rankograms are depicted as stacked vertical bar charts for all treatments (figure 10).


**Table 14: Rankogram of most common interventions in SCLC, FE**

<b>Carboplatin+Etoposide+GCSF</b>					
2.10 (0.57 – 8.09)	<b>Cisplatin+Etoposide+GCSF</b>				
2.50 (0.77 – 8.59)	1.18 (0.69 – 2.02)	<b>Cisplatin+Etoposide</b>			
2.49 (0.90 – 7.41)	1.18 (0.53 – 2.63)	1.00 (0.55 – 1.81)	<b>Carboplatin+Etoposide</b>		
2.56 (0.76 – 9.22)	1.21 (0.64 – 2.29)	1.03 (0.72 – 1.45)	1.03 (0.51 – 2.06)	<b>Cisplatin+Irinotecan</b>	
3.25 (0.96 – 11.58)	1.54 (0.84 – 2.84)	1.30 (0.97 – 1.76)	1.30 (0.68 – 2.53)	1.27 (0.80 – 2.00)	<b>Cisplatin+Topotecan</b>

**Table 15: Rankogram of most common interventions in SCLC, RE**

Carboplatin+Etoposide+GCSF					
1.90 (0.03 – 129.70)	Cisplatin+Irinotecan				
2.12 (0.02 – 220.10)	1.10 (0.05 – 28.82)	Cisplatin+Etoposide+GCSF			
2.51 (0.06 – 118.10)	1.31 (0.22 – 8.80)	1.17 (0.09 – 15.56)	Cisplatin+Etoposide		
2.51 (0.17 – 40.10)	1.31 (0.06 – 34.32)	1.16 (0.03 – 45.48)	1.00 (0.07 – 13.15)	Carboplatin+Etoposide	
3.25 (0.04 – 338.00)	1.69 (0.08 – 43.41)	1.52 (0.04 – 58.74)	1.29 (0.10 – 16.68)	1.30 (0.04 – 50.09)	Cisplatin+Topotecan

**Figure 10: Stacked Bar Chart of most common interventions in SCLC**



**Table 16: SUCRA of most common interventions in SCLC**


Treatment	SUCRA
Carboplatin+Etoposide+GCSF	0.8911
Cisplatin+Etoposide+GCSF	0.5847
Cisplatin+Irinotecan	0.5114
Cisplatin+Etoposide	0.4331
Carboplatin+Etoposide	0.4146
Cisplatin+Topotecan	0.1651

Assessment of inconsistency is crucial in the conduct of any network meta-analysis.

Inconsistency can be thought of as a conflict between ‘direct’ and ‘indirect’ evidence [27].

Similar to heterogeneity, inconsistency is caused by imbalances in effect modifiers from study to study, specifically by an imbalance in the distribution of effect modifiers in the direct and indirect evidence [27]. NetMetaXL allows users to assess inconsistency by comparing the deviance residuals and DIC statistics in fitted consistency and inconsistency models. The methods employed are described at NICE Technical Support Documents (TSD) series.

Inconsistency for this analysis was very limited, both in fixed and random effects (Figure 11).

**Figure 11: Inconsistency results of most common interventions in SCLC**

As a conclusion from this analysis, we confirmed that cisplatin+ etoposide is equal to carboplatin+etoposide. However, if we co-administrate GCSF with carboplatin+etoposide in order to minimize the well known risk of myelosuppression associated to carboplatin [45], it is revealed to be the best treatment option.

### 6.2.2 Platin-pooled data sub-analysis in ED-SCLC

Additional analysis performed only for patients with ED SCLC. All studies reporting interventions either in patients with LD-SCLC or they reported mixed LD/ED SCLC results, were excluded (Abratt RP, 1991, Altinbas M, 2004, Ardizzoni A, 2002, Artal-Cortes A, 2004, Baka S, 2008, Baka S, 2010, Bork E, 1991, Chahinian AP, 1989, Ettinger DS, 1990, Ettinger DS, 2002, Fukuoka M, 1997, Girling DJ, 1996, Grote T, 2005, Heigener DF, 2009, Hirsch FR, 1987, Hirsch FR, 2001, James LE, 1996, Joss RA, 1995, Joss RA, 1995(2), Lassen U, 1996, Lee SM, 2009, Leyvraz S, 2008, Lorigan P. 2005, Miller AA, 1995, Milroy R, 1993, Miyomoto H, 1992, Murray N, 1999, Nagel S, 2011, Postmus PE, 1992, Postmus PE, 1996, Rowland KM, 1996, Schmittel A, 2006, Sculier JP, 1990, Sculier JP, 1993, Sculier JP, 2001, Seifart U, 2005, Seifart U, 2007, Sekine I, 2003, Slevin ML, 1989, Souhami RL, 1994, Souhami RL, 1997, Steward WP, 1998, Urban T, 1999, White SC, 2001, Woll PJ, 2001).


As a result several studies were disconnected from the network and were excluded as well (De Marinis F, 2005, Kanitz E, 1992, Monnet I, 1992, Reck M, 2003).

Additional literature analysis was performed up to June 30<sup>th</sup>, 2016 and 14 new trials were identified (Supplementary table 8)

Based on previously reported outcomes [Charter 5], cisplatin and carboplatin arms were pooled together.

The network diagram is presented at figure 12.

**Figure 12: Network diagram of platin-pooled data sub-analysis in ED-SCLC**



Legend	
Drug	Abbreviation
Platin+Etoposide	A
Platin + Irinotecan	B
Platin + Paclitaxel	C
Platin + Topotecan	D
Platin + Amrubicin	E

The network includes 12 studies exploring 5 interventions. 3'961 pts were enrolled; of those, 2'315 achieved objective response rate. The network characteristics, intervention characteristics and direct comparison characteristics are presented at tables 17, 18 and 19, respectively.

**Table 17: Network characteristics of platin-pooled data sub-analysis in ED-SCLC**

Characteristic	Number
Number of Interventions	5
Number of Studies	12
Total Number of Patients in Network	3'961
Total Number of Events in Network	2'315
Total Possible Pairwise Comparisons	10
Total Number Pairwise Comparisons With Direct Data	5
Number of Two-arm Studies	12
Number of Multi-Arms Studies	0
Number of Studies With No Zero Events	12
Number of Studies With At Least One Zero Event	0
Number of Studies with All Zero Events	0

**Table 18: Intervention characteristics of platin-pooled data sub-analysis in ED-SCLC**

Treatment	# Studies	# Events	# Patients	Aggregate Rate	Min. Rate	Max. Rate
Platin+Etoposide	10	991	1764	0.5618	0.4340	0.7000
Platin + Irinotecan	7	626	1'097	0.5706	0.3911	0.8442
Platin + Paclitaxel	1	23	34	0.6765	0.6765	0.6765
Platin + Topotecan	3	442	747	0.5917	0.4167	0.6298
Platin + Ambrubicin	3	233	319	0.7304	0.6667	0.7786


**Table 19: Direct Comparison Characteristics of platin-pooled data sub-analysis in ED-SCLC**

Comparison	# Studies	# Patients	# Events
Platin+Etoposide vs. Platin + Irinotecan	6	1'810	984
Platin+Etoposide vs. Platin + Topotecan	2	1'464	861
Platin + Paclitaxel vs. Platin + Topotecan	1	46	28
Platin+Etoposide vs. Platin + Ambrubicin	2	359	231
Platin + Irinotecan vs. Platin + Ambrubicin	1	282	211



The combinations of platin with paclitaxel, platin with irinotecan and platin with topotecan were not significantly superior to platin plus etoposide. Only platin plus ambrubicin was marginally significantly better. (Figure 13)

**Figure 13: Forest plot of platin-pooled data sub-analysis in ED-SCLC**



SUCRA values supporting that combination of platin plus paclitaxel provides the best probability for ORR in ED-SCLC (SUCRA 91%), followed by platin plus ambrubicin (73%), while the combination of platin plus etoposide provides the lowest probability of success (SUCRA 15%) (Tables 20-22, Figure 14)


**Table 20: League table of platin-pooled data sub-analysis in ED-SCLC - FE**

<b>Platin + Paclitaxel</b>				
2.12 (0.51 – 9.51)	<b>Platin + Ambrubicin</b>			
2.97 (0.74 – 12.69)	1.40 (0.98 – 2.01)	<b>Platin + Irinotecan</b>		
3.05 (0.78 – 12.66)	1.44 (0.96 – 2.16)	1.03 (0.78 – 1.36)	<b>Platin + Topotecan</b>	
3.25 (0.81 – 13.77)	1.53 (1.08 – 2.17)	1.09 (0.91 – 1.31)	1.07 (0.86 – 1.32)	<b>Platin+Etoposide</b>

**Table 21: League table of platin-pooled data sub-analysis in ED-SCLC - RE**

<b>Platin + Paclitaxel</b>				
2.26 (0.36 – 14.09)	<b>Platin + Ambrubicin</b>			
2.88 (0.49 – 16.19)	1.30 (0.65 – 2.39)	<b>Platin + Irinotecan</b>		
3.11 (0.61 – 15.62)	1.40 (0.57 – 3.27)	1.07 (0.53 – 2.32)	<b>Platin + Topotecan</b>	
3.34 (0.59 – 18.27)	1.49 (0.78 – 2.72)	1.15 (0.80 – 1.74)	1.07 (0.57 – 2.00)	<b>Platin+Etoposide</b>

**Figure 14: Rankogram of platin-pooled data sub-analysis in ED-SCLC**




**Table 22: SUCRA of platin-pooled data sub-analysis in ED-SCLC**

Treatment	SUCRA
Platin + Paclitaxel	0.9093
Platin + Ambrubicin	0.7277
Platin + Irinotecan	0.3986
Platin + Topotecan	0.3093
Platin+Etoposide	0.1552

Inconsistency for this analysis was very limited, both in fixed and random effects (Figure 15).

**Figure 15: Inconsistency results of platin-pooled data sub-analysis in ED-SCLC**



### 6.2.3 Sub-analysis – New treatments

An additional analysis was performed in order to estimate the impact of investigational treatments. 11 interventions, including current standard of care were compared in 17 studies which enrolled 4'605 patients. The network characteristics and are presented at table 23.

**Table 23: Network characteristics of platin-pooled data sub-analysis compared to investigational treatments in ED-SCLC**


Characteristic	Number
Number of Interventions	11
Number of Studies	17
Total Number of Patients in Network	4'605
Total Number of Events in Network	2'649
Total Possible Pairwise Comparisons	55
Total Number Pairwise Comparisons With Direct Data	12
Number of Two-arm Studies	16
Number of Multi-Arms Studies	1
Number of Studies With No Zero Events	17
Number of Studies With At Least One Zero Event	0
Number of Studies with All Zero Events	0

Based on the SUCRA analyses (table 24, figure 16), new treatments are anticipated to improve ORR in ED-SCLC. However, are not significantly differentiated from current standard of care and further investigation is needed to drive conclusions on how these treatments would be better utilized in clinical practice.

**Table 24: SUCRA of platin-pooled data sub-analysis compared to investigational treatments in ED-SCLC**

Treatment	SUCRA
Platin + Paclitaxel + Ipilimumab	0.945
Platin+Etoposide	0.7389
Platin + Topotecan	0.6446
Platin + Etoposide + Cixutumumab	0.6014
Platin + Irinotecan	0.5939
Platin + Etoposide + Vismodegib	0.4404
Platin + Etoposide + Bevacizumab	0.3897
Platin + Etoposide + aBcl2	0.3845
Platin + Ambrubicin	0.3368
Platin + Belotecan	0.2993
Platin + Paclitaxel	0.1253

**Figure 16: Rankogram of platin-pooled data sub-analysis compared to investigational treatments in ED-SCLC**



## Charter 7 – Discussion

Herein, we have presented a comprehensive and systematic assessment of the current status of treating SCLC.

The primary aim of the present study was to explore methodological aspects related to network meta-analysis, especially in oncology field. For the purposes of this analysis, SCLC was used as an example provide an assessment of the relative effectiveness of treatments in SCLC, especially in the absence of head-to-head comparisons, and to direct future research in SCLC treatment.

We carried out a network analysis of all published RCTs in SCLC, using both “Frequentist” and “Bayesian” methodologies.

The **frequentist network analysis** involved the following steps: direct comparison of treatments, indirect comparison and combination of direct and indirect comparison. The secondary aim was to reveal the necessity to performing large RCTs for head-to-head comparisons of treatments. There are no studies involved more than 500 patients and the various chemotherapy combinations have not compared to a standard treatment such as EP. The network consisted of 91 treatments, involving 18 direct comparisons for the outcome CR, 17 for the outcome ORR and 10 for the tolerability.

The analysis of the network indicated that only two regimens might have comparable effectiveness to EP: The application of network analysis of treatments makes optimal use of all available published data and provides insight in the relative effectiveness of different treatments (monotherapies and combination therapies) [167].

However, the selection of the optimal treatment is a difficult task and network analysis may assist in quantifying the rank order of treatments in terms of efficacy/tolerability and outcomes. The direct and indirect analyses revealed two treatments with better effectiveness compared to the reference treatment (EP) for the outcome ORR: 1) combination of Cisplatin, Cyclophosphamide, Etoposide and Epirubicin and 2) combination of Cisplatin, Doxorubicin and Etoposide with Vincristine (intensified), respectively. But, the former combination showed worst tolerability than EP.

On the contrary, seven other regimens showed worse effectiveness for the ORR outcome (Cisplatin plus Doxorubicin plus Etoposide plus Vincristine (intensified), Ifosfamide plus Mesna, Carboplatin plus Pemetrexeb, Doxorubicin plus Etoposide plus Vincristine, Cyclophosphamide plus Doxorubicin plus Etoposide, Teniposide and Cisplatin) and three regimens for the both outcomes (ORR and CR) (Etoposide standard, Etoposide intensified and Carboplatin plus Ifosfamide).

In the frequentist network analysis, possible effect modifiers were not taken into account and only the unadjusted pooled ORs were calculated since data that affect the response were not provided in the individual studies. In addition, the estimated effect sizes were unadjusted for treatment dosage levels. Nevertheless, the developed methodology (and of course, the NET-MS system) cannot estimate adjusted effect sizes; though; it has the capability of subgroup analyses.

With a frequentist approach, result of analysis is presented as a point estimate with a 95% CI. However, these CIs cannot be interpreted in terms of probabilities; this shortcoming could be overcome by the use of Bayesian methods which presents probabilities that can predict and is of relevance to the decision maker [25, 165]. These methods assume prior probability distribution, prior belief of possible values of model parameter based on what is already



known on the subject. Then in the light of observed data in the study, likelihood distribution of these parameters is used to obtain a corresponding posterior probability distribution. For NMA, specific advantage is that the posterior probability distribution allows calculating the probability of the competing interventions. Results are expressed in credible intervals as opposed to the CI in case of frequentist analysis. Other advantages of Bayesian meta-analysis include the straightforward way to make predictions and possibility to incorporate different sources of uncertainty [168].

We have performed the same analysis using Bayesian statistics and in principle we reached out to similar results.

However, when we performed an adjusted analysis to ED-SCLC (12 studies, 5 interventions, 3'961 patients, and 10 possible pairwise comparisons\_5 of those with direct data), it was revealed that the combination of platin plus paclitaxel might be a good first line option for achieving ORR (SUCRA 91%), followed by platin plus ambrubicin (73%), while the combination of platin plus etoposide provides the lowest probability of success (SUCRA 15%).

An additional analysis performed including some of the investigational treatments with available results until mid-2016; now the network was increased to 17 studies, comparing 11 interventions, including 4'605 patients and 55 pairwise comparisons (12 of those with direct data). As it was expected, new treatments are improve the probabilities of achieving ORR in ED-SCLC, but not significantly differentiated from current standard of care and further investigation is needed to drive conclusions on how these treatments would be better utilized in clinical practice.

The applicability of previously mentioned analyses is questionable due to several limitations; the differences of the dosage schemes and/or treatment cycle maximum duration were ignored since we focused to the antitumor activity of each treatment based on the mode of action of each chemotherapy agent (or combination of individual chemotherapy agents). We adopted this approach since the scientific evidence of the relative anti-tumor activity of each chemotherapy agent, or combination of individual chemotherapy agents is relative scarce.

In addition, the existence of publication bias (defining as the differential magnitude of effect in large versus small studies) cannot totally be excluded [169]. However, a valid method for testing publication bias in network analysis does not exist. Also, in the network analysis, adjustments for multiple comparisons may not be applicable since the purpose of the analysis was to explore the relative significance of risk effect [170].

Data were synthesized with an objective (to assess the relative effectiveness of treatments) but not with a pre-specified key hypothesis [170-172]. An appropriate multiple test adjustment is difficult or even impossible because the investigated comparisons in the network are not independent and a clear structure in the multiple tests is missing [172].

Finally, the existence of false positive results may not be totally excluded since heterogeneity between studies within the network cannot be assessed (lack of valid methodology) and the network analysis cannot adjust for possible effect modifiers; though, synthesis of data from many studies usually is expected to reduce false discovery rate.

Although the quality of reporting of the studies included in the network-analysis was assessed, a sensitivity analysis involving the studies with high reporting quality was not considered since the aim of the assessment was to obtain an indication of the reporting quality of the current evidence in SCLC treatment; in addition, there is no established quality scales to divide “high-quality” from “low-quality” studies. Furthermore, it has been shown

that individual quality measures are not associated with treatment effect size across studies and medical areas [173].

Also, the analysis was not restricted to specific subpopulations (e.g. limited and extensive-stage SCLC) due to lack of replication and to achieve greater power in detecting significant results. Since the indirect comparisons are not randomized but observational studies across trials the differences in study populations and prognostic factors across RCTs may lead to overestimation of the treatment effects [1, 9].

In addition, the network analysis was based on grouped data from published RCTs and not on individual patient data, assuming that the relative effectiveness of a treatment is consistent in different RCTs. Therefore, the results regarding the superiority of a particular treatment should be interpreted with great caution. However, when the previous basic assumption may not be met, the results of one RCT can be not generalizable to another; though, the identification of factors that may influence the generalizability of an RCT is rather difficult [174].

## Charter 8 – Conclusions

In conclusion, network meta-analyses can be considered an extension of traditional meta-analysis by including multiple different pairwise comparisons across a range of different interventions to allow multiple treatment comparisons in the absence of head-to-head evidence. Furthermore, the methodology can combine direct and indirect treatment comparisons, thereby synthesizing a greater share of the available evidence than a traditional meta-analysis

It is essential to assess the internal validity (appropriate identification of studies that form the evidence network, quality of the individual RCTs, and extent of confounding bias due to similarity and consistency violations) and external validity (external validity of RCTs included in the evidence network) of the network , prior initiating an exercise.

From methodological perspective, Bayesian and frequentist approaches should be complementary tools; if the user's approach to a clinical problem places an emphasis on identifying causal relationships, a frequentist approach might be best suited. On the other hand, if the user takes an approach in which estimating a priori probabilities is appropriate, a Bayesian approach might be more appropriate. Ideally both approaches should be used for the same study.

# References

- [1] Higgins JPT, Green S (eds) (2008) Cochrane handbook for systematic reviews of interventions. Wiley, Chichester
- [2] Moher D, Liberati A, Tetzlaff J, et al. Preferred reporting items for systematic reviews and metaanalyses: the PRISMA statement. *Ann Intern Med*. 2009 Aug 18;151(4):264-9, W64. Epub 2009 Jul 20
- [3] Caldwell DM, Ades AE, Higgins JP. Simultaneous comparison of multiple treatments: combining direct and indirect evidence. *BMJ*. 2005 Oct 15;331(7521):897-900.
- [4] Salanti G. Indirect and mixed-treatment comparison, network, or multiple treatments meta-analysis: many names, many benefits, many concerns for the next generation evidence synthesis tool. *Research Synthesis Methods*. 2012;3:80-97.
- [5] Ioannidis JP. Integration of evidence from multiple metaanalyses: a primer on umbrella reviews, treatment networks and multiple treatments meta-analyses. *CMAJ*. 2009 Oct 13;181(8):488-9
- [6] Mills EJ, Ioannidis JP, Thorlund K, et al. How to use an article reporting a multiple treatment comparison meta-analysis. *JAMA*. 2012 Sep 26;308(12):1246-53.
- [7] Mills EJ, Thorlund K, Ioannidis JP. Demystifying trial networks and network meta-analysis. *BMJ*. 2013 May 14;346:f2914
- [8] Cipriani A, Higgins JP, Geddes JR, et al. Conceptual and technical challenges in network meta-analysis. *Ann Intern Med*. 2013 Jul 16;159(2):130-7
- [9] Higgins JP, Whitehead A. Borrowing strength from external trials in a meta-analysis. *Stat Med*. 1996 Dec 30;15(24):2733-49
- [10] Bucher HC, Guyatt GH, Griffith LE, et al. The results of direct and indirect treatment comparisons in metaanalysis of randomized controlled trials. *J Clin Epidemiol*. 1997 Jun;50(6):683-91

- [11] Lumley T. Network meta-analysis for indirect treatment comparisons. *Stat Med*. 2002 Aug 30;21(16):2313-24
- [12] Lu G, Ades AE. Combination of direct and indirect evidence in mixed treatment comparisons. *Stat Med*. 2004 Oct 30;23(20):3105-24.
- [13] Nikolakopoulou A, Chaimani A, Veroniki AA, et al. Characteristics of Networks of Interventions: A Description of a Database of 186 Published Networks. *PLOS ONE* 2014;9:e86754.
- [14] Bafeta A, Trinquart L, Seror R, et al. Analysis of the systematic reviews process in reports of network meta-analyses: methodological systematic review. *BMJ*. 2013 Jul 1;347:f3675.
- [15] Lee AW. Review of mixed treatment comparisons in published systematic reviews shows marked increase since 2009. *J Clin Epidemiol* 2014;67:138–43.
- [16] Achana F, Hubbard S, Sutton A, et al. An exploration of synthesis methods in public health evaluations of interventions concludes that the use of modern statistical methods would be beneficial. *J Clin Epidemiol*. 2014 Apr;67(4):376-90
- [17] Higgins JPT, Welton NJ. Network meta-analysis: a norm for comparative effectiveness? *Lancet Lond Engl* 2015;386:628–30.
- [18] Petropoulou M, Nikolakopoulou A, Veroniki AA et al. Bibliographic study showed improving statistical methodology of network meta-analyses published between 1999 and 2015. *J Clin Epidemiol*. 2016 Nov 15. pii: S0895-4356(16)30256-6
- [19] Song F, Loke YK, Walsh T, et al. Methodological problems in the use of indirect comparisons for evaluating healthcare interventions: survey of published systematic reviews. *BMJ*. 2009;338:b1147.
- [20] Donegan S, Williamson P, Gamble C, et al. Indirect comparisons: a review of reporting and methodological quality. *PLoS One*. 2010;5:e11054.
- [21] Baker SG, Kramer BS. The transitive fallacy for randomized trials: if A bests B and B bests C in separate trials, is A better than C? *BMC Med Res Methodol*. 2002;2:13.

- [22] Lu G, Ades AE. Assessing evidence inconsistency in mixed treatment comparisons. *J Am Stat Assoc.* 2006;101:447-59.
- [23] Salanti G, Marinho V, Higgins JP. A case study of multiple-treatments meta-analysis demonstrates that covariates should be considered. *J Clin Epidemiol.* 2009;62:857-64.
- [24] Cooper NJ, Sutton AJ, Morris D, et al. Addressing between-study heterogeneity and inconsistency in mixed treatment comparisons: application to stroke prevention treatments in individuals with non-rheumatic atrial fibrillation. *Stat Med.* 2009;28:1861-81.
- [25] Jansen JP, Schmid CH, Salanti G. Directed acyclic graphs can help understand bias in indirect and mixed treatment comparisons. *J Clin Epidemiol.* 2012; 65:798-807.
- [26] Higgins JP, Thompson SG, Deeks JJ, et al. Measuring inconsistency in meta-analyses. *BMJ.* 2003;327:557-60.
- [27] Dias S, Welton NJ, Ades AE. Study designs to detect sponsorship and other biases in systematic reviews. *J Clin Epidemiol.* 2010;63:587-8.
- [28] White IR, Barrett JK, Jackson D, et al. Consistency and inconsistency in network meta-analysis: model estimation using multivariate metaregression. *Research Synthesis Methods.* 2012;3:111-25.
- [29] Seifter EJ, Ihde DC. Therapy of small cell lung cancer: a perspective on two decades of clinical research. *Semin Oncol* 1988;15:278-99
- [30] Evans WK, Feld R, Murray N, et al. Superiority of alternating non-cross-resistant chemotherapy in extensive small cell lung cancer: a multicenter, randomized clinical trial by the National Cancer Institute of Canada. *Ann Intern Med* 1987;107(4):451-8
- [31] Fukuoka M, Furuse K, Saijo N, et al. Randomized trial of cyclophosphamide, doxorubicin, and vincristine versus cisplatin and etoposide versus alternation of these regimens in small-cell lung cancer. *J Natl Cancer Inst* 1991;83(12):855-61
- [32] Roth BJ, Johnson DH, Einhorn LH, et al. Randomized study of cyclophosphamide, doxorubicin, and vincristine versus etoposide and cisplatin versus alternation of these two regimens in extensive small-cell lung cancer: a phase III trial of the Southeastern Cancer Study Group. *J Clin Oncol* 1992;10(2):282-91

- [33] Chute JP, Chen T, Feigal E, et al. Twenty years of phase III trials for patients with extensive-stage small-cell lung cancer: perceptible progress. *J Clin Oncol* 1999;17(6):1794-801
- [34] Johnson D. Management of small cell lung cancer: current state of the art. *Chest* 1999;116(Suppl):525S-30S
- [35] Pignon JP, Arriagada R, Ihde DC et al. A meta-analysis of thoracic radiotherapy for small-cell lung cancer. *N Engl J Med*. 1992 Dec 3;327(23):1618-24
- [36] Warde P, Payne D. Does thoracic irradiation improve survival and local control in limited-stage small-cell carcinoma of the lung? A meta-analysis. *J Clin Oncol*. 1992 Jun;10(6):890-5
- [37] Fried DB, Morris DE, Poole C et al. Systematic review evaluating the timing of thoracic radiation therapy in combined modality therapy for limited-stage small cell lung cancer. *J Clin Oncol*. 2004 Dec 1;22(23):4837-45
- [38] Turrisi AT, Kim K, Blum R et al. Twice-daily compared with once-daily thoracic radiotherapy in limited small-cell lung cancer treated concurrently with cisplatin and etoposide. *N Engl J Med*. 1999 Jan 28;340(4):265-71
- [39] Auperin A, Arriagada R, Pignon JP et al. Prophylactic cranial irradiation for patients with small-cell lung cancer in complete remission *N Engl J Med*. 1999 Aug 12;341(7):476-84
- [40] Slotman B, Faivre-Finn C, Kramer G et al. Prophylactic cranial irradiation in extensive small-cell lung cancer. *N Engl J Med*. 2007 Aug 16;357(7):664-72.
- [41] Hanna NH, Einhorn LH. Small-cell lung cancer: state of the art. *Clin Lung Cancer*. 2002 Sep;4(2):87-94.
- [42] Von Pawel J, Schiller JH, Shepherd FA et al. Topotecan versus cyclophosphamide, doxorubicin, and vincristine for the treatment of recurrent small-cell lung cancer. *J Clin Oncol*. 1999 Feb;17(2):658-67
- [43] O'Brien MER, Ciuleanu TE, Tsekov H et al. Phase III trial comparing supportive care alone with supportive care with oral topotecan in patients with relapsed small-cell lung cancer. *J Clin Oncol*. 2006 Dec 1;24(34):5441-7



- [44] Chute JP, Chen T, Feigal E et al. Twenty years of phase III trials for patients with extensive-stage small cell lung cancer: perceptible progress. *J Clin Oncol*. 1999 Jun;17(6):1794-801
- [45] Johnson SW, O'Dwyer PJ. Pharmacology of cancer chemotherapy: cisplatin and its analogues. In: DeVita VT Jr, Hellman S, Rosenberg SA, editors. *Cancer: principles and practice of oncology*. 7th edition. Lippincott Williams & Wilkins; Philadelphia, PA: 2005. p. 344-58
- [46] Joss RA, Alberto P, Hu'rnny C, et al. Quality versus quantity of life in the treatment of patients with advanced small-cell lung cancer? A randomized phase III comparison of weekly carboplatin and teniposide versus cisplatin, adriamycin, etoposide alternating with cyclophosphamide, methotrexate, vincristine and lomustine: Swiss Group for Clinical Cancer Research (SAKK). *Ann Oncol*. 1995 Jan;6(1):41-8
- [47] Skarlos DV, Samantas E, Kosmidis P, et al. Randomized comparison of etoposide-cisplatin vs. etoposide-carboplatin and irradiation in small-cell lung cancer: a Hellenic Co-operative Oncology Group study. *Ann Oncol*. 1994 Sep;5(7):601-7.
- [48] Okamoto H, Watanabe K, Kunikane H, et al. Randomised phase III trial of carboplatin plus etoposide vs split doses of cisplatin plus etoposide in elderly or poor-risk patients with extensive disease small-cell lung cancer: JCOG 9702. *Br J Cancer*. 2007 Jul 16;97(2):162-9
- [49] Lee SM, James LE, Qian W, et al. Comparison of gemcitabine and carboplatin versus cisplatin and etoposide for patients with poor-prognosis small cell lung cancer. *Thorax*. 2009 Jan;64(1):75-80
- [50] Rossi A, Di Maio M, Chiodini P, et al. Carboplatin- or cisplatin-based chemotherapy in first-line treatment of small-cell lung cancer: the COCIS meta-analysis of individual patient data. *J Clin Oncol*. 2012 May 10;30(14):1692-8.
- [51] Noda K, Nishiwaki Y, Kawahara M, et al. Irinotecan plus cisplatin compared with etoposide plus cisplatin for extensive small-cell lung cancer. *N Engl J Med*. 2002 Jan 10;346(2):85-91
- [52] Hanna N, Bunn PA, Langer C, et al. Randomized phase III trial comparing irinotecan/cisplatin with etoposide/ cisplatin in patients with previously untreated extensive-stage small-cell lung cancer. *J Clin Oncol*. 2006 May 1;24(13):2038-43.

- [53] Lara PN, Natale R, Crowley J, et al. Phase III trial of irinotecan/cisplatin compared with etoposide/cisplatin in extensive-stage small-cell lung cancer: clinical and pharmacogenomic results from SWOG S0124. *J Clin Oncol*. 2009 May 20;27(15):2530-5
- [54] Zatloukal P, Cardenal F, Szxzesna A, et al. A multicenter international randomized phase III study comparing cisplatin in combination with irinotecan or etoposide in previously untreated small-cell lung cancer patients with extensive disease. *Ann Oncol*. 2010 Sep;21(9):1810-6
- [55] Kubota K, Hida T, Ishikura S, et al. Etoposide and cisplatin versus irinotecan and cisplatin in patients with limited-stage small-cell lung cancer treated with etoposide and cisplatin plus concurrent accelerated hyperfractionated thoracic radiotherapy (JCOG0202): a randomized phase 3 study. *Lancet Oncol*. 2014 Jan;15(1):106-13
- [56] Hermes A, Bergman B, Bremmew R, et al. Irinotecan plus carboplatin versus oral etoposide plus carboplatin in extensive small-cell lung cancer: a randomized phase III trial. *J Clin Oncol*. 2008 Sep 10;26(26):4261-7
- [57] Schmittel A, von Weikersthal LF, Sebastian M, et al. A randomized phase II trial of irinotecan plus carboplatin versus etoposide plus carboplatin treatment in patients with extended disease small-cell lung cancer. *Ann Oncol*. 2006 Apr;17(4):663-7
- [58] Eckardt JR, von Pawel J, Papai Z, et al. Open-label, multicenter, randomized, phase III study comparing oral topotecan/cisplatin versus etoposide/ cisplatin as treatment for chemotherapy-naïve patients with extensive-disease small-cell lung cancer. *J Clin Oncol*. 2006 May 1;24(13):2044-51
- [59] Fink TH, Huber RM, Heigener DF, et al. Topotecan/cisplatin compared with cisplatin/etoposide as first-line treatment for patients with extensive disease small-cell lung cancer. *J Thorac Oncol*. 2012 Sep;7(9):1432-9
- [60] Jiang J, Liang X, Zhou X, et al. A meta-analysis of randomized controlled trials comparing irinotecan/platinum with etoposide/platinum in patients with previously untreated extensive-stage small cell lung cancer. *J Thorac Oncol*. 2010 Jun;5(6):867-73
- [61] Lima JP, dos Santos LV, Sasse EC, et al. Camptothecins compared with etoposide in combination with platinum analog in extensive stage small cell lung cancer: systematic review with meta-analysis. *J Thorac Oncol*. 2010 Dec;5(12):1986-93

- [62] Thomas P, Castelnau O, Paillot D, et al. Phase II trial of paclitaxel and carboplatin in metastatic small-cell lung cancer: a Groupe Francais de Pneumo-Cancerologie study. *J Clin Oncol*. 2001 Mar 1;19(5):1320-5
- [63] Lyss AP, Herndon JE, Lynch TJ, et al. Novel doublets in extensive-stage small-cell lung cancer: a randomized phase II study of topotecan plus cisplatin or paclitaxel (CALGB 9430). *Clin Lung Cancer*. 2002 Feb;3(3):205-10
- [64] Leyvraz S, Pampallona S, Martinelli G, et al. A threefold dose intensity treatment with ifosfamide, carboplatin, and etoposide for patients with small cell lung cancer: a randomized trial. *J Natl Cancer Inst*. 2008 Apr 16;100(8):533-41
- [65] Lorigan P, Woll PJ, O'Brien MER, et al. Randomized phase III trial of dose-dense chemotherapy supported by whole-blood hematopoietic progenitors in better-prognosis small-cell lung cancer. *J Natl Cancer Inst*. 2005 May 4;97(9):666-74
- [66] Furuse K, Fukuoka M, Nishiwaki Y, et al. Phase III study of intensive weekly chemotherapy with recombinant human granulocyte colony-stimulating factor versus standard chemotherapy in extensive-disease small-cell lung cancer. *J Clin Oncol*. 1998 Jun;16(6):2126-32
- [67] Niell HB, Herndon JE, Miller AA, et al. Randomized phase III Intergroup trial of etoposide and cisplatin with or without paclitaxel and granulocyte colony-stimulating factor in patients with extensive-stage small-cell lung cancer: cancer and Leukemia Group B trial 9732J *Clin Oncol*. 2005 Jun 1;23(16):3752-9.
- [68] Elias A, Ibrahim J, Skarin AT, et al. Dose-intensive therapy for limited-stage small-cell lung cancer: long-term outcome. *J Cancer Res Clin Oncol*. 2011 Aug;137(8):1175-84
- [69] Roth BJ, Johnson DH, Einhorn LH, et al. Randomized study of cyclophosphamide, doxorubicin, and vincristine versus etoposide and cisplatin versus alternation of these two regimens in extensive small-cell lung cancer: a phase III trial of the Southeastern Cancer Study Group. *J Clin Oncol*. 1992 Feb;10(2):282-91
- [70] Rossi A, Garassino MC, Cinquini M, et al. Maintenance or consolidation therapy in small-cell lung cancer: a systematic review and meta-analysis. *Lung Cancer*. 2010 Nov;70(2):119-28

- [71] Schiller JH, Adak S, Cella D, et al. Topotecan versus observation after cisplatin plus etoposide in extensive-stage small-cell lung cancer: E7593 - a phase III trial of the Eastern Cooperative Oncology Group. *J Clin Oncol*. 2001 Apr 15;19(8):2114-22
- [72] Hanna NH, Einhorn LH. Small-cell lung cancer: state of the art. *Clin Lung Cancer*. 2002 Sep;4(2):87-94
- [73] Yana T, Negoro S, Takada M, et al. Phase II study of amrubicin in previously untreated patients with extensive-disease small cell lung cancer: west Japan Thoracic Oncology Group study. *Invest New Drugs*. 2007 Jun;25(3):253-8
- [74] Ohe Y, Negoro S, Matsui K, et al. Phase I-II study of amrubicin and cisplatin in previously untreated patients with extensive-stage small cell lung cancer. *Ann Oncol*. 2005 Mar;16(3):430-6
- [75] Satouchi M, Kotani Y, Shibata T, et al. Phase III study comparing amrubicin plus cisplatin with irinotecan plus cisplatin in the treatment of extensive-disease small-cell lung cancer: JCOG 0509. *J Clin Oncol*. 2014 Apr 20;32(12):1262-8
- [76] Onoda S, Masuda N, Seto T, et al. Phase II trial of amrubicin for the treatment of refractory or relapsed small-cell lung cancer: thoracic Oncology Research Group study 0301. *J Clin Oncol*. 2006 Dec 1;24(34):5448-53
- [77] Igawa S, Yamamoto N, Ueda S, et al. Evaluation of the recommended dose and efficacy of amrubicin as second- and third-line chemotherapy for small cell lung cancer. *J Thorac Oncol*. 2007 Aug;2(8):741-4
- [78] Ettinger DS, Jotte R, Lorigan P, et al. Phase II study of amrubicin as second-line therapy in patients with platinum-refractory small-cell lung cancer. *J Clin Oncol*. 2010 May 20;28(15):2598-603
- [79] Shimokawa T, Shibuya M, Kitamura K, et al. Retrospective analysis of efficacy and safety of amrubicin in refractory and relapsed small-cell lung cancer. *Int J Clin Oncol*. 2009 Feb;14(1):63-9
- [80] Inoue A, Sugawara S, Yamazaki K, et al. Randomized phase II trial comparing amrubicin with topotecan in patients with previously treated small-cell lung cancer: north Japan Lung Cancer Study Group trial 0402. *J Clin Oncol*. 2008 Nov 20;26(33):5401-6

- [81] Jotte R, Conkling P, Reynolds C, et al. Randomized phase II trial of single-agent amrubicin or topotecan as second-line treatment in patients with small-cell lung cancer sensitive to first-line platinum-based chemotherapy. *J Clin Oncol*. 2011 Jan 20;29(3):287-93
- [82] Jotte R, von Pawel J, Spigel DR, et al. Randomized phase III trial of amrubicin versus topotecan as second-line treatment for small cell lung cancer. *J Clin Oncol* 2011;29(15S):453s
- [83] Sharp A, Bhosle J, Abdelraouf F, et. al. Development of molecularly targeted agents and immunotherapies in small cell lung cancer. *Eur J Cancer*. 2016 Jun;60:26-39
- [84] Belani C, Dahlberg S, Rudin C, et al. Three-arm randomized phase II study of cisplatin and etoposide (CE) versus CE with either vismodegib (V) or cixutumumab (Cx) for patients with extensive stage-small cell lung cancer (ES-SCLC) (ECOG 1508). *J Clin Oncol* 2013;31.
- [85] Johnson BE, Fischer T, Fischer B, et al. Phase II study of imatinib in patients with small cell lung cancer. *Clin Cancer Res*. 2003 Dec 1;9(16 Pt 1):5880-7
- [86] Langer CJ, Albert I, Ross HJ, et al. Randomized phase II study of carboplatin and etoposide with or without obatoclox mesylate in extensive-stage small cell lung cancer. *Lung Cancer* 2014;85(3):420e8.
- [87] Owonikoko TK, Zhang G, Deng X, et al. Poly (ADP) ribose polymerase enzyme inhibitor, veliparib, potentiates chemotherapy and radiation in vitro and in vivo in small cell lung cancer. *Cancer Med*. 2014 Dec;3(6):1579-94
- [88] Pandya KJ, Dahlberg S, Hidalgo M, et al. A randomized, phase II trial of two dose levels of temsirolimus (CCI-779) in patients with extensive-stage small-cell lung cancer who have responding or stable disease after induction chemotherapy: a trial of the Eastern Cooperative Oncology Group (E1500). *J Thorac Oncol* 2007;2(11):1036e41.
- [89] Patton D, Spigel R, Greco FA, et al. Irinotecan (I), carboplatin (C), and radio-therapy (RT) followed by maintenance bevacizumab (B) in the treatment (tx) of limited-stage small cell lung cancer (LS-SCLC): update of a phase II trial of the Minnie Pearl Cancer Research Network. *J Clin Oncol* 2006;24(18S):7085.

- [90] Ready NE, Dudek AZ, Pang HH, et al. Cisplatin, irinotecan, and bevacizumab for untreated extensive-stage small-cell lung cancer: CALGB 30306, a phase II study. *J Clin Oncol* 2011;29(33):4436e41.
- [91] Ready NE, Pang HH, Gu L, et al. Chemotherapy with or without maintenance sunitinib for untreated extensive-stage small-cell lung cancer: a randomized, double-blind, placebo-controlled phase II Study. *J Clin Oncol* 2015 May 20;33(15):1660e5.
- [92] Rudin CM, Salgia R, Wang X, et al. Randomized phase II Study of carboplatin and etoposide with or without the bcl-2 antisense oligonucleotide oblimersen for extensive-stage small-cell lung cancer: CALGB 30103. *J Clin Oncol* 2008;26(6):870e6.
- [93] Schneider BJ, Kalemkerian GP, Ramnath N, et al. Phase II trial of imatinib main-tenance therapy after irinotecan and cisplatin in patients with c-Kit-positive, extensive-stage small-cell lung cancer. *Clin Lung Cancer* 2010;11(4):223e7.
- [94] Sharma N, Pennell N, Nickolich M, et al. Phase II trial of sorafenib in conjunction with chemotherapy and as maintenance therapy in extensive-stage small cell lung cancer. *Invest New Drugs* 2014;32(2):362e8.
- [95] Spigel DR, Greco FA, Rubin MS, et al. Phase II study of maintenance sunitinib following irinotecan and carboplatin as first-line treatment for patients with extensive-stage small-cell lung cancer. *Lung Cancer* 2012;77(2):359e64.
- [96] Spigel DR, Greco FA, Zubkus JD, et al. Phase II trial of irinotecan, carboplatin, and bevacizumab in the treatment of patients with extensive-stage small-cell lung cancer. *J Thorac Oncol* 2009;4(12):1555e60.
- [97] Spigel DR, Hainsworth JD, Farley C, et al. Tracheoesophageal (TE) fistula development in a phase II trial of concurrent chemoradiation (CRT) and bev-acizumab (B) in limited-stage small-cell lung cancer (LS-SCLC). *J Clin Oncol* 2008;26(15S):7554.
- [98] Spigel DR, Hainsworth JD, Simons L, et al. Irinotecan, carboplatin, and imatinib in untreated extensive-stage small-cell lung cancer: a phase II trial of the Minnie Pearl Cancer Research Network. *J Thorac Oncol* 2007;2(9):854e61.

- [99] Spigel DR, Townley PM, Waterhouse DM, et al. Randomized phase II study of bevacizumab in combination with chemotherapy in previously untreated extensive-stage small-cell lung cancer: results from the SALUTE trial. *J Clin Oncol* 2011;29(16):2215e22.
- [100] Allen JW, Moon J, Redman M, et al. Southwest Oncology Group S0802: a randomized, phase II trial of weekly topotecan with and without ziv-aflibercept in patients with platinum-treated small-cell lung cancer. *J Clin Oncol* 2014;32(23):2463e70.
- [101] de Marinis F, Atmaca A, Tiseo M, et al. A phase II study of the histone deacetylase inhibitor panobinostat (LBH589) in pretreated patients with small-cell lung cancer. *J Thorac Oncol* 2013;8(8):1091e4.
- [102] Gandhi L, Heist R, Lucca J, et al. A phase II trial of pazopanib in relapsed/refractory small cell lung cancer (SCLC). *J Clin Oncol* 2012;30.
- [103] Gitlitz BJ, Moon J, Glisson BS, et al. Sorafenib in platinum-treated patients with extensive stage small cell lung cancer: a Southwest Oncology Group (SWOG 0435) phase II trial. *J Thorac Oncol* 2010;5(11):1835e40.
- [104] Han JY, Kim HY, Lim KY, et al. A phase II study of sunitinib in patients with relapsed or refractory small cell lung cancer. *Lung Cancer* 2013;79(2):137e42.
- [105] Heist RS, Fain J, Chinnasami B, et al. Phase I/II study of AT-101 with topotecan in relapsed and refractory small cell lung cancer. *J Thorac Oncol* 2010;5(10):1637e43.
- [106] Jalal S, Bedano P, Einhorn L, et al. Paclitaxel plus bevacizumab in patients with chemosensitive relapsed small cell lung cancer: a safety, feasibility, and efficacy study from the Hoosier Oncology Group. *J Thorac Oncol* 2010; 5(12):2008e11.
- [107] Krug LM, Crapanzano JP, Azzoli CG, et al. Imatinib mesylate lacks activity in small cell lung carcinoma expressing c-kit protein: a phase II clinical trial. *Cancer* 2005;103(10):2128e31.
- [108] Lara Jr PN, Chansky K, Davies AM, et al. Bortezomib (PS-341) in relapsed or refractory extensive stage small cell lung cancer: a Southwest Oncology Group phase II trial (S0327). *J Thorac Oncol* 2006;1(9):996e1001.

- [109] Moore AM, Einhorn LH, Estes D, et al. Gefitinib in patients with chemo-sensitive and chemo-refractory relapsed small cell cancers: a Hoosier Oncology Group phase II trial. *Lung Cancer* 2006;52(1):93e7.
- [110] Mountzios G, Emmanouilidis C, Vardakis N, et al. Paclitaxel plus bevacizumab in patients with chemoresistant relapsed small cell lung cancer as salvage treatment: a phase II multicenter study of the Hellenic Oncology Research Group. *Lung Cancer* 2012;77(1):146e50.
- [111] Rudin CM, Hann CL, Garon EB, et al. Phase II study of single-agent navitoclax (ABT-263) and biomarker correlates in patients with relapsed small cell lung cancer. *Clin Cancer Res* 2012;18(11): 3163e9.
- [112] Tarhini A, Kotsakis A, Gooding W, et al. Phase II study of everolimus (RAD001) in previously treated small cell lung cancer. *Clin Cancer Res* 2010; 16(23):5900e7.
- [113] Wainberg Z, Rafii S, Ramanathan R, et al. Safety and antitumor activity of the PARP inhibitor BMN673 in a phase 1 trial recruiting metastatic small-cell lung cancer (SCLC) and germline BRCA-mutation carrier cancer patients. *J Clin Oncol* 2014;32:5S.
- [114] Waterhouse DM, Morgan SK, Spigel DR, et al. Phase II study of oral topotecan plus bevacizumab (topo-bev) for second-line treatment of small cell lung cancer (SCLC). *J Clin Oncol* 2010;28(15).
- [115] Putton JL, Lavole A, Quoix E, et al. Randomized phase II-III study of bevacizumab in combination with chemotherapy in previously untreated extensive small-cell lung cancer: results from the IFCT-0802 trial dagger. *Ann Oncol* 2015;26(5):908e14.
- [116] Rudin CM, Salgia R, Wang X, et al. Randomized phase II Study of carboplatin and etoposide with or without the bcl-2 antisense oligonucleotide oblimersen for extensive-stage small-cell lung cancer: CALGB 30103. *J Clin Oncol*. 2008 Feb 20;26(6):870-6.
- [117]. Langer CJ, Albert I, Ross HJ et al. Randomized phase II study of carboplatin and etoposide with or without obatoclax mesylate in extensive-stage small cell lung cancer, *Lung Cancer*. 2014 Sep;85(3):420-8.



- [118] Reck M, Bondarenko I, Luft A et al. Ipilimumab in combination with paclitaxel and carboplatin as first-line therapy in extensive-disease-small-cell lung cancer: results from a randomized, double-blind, multicenter phase 2 trial. *Ann Oncol*. 2013 Jan;24(1):75-83.
- [119] Antonia SJ, López-Martin JA, Bendell J et. al. Nivolumab alone and nivolumab plus ipilimumab in recurrent small-cell lung cancer (CheckMate 032): a multicentre, open-label, phase 1/2 trial. *Lancet Oncol*. 2016 Jul;17(7):883-95.
- [120] Beck A1, Lambert J, Sun M,et al. Fourth World Antibody-Drug Conjugate Summit: February 29-March 1, 2012, Frankfurt, Germany.MAbs. 2012 Nov-Dec;4(6):637-47.
- [121] Rudin CM, Ismaila N, Hann CL et al. Treatment of Small-Cell Lung Cancer: American Society of Clinical Oncology Endorsement of the American College of Chest Physicians Guideline. *J Clin Oncol*. 2015 Dec 1;33(34):4106-11.
- [122] Miller AB, Hoogstraten B, Staquet M, et al. Reporting results of cancer treatment. *Cancer*. 1981 Jan 1;47(1):207-14
- [123] Eisenhauer EA, Therasse P, Bogaerts J, et al. New response evaluation criteria in solid tumours: Revised RECIST guideline (version 1.1). *Eur J Cancer*. 2009 Jan;45(2):228-47
- [124] Peppercorn JM, Smith TJ, Helft PR, et al. American Society of Clinical Oncology statement: Toward individualized care for patients with advanced cancer. *J Clin Oncol*. 2011 Feb 20;29(6):755-60
- [125] Gill S, Sargent D. End points for adjuvant therapy trials: Has the time come to accept disease-free survival as a surrogate end point for overall survival? *Oncologist*. 2006 Jun;11(6):624-9.
- [126] Prentice RL. Surrogate endpoints in clinical trials: Definition and operational criteria. *Stat Med*. 1989 Apr;8(4):431-40.
- [127] Freedman LS, Graubard BI, Schatzkin A. Statistical validation of intermediate endpoints for chronic diseases. *Stat Med*. 1992 Jan 30;11(2):167-78
- [128] Buyse M, Molenberghs G. Criteria for the validation of surrogate endpoints in randomized experiments. *Biometrics*. 2009 Jun;65(2):521-9.

- [129] Begg CB, Leung D. On the use of surrogate end points in randomized trials. *Stat Med*. 2011 Jan 30;30(2):186-96
- [130] Foster NR, Qi Y, Shi Q et al. Tumor response and progression-free survival as potential surrogate endpoints for overall survival in extensive stage small-cell lung cancer: findings on the basis of North Central Cancer Treatment Group trials. *Cancer*. 2011 Mar 15;117(6):1262-71.
- [131] Foster NR, Renfro LA, Schild SE, et al. Multitrial Evaluation of Progression-Free Survival as a Surrogate End Point for Overall Survival in First-Line Extensive-Stage Small-Cell Lung Cancer. *J Thorac Oncol*. 2015 Jul;10(7):1099-106
- [132] Booth CM, Eisenhauer EA. Progression-free survival: meaningful or simply measurable? *J Clin Oncol*. 2012 Apr 1;30(10):1030-3
- [133] Burzykowski T, Molenberghs G, Buyse M, et al. Validation of surrogate endpoints in multiple randomized clinical trials with failure time endpoints. *Appl Stat* 2001;50:405–422.
- [134] Buyse M, Burzykowski T, Carroll K, et al. Progression-free survival is a surrogate for survival in advanced colorectal cancer. *J Clin Oncol*. 2007 Nov 20;25(33):5218-24
- [135] Tang PA, Bentzen SM, Chen EX, et al. Surrogate end points for median overall survival in metastatic colorectal cancer: literature-based analysis from 39 randomized controlled trials of first-line chemotherapy. *J Clin Oncol*. 2007 Oct 10;25(29):4562-8.
- [136] Bast RC, Thigpen JT, Arbuck SG, et al. Clinical trial endpoints in ovarian cancer: report of an FDA/ASCO/AACR Public Workshop. *Gynecol Oncol*. 2007 Nov;107(2):173-6..
- [137] Sherrill B, Amonkar M, Wu Y, et al. Relationship between effects on time-to-disease progression and overall survival in studies of metastatic breast cancer. *Br J Cancer*. 2008 Nov 18;99(10):1572-8
- [138] Hackshaw A, Knight A, Barrett-Lee P, et al. Surrogate markers and survival in women receiving first-line combination anthracycline chemotherapy for advanced breast cancer. *Br J Cancer*. 2005 Nov 28;93(11):1215-21.
- [139] Burzykowski T, Buyse M, Piccart-Gebhart MJ, et al. Evaluation of tumor response, disease control, progression-free survival, and time to progression as potential surrogate end points in metastatic breast cancer. *J Clin Oncol*. 2008 Apr 20;26(12):1987-92

- [140] Miksad RA, Zietemann V, Gothe R, et al. Progression-free survival as a surrogate endpoint in advanced breast cancer. *Int J Technol Assess Health Care*. 2008 Fall;24(4):371-83.
- [141] Buyse M, Vangeneugden T, Bijnens L, et al. Validation of biomarkers as surrogates for clinical endpoints. In Bloom JC, Dean RA (Eds.), *Biomarkers in Clinical Drug Development*. New York, NY: Marcel Dekker, 2003. pp 149–168.
- [142] Collette L, Burzykowski T, Carroll KJ, et al. Is prostate-specific antigen a valid surrogate end point for survival in hormonally treated patients with metastatic prostate cancer? Joint research of the European Organisation for Research and Treatment of Cancer, the Limburgs Universitair Centrum, and AstraZeneca Pharmaceuticals. *J Clin Oncol*. 2005 Sep 1;23(25):6139-48
- [143] Paoletti X, Oba K, Bang YJ, et al.; GASTRIC group. Progression-free survival as a surrogate for overall survival in advanced/recurrent gastric cancer trials: a meta-analysis. *J Natl Cancer Inst*. 2013 Nov 6;105(21):1667-70.
- [144] Schmittel A, Sebastian M, Fischer von Weikersthal L et al. A German multicenter, randomized phase III trial comparing irinotecan-carboplatin with etoposide-carboplatin as first-line therapy for extensive-disease small-cell lung cancer. *Ann Oncol*. 2011 Aug;22(8):1798-804.
- [145] Socinski MA, Smit EF, Lorigan P, et al. Phase III study of pemetrexed plus carboplatin compared with etoposide plus carboplatin in chemotherapy-naïve patients with extensive-stage small-cell lung cancer. *J Clin Oncol*. 2009 Oct 1;27(28):4787-92.
- [146] Socinski MA, Weissman C, Hart LL, et al. Randomized phase II trial of pemetrexed combined with either cisplatin or carboplatin in untreated extensive-stage small-cell lung cancer. *J Clin Oncol*. 2006 Oct 20;24(30):4840-7.
- [147] Sorensen M, Pijls-Johannasma M, Felip E; on behalf of the ESMO Guidelines Working Group. Small-cell-lung cancer: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow up. *Ann Oncol* 2010;21(Suppl 5):v120–5.
- [148] DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials* 1986;7(3):177-88.

- [149] Zintzaras E, Lau J. Synthesis of genetic association studies for pertinent gene-disease associations requires appropriate methodological and statistical approaches. *J Clin Epidemiol.* 2008;61(7):634-45.
- [150] Glenny AM, Altman DG, Song F, et al. International Stroke Trial Collaborative Group. Indirect comparisons of competing interventions. *Health Technol Assess* 2005; 9(26): 1-134, iii-iv.
- [151] Whitehead A. *Meta-Analysis of Controlled Clinical Trials*. Chichester, UK: John Wiley & Sons, Ltd, 2002.
- [152] yWorks GmbH, Tübingen, Germany
- [153] Wallace BC, Schmid CH, Lau J, Trikalinos TA. Meta-Analyst: software for meta-analysis of binary, continuous and diagnostic data. *BMC Med Res Methodol.* 2009 Dec 4;9:80.
- [154] Loehrer PJ Sr, Ansari R, Gonin R, et al. Cisplatin plus etoposide with and without ifosfamide in extensive small-cell lung cancer: a Hoosier Oncology Group study. *J Clin Oncol.* 1995;13(10):2594-9.
- [155] Artal-Cortés A, Gomez-Codina J, Gonzalez-Larriba JL, et al. Prospective randomized phase III trial of etoposide/cisplatin versus high-dose epirubicin/cisplatin in small-cell lung cancer. *Clin Lung Cancer.* 2004;6(3):175-83.
- [156] Baka S, Califano R, Ferraldeschi R, et al. Phase III randomized trial of doxorubicin-based chemotherapy compared with platinum-based chemotherapy in small-cell lung cancer. *Br J Cancer* 2008;99(3):442-7
- [157] Grote T, Yeilding AL, Castillo R, et al. Efficacy and safety analysis of epoetin alfa in patients with small-cell lung cancer: a randomized, double-blind, placebo-controlled trial. *J Clin Oncol.* 2005;23(36):9377-86.
- [158] Mavroudis D, Papadakis E, Veslemes M, et al; Greek Lung Cancer Cooperative Group. A multicenter randomized clinical trial comparing paclitaxel-cisplatin-etoposide versus cisplatin-etoposide as first-line treatment in patients with small-cell lung cancer. *Ann Oncol.* 2001;12(4):463-70.

- [159] Miller AA, Herndon JE 2nd, Hollis DR, et al. Schedule dependency of 21-day oral versus 3-day intravenous etoposide in combination with intravenous cisplatin in extensive-stage small-cell lung cancer: a randomized phase III study of the Cancer and Leukemia Group B. *J Clin Oncol*. 1995;13(8):1871-9.
- [160] Miyamoto H, Nakabayashi T, Isobe H, et al. A phase III comparison of etoposide/cisplatin with or without added ifosfamide in small-cell lung cancer. *Oncology*. 1992;49(6):431-5.
- [161] Pujol JL, Daurès JP, Rivière A, et al. Etoposide plus cisplatin with or without the combination of 4'-epidoxorubicin plus cyclophosphamide in treatment of extensive small-cell lung cancer: a French Federation of Cancer Institutes multicenter phase III randomized study. *J Natl Cancer Inst*. 2001;93(4):300-8.
- [162] Rowland KM Jr, Loprinzi CL, Shaw EG, et al. Randomized double-blind placebo-controlled trial of cisplatin and etoposide plus megestrol acetate/placebo in extensive-stage small-cell lung cancer: a North Central Cancer Treatment Group study. *J Clin Oncol*. 1996;14(1):135-41.
- [163] Ruotsalainen TM, Halme M, Tamminen K, et al. Concomitant chemotherapy and IFN-alpha for small cell lung cancer: a randomized multicenter phase III study. *J Interferon Cytokine Res*. 1999;19(3):253-9.
- [164] Wolf M, Havemann K, Holle R, et al. Cisplatin/etoposide versus ifosfamide/etoposide combination chemotherapy in small-cell lung cancer: a multicenter German randomized trial. *J Clin Oncol*. 1987;5(12):1880-9.
- [165] Spiegelhalter DJ, Thomas A, Best NG, et al. WinBUGS User Manual, Version 1.4. Cambridge, UK: MRC Biostatistics Unit, 2002. OpenBUGS. Available from: [www.openbugs.info](http://www.openbugs.info). (Accessed on December 12<sup>th</sup>, 2016)
- [166] Brown S, Hutton B, Clifford T et al. A Microsoft-Excel-based tool for running and critically appraising network meta-analyses—an overview and application of NetMetaXL. *Syst Rev*. 2014 Sep 29;3:110
- [167] van der Valk R, Webers CA, Lumley T, et al. A network meta-analysis combined direct and indirect comparisons between glaucoma drugs to rank effectiveness in lowering intraocular pressure. *J Clin Epidemiol* 2009;62(12):1279-83

- [168] Sutton AJ, Abrams KR, Jones DR, et al. Wiley; 2000. *Methods for Meta-Analysis in Medical Research*.
- [169] Zintzaras E, Voulgarelis M, Moutsopoulos HM. The risk of lymphoma development in autoimmune diseases: a meta-analysis. *Arch Intern Med*. 2005 Nov 14;165(20):2337-44
- [170] Rothman KJ. No adjustments are needed for multiple comparisons. *Epidemiology*. 1990 Jan;1(1):43-6.
- [171] Savitz DA, Olshan AF. Multiple comparisons and related issues in the interpretation of epidemiologic data. *Am J Epidemiol*. 1995 Nov 1;142(9):904-8
- [172] Bender R, Lange S. Adjusting for multiple testing-when and how? *J Clin Epidemiol*. 2001 Apr;54(4):343-9
- [173] Balk EM, Bonis PA, Moskowitz H, Schmid CH, Ioannidis JP, Wang C, Lau J. Correlation of quality measures with estimates of treatment effect in meta-analyses of randomized controlled trials. *JAMA*. 2002 Jun 12;287(22):2973-82
- [174] Black N. Why we need observational studies to evaluate the effectiveness of health care. *BMJ*. 1996 May 11;312(7040):1215-8

# Appendices

## Supplementary Tables

**Supplementary Table 1: List of network meta-analyses in oncology reported 2006-2014**

NMAs in oncology reported 2006-2014	
2006	<ul style="list-style-type: none"> <li>Kyrgiou, M., G. Salanti, et al. (2006). "Survival benefits with diverse chemotherapy regimens for ovarian cancer: meta-analysis of multiple treatments." <u>J Natl Cancer Inst</u> <b>98</b>(22): 1655-1663.</li> </ul>
2007	<ul style="list-style-type: none"> <li>Purkayastha, S., T. Athanasiou, et al. (2007). "Magnetic resonance colonography vs computed tomography colonography for the diagnosis of colorectal cancer: an indirect comparison." <u>Colorectal Dis</u> <b>9</b>(2): 100-111.</li> </ul>
2008	<ul style="list-style-type: none"> <li>Coleman, C. I., W. L. Baker, et al. (2008). "Antihypertensive medication and their impact on cancer incidence: a mixed treatment comparison meta-analysis of randomized controlled trials." <u>J Hypertens</u> <b>26</b>(4): 622-629.</li> <li>Mauri, D., N. P. Polyzos, et al. (2008). "Multiple-treatments meta-analysis of chemotherapy and targeted therapies in advanced breast cancer." <u>J Natl Cancer Inst</u> <b>100</b>(24): 1780-1791.</li> </ul>
2009	<ul style="list-style-type: none"> <li>Golfiniopoulos, V., G. Pentheroudakis, et al. (2009). "Comparative survival with diverse chemotherapy regimens for cancer of unknown primary site: multiple-treatments meta-analysis." <u>Cancer Treat Rev</u> <b>35</b>(7): 570-573.</li> <li>Hawkins, N., D. A. Scott, et al. (2009). "No study left behind: a network meta-analysis in non-small-cell lung cancer demonstrating the importance of considering all relevant data." <u>Value Health</u> <b>12</b>(6): 996-1003.</li> <li>Lim, E., G. Harris, et al. (2009). "Preoperative versus postoperative chemotherapy in patients with resectable non-small cell lung cancer: systematic review and indirect comparison meta-analysis of randomized trials." <u>J Thorac Oncol</u> <b>4</b>(11): 1380-1388.</li> <li>Mills, E. J., B. Rachlis, et al. (2009). "Metastatic renal cell cancer treatments: an indirect comparison meta-analysis." <u>BMC Cancer</u> <b>9</b>: 34</li> </ul>
2011	<ul style="list-style-type: none"> <li>Bangalore, S., S. Kumar, et al. (2011). "Antihypertensive drugs and risk of cancer: network meta-analyses and trial sequential analyses of 324,168 participants from randomised trials." <u>Lancet Oncol</u> <b>12</b>(1): 65-82.</li> </ul>

	<ul style="list-style-type: none"> <li>• Jansen, J. P. (2011). "Network meta-analysis of survival data with fractional polynomials." <i>BMC Med Res Methodol</i> <b>11</b>: 61.</li> <li>• Leung, H. W. and A. L. Chan (2011). "Multikinase inhibitors in metastatic renal cell carcinoma: indirect comparison meta-analysis." <i>Clin Ther</i> <b>33</b>(6): 708-716.</li> <li>• Nuijten, M. J., R. Aultman, et al. (2011). "An indirect comparison of the efficacy of bevacizumab plus carboplatin and paclitaxel versus pemetrexed with cisplatin in patients with advanced or recurrent non-squamous adenocarcinoma non-small cell lung cancer." <i>Curr Med Res Opin</i> <b>27</b>(11): 2193-2201.</li> <li>• Ziogas, D. C., M. Voulgarelis, et al. (2011). "A network meta-analysis of randomized controlled trials of induction treatments in acute myeloid leukemia in the elderly." <i>Clin Ther</i> <b>33</b>(3): 254-279.</li> </ul>
2012	<ul style="list-style-type: none"> <li>• Alberton, M., P. Wu, et al. (2012). "Adverse events associated with individual statin treatments for cardiovascular disease: an indirect comparison meta-analysis." <i>QJM</i> <b>105</b>(2): 145-157.</li> <li>• Cheng, M. M., B. Goulart, et al. (2012). "A network meta-analysis of therapies for previously untreated chronic lymphocytic leukemia." <i>Cancer Treat Rev</i> <b>38</b>(8): 1004-1011.</li> <li>• Dequen, P., P. Lorigan, et al. (2012). "Systematic review and network meta-analysis of overall survival comparing 3 mg/kg ipilimumab with alternative therapies in the management of pretreated patients with unresectable stage III or IV melanoma." <i>Oncologist</i> <b>17</b>(11): 1376-1385.</li> <li>• Ford, J. A., R. Jones, et al. (2013). "Denosumab for treatment of bone metastases secondary to solid tumours: systematic review and network meta-analysis." <i>Eur J Cancer</i> <b>49</b>(2): 416-430.</li> <li>• Jandhyala, R. and J. Fullarton (2012). "Various formulations of oral transmucosal fentanyl for breakthrough cancer pain: an indirect mixed treatment comparison meta-analysis." <i>BMJ Support Palliat Care</i> <b>2</b>(2): 156-162.</li> <li>• Jansen, J. P. and S. Cope (2012). "Meta-regression models to address heterogeneity and inconsistency in network meta-analysis of survival outcomes." <i>BMC Med Res Methodol</i> <b>12</b>: 152.</li> <li>• Kho, M. E. and M. C. Brouwers (2012). "Application of the systematic review and bibliometric network analysis (SeBriNA) methodology contextualizes evidence. Part 2: rituximab for non-Hodgkin's lymphoma." <i>J Clin Epidemiol</i> <b>65</b>(9): 996-1009.</li> <li>• Mhaskar, R., J. Redzepovic, et al. (2012). "Bisphosphonates in multiple myeloma: a network meta-analysis." <i>Cochrane Database Syst Rev</i>(5): CD003188.</li> </ul>



	<ul style="list-style-type: none"> <li>• Terasawa, T., N. A. Trikalinos, et al. (2013). "Comparative efficacy of first-line therapies for advanced-stage chronic lymphocytic leukemia: a multiple-treatment meta-analysis." <i>Cancer Treat Rev</i> 39(4): 340-349.</li> <li>• Wang, S. Y., H. Chu, et al. (2012). "Network meta-analysis of margin threshold for women with ductal carcinoma in situ." <i>J Natl Cancer Inst</i> 104(7): 507-516.</li> </ul>
2013	<ul style="list-style-type: none"> <li>• Bachelot, T., R. McCool, et al. (2014). "Comparative efficacy of everolimus plus exemestane versus fulvestrant for hormone-receptor-positive advanced breast cancer following progression/recurrence after endocrine therapy: a network meta-analysis." <i>Breast Cancer Res Treat</i> 143(1): 125-133.</li> <li>• Bakalos, G., M. Miligkos, et al. (2013). "Assessing the relative effectiveness and tolerability of treatments in small cell lung cancer: a network meta-analysis." <i>Cancer Epidemiol</i> 37(5): 675-682.</li> <li>• Cope, S., M. J. Ouwens, et al. (2013). "Progression-free survival with fulvestrant 500 mg and alternative endocrine therapies as second-line treatment for advanced breast cancer: a network meta-analysis with parametric survival models." <i>Value Health</i> 16(2): 403-417.</li> <li>• Dranitsaris, G., S. Schmitz, et al. (2013). "Small molecule targeted therapies for the second-line treatment for metastatic renal cell carcinoma: a systematic review and indirect comparison of safety and efficacy." <i>J Cancer Res Clin Oncol</i> 139(11): 1917-1926.</li> <li>• Fang, Y., Y. Ding, et al. (2013). "Radioiodine therapy for patients with differentiated thyroid cancer after thyroidectomy: direct comparison and network meta-analyses." <i>J Endocrinol Invest</i> 36(10): 896-902.</li> <li>• Ford, J. A., R. Jones, et al. (2013). "Denosumab for treatment of bone metastases secondary to solid tumours: systematic review and network meta-analysis." <i>Eur J Cancer</i> 49(2): 416-430.</li> <li>• Gupta, A. K. and M. Paquet (2013). "Network meta-analysis of the outcome 'participant complete clearance' in nonimmunosuppressed participants of eight interventions for actinic keratosis: a follow-up on a Cochrane review." <i>Br J Dermatol</i> 169(2): 250-259.</li> <li>• Palmieri, C., J. R. Fullarton, et al. (2013). "Comparative efficacy of bisphosphonates in metastatic breast and prostate cancer and multiple myeloma: a mixed-treatment meta-analysis." <i>Clin Cancer Res</i> 19(24): 6863-6872.</li> <li>• Pechlivanoglou, P., H. H. Le, et al. (2014). "Mixed treatment comparison of prophylaxis against invasive fungal infections in neutropenic patients receiving</li> </ul>

	<p>therapy for haematological malignancies: a systematic review." <i>J Antimicrob Chemother</i> 69(1): 1-11.</p> <ul style="list-style-type: none"> <li>Robertson, C., A. Close, et al. (2013). "Relative effectiveness of robot-assisted and standard laparoscopic prostatectomy as alternatives to open radical prostatectomy for treatment of localised prostate cancer: a systematic review and mixed treatment comparison meta-analysis." <i>BJU Int</i> 112(6): 798-812.</li> <li>Signorovitch, J., R. Ayyagari, et al. (2014). "Major molecular response during the first year of dasatinib, imatinib or nilotinib treatment for newly diagnosed chronic myeloid leukemia: a network meta-analysis." <i>Cancer Treat Rev</i> 40(2): 285-292.</li> <li>Skoetz, N., S. Trelle, et al. (2013). "Effect of initial treatment strategy on survival of patients with advanced-stage Hodgkin's lymphoma: a systematic review and network meta-analysis." <i>Lancet Oncol</i> 14(10): 943-952.</li> <li>Terasawa, T., N. A. Trikalinos, et al. (2013). "Comparative efficacy of first-line therapies for advanced-stage chronic lymphocytic leukemia: a multiple-treatment meta-analysis." <i>Cancer Treat Rev</i> 39(4): 340-349.</li> <li>Zeppetella, G., A. Davies, et al. (2014). "A network meta-analysis of the efficacy of opioid analgesics for the management of breakthrough cancer pain episodes." <i>J Pain Symptom Manage</i> 47(4): 772-785 e775.</li> </ul>
2014	<ul style="list-style-type: none"> <li>Bachelot, T., R. McCool, et al. (2014). "Comparative efficacy of everolimus plus exemestane versus fulvestrant for hormone-receptor-positive advanced breast cancer following progression/recurrence after endocrine therapy: a network meta-analysis." <i>Breast Cancer Res Treat</i> 143(1): 125-133.</li> <li>Chan, K., K. Shah, et al. (2014). "A Bayesian meta-analysis of multiple treatment comparisons of systemic regimens for advanced pancreatic cancer." <i>PLoS One</i> 9(10): e108749.</li> <li>Chen, Y. P., Z. X. Wang, et al. (2015). "A Bayesian network meta-analysis comparing concurrent chemoradiotherapy followed by adjuvant chemotherapy, concurrent chemoradiotherapy alone and radiotherapy alone in patients with locoregionally advanced nasopharyngeal carcinoma." <i>Ann Oncol</i> 26(1): 205-211.</li> <li>Gresham, G. K., G. A. Wells, et al. (2014). "Chemotherapy regimens for advanced pancreatic cancer: a systematic review and network meta-analysis." <i>BMC Cancer</i> 14: 471.</li> <li>Gupta, T., S. Kannan, et al. (2015). "Concomitant chemoradiotherapy versus altered fractionation radiotherapy in the radiotherapeutic management of</li> </ul>

	<p>locoregionally advanced head and neck squamous cell carcinoma: An adjusted indirect comparison meta-analysis." <i>Head Neck</i> 37(5): 670-676.</p> <ul style="list-style-type: none"> <li>• Hutton, B., L. Joseph, et al. (2014). "Checking whether there is an increased risk of post-transplant lymphoproliferative disorder and other cancers with specific modern immunosuppression regimens in renal transplantation: protocol for a network meta-analysis of randomized and observational studies." <i>Syst Rev</i> 3: 16.</li> <li>• Kelly, M. E., G. Spolverato, et al. (2015). "Synchronous colorectal liver metastasis: a network meta-analysis review comparing classical, combined, and liver-first surgical strategies." <i>J Surg Oncol</i> 111(3): 341-351.</li> <li>• Kumagai, K., I. Rouvelas, et al. (2015). "Survival benefit and additional value of preoperative chemoradiotherapy in resectable gastric and gastro-oesophageal junction cancer: a direct and adjusted indirect comparison meta-analysis." <i>Eur J Surg Oncol</i> 41(3): 282-294.</li> <li>• Ladyzynski, P., M. Molik, et al. (2015). "A network meta-analysis of progression free survival and overall survival in first-line treatment of chronic lymphocytic leukemia." <i>Cancer Treat Rev</i> 41(2): 77-93.</li> <li>• Liang, W., X. Wu, et al. (2014). "Network meta-analysis of erlotinib, gefitinib, afatinib and icotinib in patients with advanced non-small-cell lung cancer harboring EGFR mutations." <i>PLoS One</i> 9(2): e85245.</li> <li>• Mulla, S. M., D. N. Buckley, et al. (2014). "Management of chronic neuropathic pain: a protocol for a multiple treatment comparison meta-analysis of randomised controlled trials." <i>BMJ Open</i> 4(11): e006112.</li> <li>• Nagayama, A., T. Hayashida, et al. (2014). "Comparative effectiveness of neoadjuvant therapy for HER2-positive breast cancer: a network meta-analysis." <i>J Natl Cancer Inst</i> 106(9).</li> <li>• Pechlivanoglou, P., H. H. Le, et al. (2014). "Mixed treatment comparison of prophylaxis against invasive fungal infections in neutropenic patients receiving therapy for haematological malignancies: a systematic review." <i>J Antimicrob Chemother</i> 69(1): 1-11.</li> <li>• Popat, S., A. Mellemaard, et al. (2015). "Nintedanib plus docetaxel as second-line therapy in patients with non-small-cell lung cancer: a network meta-analysis." <i>Future Oncol</i> 11(3): 409-420.</li> <li>• Popat, S., T. Mok, et al. (2014). "Afatinib in the treatment of EGFR mutation-positive NSCLC--a network meta-analysis." <i>Lung Cancer</i> 85(2): 230-238.</li> <li>• Sekine, L., V. D. Morais, et al. (2015). "Conventional and high-dose daunorubicin and idarubicin in acute myeloid leukaemia remission induction treatment: a mixed</li> </ul>
--	--

	<p>treatment comparison meta-analysis of 7258 patients." <i>Hematol Oncol</i> 33(4): 212-219.</p> <ul style="list-style-type: none"> <li>• Signorovitch, J., R. Ayyagari, et al. (2014). "Major molecular response during the first year of dasatinib, imatinib or nilotinib treatment for newly diagnosed chronic myeloid leukemia: a network meta-analysis." <i>Cancer Treat Rev</i> 40(2): 285-292.</li> <li>• Vegter, S. and K. Tolley (2014). "A network meta-analysis of the relative efficacy of treatments for actinic keratosis of the face or scalp in Europe." <i>PLoS One</i> 9(6): e96829.</li> <li>• Wang, J. C., J. H. Tian, et al. (2014). "Which is the best Chinese herb injection based on the FOLFOX regimen for gastric cancer? A network meta- analysis of randomized controlled trials." <i>Asian Pac J Cancer Prev</i> 15(12): 4795-4800.</li> <li>• Xiong, T., R. M. Turner, et al. (2014). "Comparative efficacy and safety of treatments for localised prostate cancer: an application of network meta-analysis." <i>BMJ Open</i> 4(5): e004285.</li> <li>• Zeppetella, G., A. Davies, et al. (2014). "A network meta-analysis of the efficacy of opioid analgesics for the management of breakthrough cancer pain episodes." <i>J Pain Symptom Manage</i> 47(4): 772-785 e775.</li> <li>• Zhang, Y. W., Y. L. Zhang, et al. (2014). "Chemotherapy for patients with gastric cancer after complete resection: a network meta-analysis." <i>World J Gastroenterol</i> 20(2): 584-592.</li> <li>• Zhu, G. Q., K. Q. Shi, et al. (2014). "Systematic review with network meta-analysis: adjuvant therapy for resected biliary tract cancer." <i>Aliment Pharmacol Ther</i> 40(7): 759-770.</li> <li>• Zhu, Z., J. Zhang, et al. (2014). "Efficacy and toxicity of external-beam radiation therapy for localised prostate cancer: a network meta-analysis." <i>Br J Cancer</i> 110(10): 2396-2404.</li> </ul>
--	--

**Supplementary Table 2: The characteristics of the individual RCTs included in network meta-analysis for assessing the relative effectiveness of treatments in SCLC**

	Author	Year Published	Country	Stage	Tx	Demographics			Tumor Response	Survival	Common Hematologic Toxicities (Grade 3 & 4)			
						Randomized Patients (n)	Gender, male:female (%)	Age, mo(s). mean, $\pm$ SD or median, min-max)			Anemia (%)	Leukopenia (%)	Neutropenia (%)	Thrombocytopenia (%)
1	Abratt RP	1991	South Africa	LD	C1	40	65.0 : 35.0	58 (35-71)	75.0	14 (n.a.-n.a.)				
					C2	38	63.2 : 36.8	59 (33-72)	52.6	11 (n.a.-n.a.)				
2	Abratt RP	1995	South Africa	LD	C1	43	65.1 : 34.9	59 (33-72)	74.4	14.5 (n.a.-n.a.)		9.3		
					C3	38	71.1 : 28.9	60 (35-71)	60.5	12 (n.a.-n.a.)		28.9		
3	Altinbas M	2004	Turkey	LD + ED	C4	42	83.3 : 16.7	58 (37-75)	40.5	LD: 10 (7.3 - 12.7), ED: 8 (6.1 - 9.9)			2.4	
				LD + ED	C5	42	78.6 : 21.4	57.5 (34-74)	64.3%	LD: 16 (8.3 - 23.7), ED: 13 (9.8 - 16.2)			11.9	
4	Ansari R	1995	USA	ED	C6	84	67.9 : 32.1	61 (45-76)	67.1	7.3 (n.a.-n.a.)	15.9	47.6		22.0
					C7	87	66.7 : 33.3	63 (32-78)	72.8	9.1 (n.a.-n.a.)	52.5	71.3		35.0
5	Ardizzoni A	2002	Multi-countries	LD + ED	C8	119	70.6 : 29.4	59 (33-69)	79.0	13.5 (11.8 - 15.8)				
					C9	125	70.4 : 29.6	59 (35-70)	88.2	13 (11.3 - 15.3)				

6	Artal-Cortes A	2004	Spain	LD + ED	C6	202	98.5 : 1.5	60 (39-75)	68.8	LD: 12.9 (11.4 - 14.5), ED: 7.9 (7.0 - 9.0)	14.4		57.4	13.9
					C10	200	99.0 : 1.0	63 (35-75)	74.5	LD: 12.9 (11.7 - 14.6), ED: 8.1 (6.8 - 9.5)	14.5		42.5	16.0
7	Baka S	2008	UK	ED	C9	139	48.2 : 51.8	66 (38-81)	69.8	9.7 (n.a.-n.a.)	27.7		91.2	56.2
					C6	141	53.2 : 46.8	65 (39-89)	75.2	10.6 (n.a.-n.a.)	19.3		58.6	49.3
8	Baka S	2010	Greece	ED	C11	183	88.0 : 12.0	63 (35-78)	51.4	10.9 (0.5 - 86.2)	13.1		55.7	19.7
					C12	181	94.5 : 5.5	64 (42-82)	55.2	9.8 (0.5 - 86.1)	11.6		54.7	23.2
9	Bork E	1991	Sweden	LD + ED	C13	46	58.7 : 41.3	74 (70-82)	65.2	8.5 (n.a.-n.a.)				
					C14	48	68.8 : 31.2	73 (70-83)	70.8	11.3 (n.a.-n.a.)				
10	Chahinian AP	1989	USA	LD + ED	C15	86	72.0 : 28.0		51.2	7.9 (n.a.-n.a.)				
					C16	103	68.0 : 32.0		67.0	9.3 (n.a.-n.a.)				
					C17	105	64.0 : 36.0		47.6	7.9 (n.a.-n.a.)				
11	De Marinis	2005	Italy	LD + ED	C18	70	80.0 : 20.0	63 (52-71)	62.9	LD: 1 (11 - 30), ED: 8 (1 - 27)	12.9	12.9	41.4	41.4
					C19	70	88.6 : 11.4	61 (48-75)	57.1	LD: 12 (4 - 28), ED: 9 (1-23)	10.0	4.3	24.3	27.1
12	Eckardt JR	2006	Multi-countries	LD + ED	C20	389	79.4 : 20.6	59.7 ( $\pm$ 9.1)	63.0	9.8 (9.4 - 10.6)	37.9	29.9	58.4	45.9
					C6	395	78.7 : 21.3	59.6 ( $\pm$ 9.6)	68.9	10.1 (9.3 - 10.9)	44.3	44.3	83.6	14.5
13	Ettinger DS	1990	Multi-countries	ED	C3	294	67.3 : 32.7		61.2	10.7 (n.a.-n.a.)				
					C82	283	67.5 : 32.5		64.0	11.5 (n.a.-n.a.)				

14	Ettinger DS	2002	USA	ED	C3	46	54.3 : 45.7	64 (36-79)	56.5	10.5 (n.a.-n.a.)				
					C21	43	60.5 : 39.5	61 (33-79)	48.8	10.8 (n.a.-n.a.)				
					C13	46	65.3 : 21.7	61 (37-79)	43.5	9.5 (n.a.-n.a.)				
15	Fukuoka M	1997	Japan	ED	C22	31	96.8 : 3.2	61 (42-73)	83.9	14.8 (11.4-22.7)				
					C23	32	78.1 : 21.9	61 (44-73)	96.9	8.0 (6.1-10.4)				
16	Furuse K	1998	Japan	ED	C23	114	85.1 : 14.9	64 (32-74)	85.1	11.6 (9.8-12.7)	86.8	83.3		72.8
					C24	113	82.3 : 17.7	64 (38-74)	76.1	10.9 (6.7-12.0)	41.6	82.3		25.7
17	Girling DJ	1996	UK	LD + ED	C25	154	63.0 : 37.0		55.3	4.6 (n.a.-n.a.)				
					C26	156	62.8 : 37.2		53.7	4.7 (n.a.-n.a.)				
18	Greco FA	2005	USA	ED	C27	60	65.0 : 35.0	60 (42-78)	78.3	10.6 (9.5-12.0)	13.3	70.0	81.7	48.3
					C28	60	48.3 : 51.7	62 (38-79)	48.3	9.1 (7.7-10.4)	10.0	60.0	65.0	15.0
19	Grote T	2005	USA	LD + ED	C30	109	54.0 : 45.9	64.4 ( $\pm$ 8.7)	59.6	10.5 (n.a.-n.a.)				
					C6	115	55.7 : 44.3	63.2 ( $\pm$ 8.9)	55.7	10.4 (n.a.-n.a.)				
20	Heigener DF	2009	Germany	ED	C31	37	64.9 : 35.1	60 (46-72)	62.2	11.2 (9.1-15.2)	18.9		70.3	29.7
					C32	40	77.5 : 22.5	61 (41-76)	80.0	11.9 (8.8-14.7)	32.5		37.5	62.5
21	Hirsch FR	1987	Multi-countries	ED	C33	89	73.0 : 27.0	61 ( $\pm$ n.a.)	68.5	5.8 (n.a.-n.a.)				
					C34	92	80.4 : 19.6	60 ( $\pm$ n.a.)	66.3	6.8 (n.a.-n.a.)				
					C35	88	65.9 : 34.1	62 ( $\pm$ n.a.)	73.9	8.3 (n.a.-n.a.)				
22	Hirsch FR	2001	Denmark	LD + ED	C86	136	53.0 : 47.0	59 (34-69)	72.1	10.4 (n.a.-n.a.)				
					C87	134	63.0 : 37.0	59 (40-69)	71.6	9.8 (n.a.-n.a.)				

23	Hong WK	1989	USA	LD + ED	C3	126	81.0 : 19.0	61 (37-79)	54.0	LD: 13.8 (0.5-65.5), ED: 7.8 (0.3-67.8)				
					C36	141	70.0 : 30.0	62 (39-78)	54.6	LD: 13.8 (0.5-89.5), ED: 9.8 (0.3-67.8)				
					C37	86	71.0 : 29.0	62 (43-80)	43.0	LD: 10.3 (0.5-65.5), ED: 7.3 (0.2-80.7)				
24	James LE	1996	UK	ED	C38	89	64.0 : 36.0	63 (38 - 74)	44.9	5.8 (4.0-6.6)		4.5		2.2
					C24	78	65.4 : 34.6	63 (39 - 75)	58.9	6.4 (4.9-7.3)		20.5		2.7
25	Joss RA (Annals of Oncology 6: 41-48, 1995)	1995	Switzerland	ED	C39	32	96.9 : 3.1	59 (n.a. - n.a.)	28.1	4.9 (n.a.-n.a.)				
					C81	27	85.2 : 14.8	56 (n.a. - n.a.)	63.0	8.7 (n.a.-n.a.)				
26	Joss RA (2) (Annals of Oncology 6: 157-166, 1995.)	1995	Switzerland	LD + ED	C81	202	86.6 : 13.4	58 (n.a. - n.a.)	88.4	13.5 (n.a.-n.a.)				
					C89	204	87.7 : 12.3	58 (n.a. - n.a.)	87.4%	11.5 (n.a.-n.a.)				
27	Kanitz E	1992	Multi-countries	ED	C40	52	80.8 : 19.2	55 (33-69)	63.6	6.6 (± n.a.)		19.2		
					C41	59	88.1 : 11.9	56 (33-70)	69.2	27.7 (± n.a.)		8.8		
28	Lassen U	1996	Multi-countries	LD + ED	C84	160	58.7 : 41.3	61 (34-70)	62.5	11.2 (n.a.-n.a.)		53.0		29.0
					C83	158	69.0 : 31.0	62 (36-70)	72.2	11.3 (n.a.-n.a.)		58.0		27.0
					C88	157	59.9 : 40.1	63 (37-70)	64.3	9.8 (n.a.-n.a.)		77.0		10.0
29	Lee SM	2009	UK	LD + ED	C42	121	56.2 : 43.8	62 (37-80)	62.8	8 (n.a.-n.a.)	14.0	31.4	38.8	21.5
					C6	120	56.7 : 43.3	62,5 (27-81)	62.5	8,1 (n.a.-n.a.)	2.5	13.3	28.3	4.2
30	Leyvraz S	2008	Multi-countries	LD + ED	C43	69	75.4 : 24.6		78.3	18.1 (n.a.-n.a.)				
					C44	71	71.8 : 28.2		67.6	14.4 (n.a.-n.a.)				



31	Lorigan P	2005	UK	ED	C44	159	67.3 : 32.7	58 (36-73)	76.1	13.9 (12.9-15.8)	45.8	95.4		82.5
					C43	159	59.1 : 40.9	58 (35-70)	84.3	14.4 (12.7-16.0)	71.5	94.7		95.3
32	Lowenbraun S	1984	USA	ED	C3	106	75.5 : 24.5		71.7	10.5 (n.a.-n.a.)				
					C45	108	75.9 : 24.1		74.1	10.6 (n.a.-n.a.)				
33	Lyss	2002	USA	ED	C20	12	83.3 : 16.7	60.8 ( $\pm$ 8.5)	41.6	5.7 (n.a.-n.a.)				
					C28	13	46.2 : 53.8	64.7 ( $\pm$ 7.9)	53.8	13.8 (n.a.-n.a.)				
					C29	32	53.1 : 46.9	58.0 ( $\pm$ 11.1)	68.8	9.9 (n.a.-n.a.)				
34	Mavroudis D	2001	Greece	LD + ED	C91	62	93.5 : 6.5	62 (36-75)	50.0	LD: 14 (0.5-24.0), ED: 7 (0.5-27.0)	3.2		43.5	17.7
					C6	71	90.1 : 9.9	64 (37-75)	47.9	LD: 12.5 (1.0-25.0), ED: 9.5 (1.0-30.0)	9.8		39.4	5.6
35	Miller AA	1995	USA	ED	C6	156	63.5 : 36.5		53.8	9.5 (8.4-11.0)	32.0	62.0	85.0	32.0
					C46	150	74.7 : 25.3		60.0	9.9 (7.9-11.2)	55.0	83.0	83.0	52.0
36	Milroy R	1993	UK	LD + ED	C47	111	55.9 : 44.1	59 (35-70)	72.1	10.3 (9.0-12.0)				
					C45	109	59.6 : 40.4	59 (37-69)	70.6	11 (9.0-12.3)				
37	Miyomoto H	1992	Japan	LD + ED	C6	42	76.2 : 23.8	63 (40-78)	66.7	13.8 (n.a.-n.a.)	19.0	42.9	21.4	
					C7	47	76.6 : 23.4	63 (37-76)	59.6	14 (n.a.-n.a.)	36.2	72.3	25.5	
38	Monnet I	1992	France	ED	C48	30	96.7 : 3.3	58 ( $\pm$ n.a.)	75.9	9.7 (n.a.-n.a.)				
					C49	30	86.7 : 13.3	57 ( $\pm$ n.a.)	28.6	10.4 (n.a.-n.a.)				
39	Murray N	1999	Multi-countries	ED	C22	110	69.1 : 30.9	59.3 ( $\pm$ n.a.)	87.1	11.8 (n.a.-n.a.)				
					C24	109	63.3 : 36.7	59.3 ( $\pm$ n.a.)	69.7	10.9 (n.a.-n.a.)				

40	Nagel S	2011	Germany	LD + ED	C32	36	77.8 : 22.2	61 (± n.a.)	72.2	11 (6.9-15.0)		11.1		27.8
					C31	36	63.9 : 36.1	59 (± n.a.)	66.7	11 (4.2-17.7)		16.7		25.0
41	Niell HB	2005	USA	ED	C6	282	46.1 : 53.9	62 (n.a.-n.a.)	65.6	9.9 (9.2-10.8)	17.4	34.4	66.7	13.8
					C91	283	45.2 : 54.8	61 (n.a.-n.a.)	73.1	10.6 (9.9-11.2)	19.8	40.3	44.5	22.3
42	Noda K	2002	Japan	ED	C6	77	89.6 : 10.4	63 (41-70)	67.5	9.4 (8.1-10.8)	29.9	51.9	92.2	18.2
					C50	77	81.8 : 18.2	63 (33-70)	84.4	12.8 (11.7-15.2)	26.0	26.0	63.6	5.2
43	Okamoto H	2007	Japan	ED	C31	110	86.4 : 13.6	74 (56-86)	72.7	10.6 (n.a.-n.a.)	29.1	53.6	94.5	55.5
					C6	110	89.1 : 10.9	73.5 (55-85)	73.4	9.9 (n.a.-n.a.)	24.8	51.4	89.9	15.6
44	Postmus PE	1992	Multi-countries	LD + ED	C8	63	77.8 : 22.2	59 (39-70)	83.3	12.3 (n.a.-n.a.)		41.6		16.6
					C51	55	83.6 : 16.4	59 (38-69)	76.5	9.3 (n.a.-n.a.)		52.9		49.0
					C49	60	78.3 : 21.7	57 (39-70)	60.3	10.5 (n.a.-n.a.)		6.8		18.9
45	Postmus PE	1996	Multi-countries	ED	C8	73	84.9 : 15.1	61 (41-73)	68.5	7.6 (n.a.-n.a.)	11.0	90.4		20.5
					C85	70	78.6 : 21.4	61 (29-74)	70.0	8.7 (n.a.-n.a.)	32.9	90.0		62.9
46	Pujol JL	2001	France	ED	C6	109	77.1 : 22.9	59.3 (± 8.9)	60.6	9.3 (n.a.-n.a.)	18.0		85.0	18.0
					C52	117	83.5 : 14.5	58.5 (± 9.2)	76.1	10.5 (n.a.-n.a.)	51.0		99.0	78.0
47	Quoix E	2005	Multi-countries	ED	C20	41	70.7 : 29.3	61 (± n.a.)	63.4	10.4 (1.0-24.3)	46.3	46.3	87.8	31.7
					C53	41	68.3 : 31.7	61 (± n.a.)	61.0	10.9 (0.4-21.9)	19.5	65.9	87.8	19.5
48	Reck M	2003	Germany	LD + ED	C54	307	76.0 : 24.0	60 (35-75)	69.4	11.7 (10.9-12.6)				
					C55	301	75.0 : 25.0	60 (30-75)	72.1	12.7 (11.2-14.1)				

49	Roth BJ	1992	USA	LD + ED	C6	159	77.4 : 22.6	62.7 (± n.a.)	60.7	8.6 (n.a.-n.a.)	25.0	29.3		9.3
					C3	156	80.1 : 19.9	61.7 (± n.a.)	50.7	8.3 (n.a.-n.a.)	12.9	42.9		3.6
					C24	162	75.3 : 24.7	62.6 (± n.a.)	59.4	8.1 (n.a.-n.a.)	26.1	39.9		16.7
50	Rowland KM	1996	USA	LD + ED	C56	122	56.0 : 44.0		68.0					
					C6	121	64.0 : 36.0		79.3					
51	Ruotsalainen TM	1999	Multi-countries	LD + ED	C6	78	59.0 : 41.0	60.5 (± n.a.)	60.3	10.2 (n.a.-n.a.)				
					C57	75	64.0 : 36.0	59.6 (± n.a.)	58.7	10.0 (n.a.-n.a.)				
					C58	66	66.7 : 33.3	59.9 (± n.a.)	60.6	10.1 (n.a.-n.a.)				
52	Schmittel A	2006	Germany	ED	C59	35	71.4 : 28.6	59 (34-77)	62.9		14.3	31.4	25.7	31.4
					C31	35	71.4 : 28.6	63 (48-74)	57.1		34.3	65.7	51.4	31.4
53	Sculier JP	1990	Belgium	LD + ED	C60	95	89.5 : 10.5	61 (37-75)	74.1	11.3 (n.a.-n.a.)		45.0		7.1
					C61	106	90.6 : 9.4	62 (35-74)	55.1	10.0 (n.a.-n.a.)		34.7		7.1
54	Sculier JP	1993	Belgium	LD + ED	C90	107	89.7 : 10.3	61 (35-74)	67.3	12.3 (n.a.-n.a.)		58.9		6.5
					C8	108	89.8 : 10.2	61 (33-74)	61.4	10.8 (n.a.-n.a.)		75.9		16.6
55	Sculier JP	2001	Belgium	ED	C62	78	89.7 : 10.3	61 (37-75)	57.9	9.5 (7.8-11.6)		84.6		15.4
					C63	78	83.3 : 16.7	64 (35-74)	73.6	8.8 (7.3-10.2)		84.6		44.9
56	Seifart U	2007	Germany	ED	C66	51	66.7 : 33.3	60.8 (± n.a.)	78.4	11.8 (n.a.-n.a.)	21.6	27.5		19.6
					C67	49	67.3 : 32.7	63.8 (± n.a.)	73.5	11.6 (n.a.-n.a.)	22.4	20.4		22.4
57	Seifart U	2005	Germany	ED	C64	42	78.6 : 21.4	61.5 (40-72)	38.1	8.7 (n.a.-n.a.)	42.9	64.3		52.4
					C65	42	81.0 : 19.0	60.5 (37-72)	31.0	7.6 (n.a.-n.a.)	21.4	47.6		40.5

58	Sekine I	2003	Japan	ED	C68	30	90.0 : 10.0	64 (47-70)	83.3	8.9 (n.a.-n.a.)	56.7	50.0	56.7	26.7
					C69	30	90.0 : 10.0	63 (46-68)	76.7	12.9 (n.a.-n.a.)	46.7	53.3	86.7	10.0
59	Sekine I	2008	Japan	ED	C50	54	79.6 : 20.4	63 (42-70)	75.9	12.4 (n.a.-n.a.)	33.3	18.5	51.9%	3.7
					C70	55	85.5 : 14.5	62 (48-70)	87.3	13.7 (n.a.-n.a.)	45.5	52.7	94.5%	23.6
60	Slevin ML	1989	UK	ED	C71	20			10.0					
					C14	19			89.5					
61	Socinski MA	2009	Multi-countries	ED	C72	453	71.7 : 28.3	62.5 (35.0-88.5)	24.9	8.1 (n.a.-n.a.)	10.8	4.0	10.2	9.1
					C31	455	72.5 : 27.5	62.5 (38.5 - 86.5)	44.0	10.6 (n.a.-n.a.)	7.3	8.1	46.2	10.1
62	Socinski MA	2006	USA	ED	C72	38	42.1 : 57.9	66 (47-75)	39.5	10.4 (n.a.-n.a.)	11.4		17.1	14.3
					C73	40	50.0 : 50.0	66 (46.82)	35.0	7.6 (n.a.-n.a.)	5.3		18.4	21.1
63	Souhami RL	1994	UK	LD + ED	C74	221	64.3 : 35.7	62 (34-73)	82.4	9.6 (n.a.-n.a.)				
					C24	217	67.7 : 32.3	63 (32-74)	81.1	8.8 (n.a.-n.a.)				
64	Souhami RL	1997	UK	LD + ED	C38	80	52.5 : 47.5	67 (49-80)	46.3	5.9 (n.a.-n.a.)				
					C14	75	56.0 : 44.0	66 (50-86)	32.8	4.8 (n.a.-n.a.)				
65	Steward WP	1998	Multi-countries	LD + ED	C75	153	73.9 : 26.1	59 (38-75)	64.1	11.7 (n.a.-n.a.)				
					C76	147	73.5 : 26.5	60 (37-75)	77.6	14.8 (n.a.-n.a.)				
66	Urban T	1999	France	LD + ED	C8	228	91.7 : 8.3	57 ( $\pm$ 9)	52.2	8.9 (7.9-9.3)				
					C77	229	92.1 : 7.9	56 ( $\pm$ 10)	72.1	9.0 (8.2-10.1)				
67	Wampler GL	1991	USA	ED	C3	79	65.4 : 34.6		43.0					
					C78	82	69.8 : 30.2		48.8					

68	White SC	2001	UK	ED	C3	59	57.6 : 42.4	70 (46-85)	37.9	4.3 (n.a.-n.a.)	8.6	27.6		3.4
					C79	60	48.3 : 51.7	70 (54-76)	25.4	3.9 (n.a.-n.a.)	16.9	25.4		27.1
69	Wolf M	1987	Germany	LD + ED	C6	72	89.0 : 11.0	57 (35-70)	65.3	11.6 (n.a.-n.a.)				
					C80	64	91.0 : 9.0	57 (25-71)	68.8	9.4 (n.a.-n.a.)				
70	Woll PJ	2001	UK	LD + ED	C44	25	80.0 : 20.0	61 (40-69)	76.0	11.8 (n.a.-n.a.)				
					C43	25	56.0 : 44.0	55 (47-68)	80.0	12.7 (n.a.-n.a.)				
71	Zatloukal P	2010	Multi-countries	ED	C50	202	76.2 : 23.8	60 (34-79)	39.1	10.2 (9.0-11.7)	6.9	6.4%	5.4	38.1
					C6	203	76.4 : 23.6	61 (40-75)	46.3	9.7 (8.9-11.1)	6.4	9.9	4.4	59.6

**Supplementary Table 3: Quality assessment of the individual RCTs included in network meta-analysis for assessing the relative effectiveness of treatments in SCLC**

	Author	Year Published	Country	Stage	Precise details of the interventions in each arm	Description of study end-points	Description of sample size estimation	Method of randomization (sequence generation)	Implementation of randomization	Blinding (masking)	Participant flow
1	Abratt RP	1991	South Africa	LD	Yes	Yes	No	Yes	No	n.a.	Yes
2	Abratt RP	1995	South Africa	LD	Yes	Yes	No	Yes	No	n.a.	Yes
3	Altinbas M	2004	Turkey	LD + ED	Yes	Yes	Yes	Yes	No	n.a.	Yes
4	Ansari R	1995	USA	ED	Yes	Yes	No	Yes	No	n.a.	Yes
5	Ardizzoni A	2002	Multi-countries	LD + ED	Yes	Yes	Yes	Yes	No	n.a.	Yes
6	Artal-Cortes A	2004	Spain	LD + ED	Yes	Yes	Yes	Yes	No	n.a.	Yes
7	Baka S	2008	UK	ED	Yes	Yes	Yes	Yes	No	n.a.	Yes
8	Baka S	2010	Greece	ED	Yes	Yes	Yes	No	No	n.a.	Yes
9	Bork E	1991	Sweden	LD + ED	Yes	Yes	Yes	No	No	n.a.	Yes
10	Chahinian AP	1989	USA	LD + ED	Yes	Yes	Yes	No	No	n.a.	Yes
11	De Marinis	2005	Italy	LD + ED	Yes	Yes	Yes	No	No	n.a.	Yes
12	Eckardt JR	2006	Multi-countries	LD + ED	Yes	Yes	Yes	No	No	n.a.	Yes
13	Ettinger DS	1990	Multi-countries	ED	Yes	No	Yes	No	No	n.a.	No
14	Ettinger DS	2002	USA	ED	Yes	Yes	Yes	No	No	n.a.	Yes
15	Fukuoka M	1997	Japan	ED	Yes	Yes	No	No	No	n.a.	Yes
16	Furuse K	1998	Japan	ED	Yes	Yes	Yes	No	No	n.a.	Yes
17	Girling DJ	1996	UK	LD + ED	Yes	Yes	Yes	No	No	n.a.	Yes
18	Greco FA	2005	USA	ED	Yes	Yes	Yes	No	No	n.a.	Yes
19	Grote T	2005	USA	LD + ED	Yes	Yes	Yes	No	No	No	Yes
20	Heigener DF	2009	Germany	ED	Yes	Yes	Yes	No	No	n.a.	Yes
21	Hirsch FR	1987	Multi-countries	ED	Yes	Yes	No	No	No	n.a.	No

22	Hirsch FR	2001	Denmark	LD + ED	Yes	Yes	Yes	No	No	n.a.	Yes
23	Hong WK	1989	USA	LD + ED	Yes	No	Yes	No	No	n.a.	Yes
24	James LE	1996	UK	ED	Yes	No	Yes	Yes	No	n.a.	Yes
25	Joss RA (Annals of Oncology 6: 41-48, 1995)	1995	Switzerland	ED	Yes	No	Yes	No	No	n.a.	Yes
26	Joss RA (2) (Annals of Oncology 6: 157-	1995	Switzerland	LD + ED	Yes	Yes	Yes	Yes	No	n.a.	Yes
27	Kanitz E	1992	Multi-countries	ED	Yes	Yes	No	No	No	n.a.	Yes
28	Lassen U	1996	Multi-countries	LD + ED	Yes	No	No	No	No	n.a.	Yes
29	Lee SM	2009	UK	LD + ED	Yes	Yes	Yes	Yes	No	n.a.	Yes
30	Leyvraz S	2008	Multi-countries	LD + ED	Yes	Yes	Yes	No	No	n.a.	Yes
31	Lorigan P	2005	UK	ED	Yes	Yes	Yes	Yes	No	n.a.	Yes
32	Lowenbraun S	1984	USA	ED	Yes	No	Yes	No	No	n.a.	Yes
33	Lyss	2002	USA	ED	Yes	Yes	Yes	Yes	Yes	n.a.	Yes
34	Mavroudis D	2001									
35	Miller AA	1995	USA	ED	Yes	Yes	Yes	Yes	No	n.a.	Yes
36	Milroy R	1993	UK	LD + ED	Yes	No	Yes	Yes	No	n.a.	Yes
37	Miyomoto H	1992	Japan	LD + ED	Yes	No	No	No	No	n.a.	Yes
38	Monnet I	1992	France	ED	Yes	No	Yes	No	No	n.a.	Yes
39	Murray N	1999	Multi-countries	ED	Yes	Yes	Yes	No	No	n.a.	Yes
40	Nagel S	2011	Germany	LD + ED	Yes	Yes	Yes	Yes	Yes	n.a.	Yes
41	Niell HB	2005	USA	ED	Yes	Yes	No	No	No	n.a.	Yes
42	Noda K	2002	Japan	ED	Yes	Yes	Yes	No	No	n.a.	Yes
43	Okamoto H	2007	Japan	ED	Yes	Yes	Yes	No	No	n.a.	Yes
44	Postmus PE	1992	Multi-countries	LD + ED	Yes	No	No	No	No	n.a.	Yes
45	Postmus PE	1996	Multi-countries	ED	Yes	No	Yes	Yes	No	n.a.	Yes

46	Pujol JL	2001	France	ED	Yes	No	Yes	Yes	No	n.a.	Yes
47	Quoix E	2005	Multi-countries	ED	Yes	Yes	Yes	Yes	No	n.a.	Yes
48	Reck M	2003	Germany	LD + ED	Yes	Yes	Yes	Yes	No	n.a.	Yes
49	Roth BJ	1992	USA	LD + ED	Yes	Yes	Yes	No	No	n.a.	Yes
50	Rowland KM	1996	USA	LD + ED	Yes	No	Yes	No	No	n.a.	Yes
51	Ruotsalainen TM	1999	Multi-countries	LD + ED	Yes	Yes	Yes	No	No	n.a.	Yes
52	Schmittel A	2006	Germany	ED	Yes	Yes	Yes	Yes	No	n.a.	Yes
53	Sculier JP	1990	Belgium	LD + ED	Yes	No	No	No	No	n.a.	No
54	Sculier JP	1993	Belgium	LD + ED	Yes	No	No	No	No	n.a.	No
55	Sculier JP	2001	Belgium	ED	Yes	No	No	No	No	n.a.	No
56	Seifart U	2007	Germany	ED	Yes	Yes	Yes	Yes	No	n.a.	Yes
57	Seifart U	2005	Germany	ED	Yes	Yes	Yes	Yes	No	n.a.	Yes
58	Sekine I	2003	Japan	ED	Yes	Yes	Yes	Yes	No	n.a.	Yes
59	Sekine I	2008	Japan	ED	Yes	No	No	Yes	No	n.a.	Yes
60	Slevin ML	1989	UK	ED	Yes	No	No	No	No	n.a.	Yes
61	Socinski MA	2009	Multi-countries	ED	Yes	Yes	Yes	Yes	No	n.a.	Yes
62	Socinski MA	2006	USA	ED	Yes	Yes	Yes	Yes	No	n.a.	Yes
63	Souhami RL	1994	UK	LD + ED	Yes	Yes	Yes	Yes	No	n.a.	Yes
64	Souhami RL	1997	UK	LD + ED	Yes	Yes	Yes	Yes	No	n.a.	Yes
65	Steward WP	1998	Multi-countries	LD + ED	Yes	Yes	Yes	No	No	n.a.	Yes
66	Urban T	1999	France	LD + ED	Yes	Yes	Yes	Yes	No	n.a.	Yes
67	Wampler GL	1991	USA	ED	Yes	No	No	No	No	n.a.	Yes
68	White SC	2001	UK	ED	Yes	Yes	Yes	Yes	No	n.a.	Yes
69	Wolf M	1987	Germany	LD + ED	Yes	No	No	No	No	n.a.	Yes



<b>70</b>	<b>Woll PJ</b>	<b>2001</b>	<b>UK</b>	<b>LD + ED</b>	<b>Yes</b>	<b>No</b>	<b>No</b>	<b>No</b>	<b>No</b>	<b>n.a.</b>	<b>Yes</b>
<b>71</b>	<b>Zatloukal P</b>	<b>2010</b>	<b>Multi- countries</b>	<b>ED</b>	<b>Yes</b>	<b>Yes</b>	<b>Yes</b>	<b>Yes</b>	<b>No</b>	<b>n.a.</b>	<b>Yes</b>

**Supplementary Table 4: Proportion of reporting of 24 data items in a total of 81 randomized clinical trials in small cell lung cancer by publication period (pre- and post-CONSORT and combined)\***

Data items	Combined 1984-2011 (n = 71) <sup>†</sup>	Pre- CONSORT 1984-1995 (n = 24)	Post- CONSORT 1996-2011 (n = 47)	ΔPost- CONSORT – Pre- CONSORT	OR, 95% CI <sup>¥</sup>	P-value <sup>‡</sup> FET Two- tailed
<b>TITLE/ABSTRACT</b>						
1. Randomized in title/abstract	0.83 (59)	0.83 (20)	0.83 (39)	0.95 (19)	0.9750 (0.2616 – 3.6342)	1.0000
<b>INTRODUCTION</b>						
2. Scientific background in introduction	0.89 (63)	0.83 (20)	0.91 (23)	0.65 (13)	2.150 (0.4875 – 9.4826)	0.4296
<b>METHODS</b>						
3. Eligibility criteria for participants	1.00 (71)	1.00 (24)	1.00 (47)	0.54 (13)	NaN	1.0000
4. Precise details of the interventions in each arm	1.00 (71)	1.00 (24)	1.00 (47)	0.54 (13)	NaN	1.0000
5. Objectives	0.92 (65)	0.88 (21)	0.94 (44)	0.62 (13)	2.095 (0.389 – 11.272)	0.3992
6. End-points	0.68 (48)	0.46 (11)	0.79 (37)	2.36 (26)	4.373 (1.508 – 12.676)	0.0075
7. Sample size	0.76 (54)	0.54 (13)	0.93 (41)	2.15 (28)	5.782 (1.787 – 18.708)	0.0033
8. Method of randomization (sequence generation)	0.45 (32)	0.29 (7)	0.53 (25)	2.57 (18)	0.760 (0.965 – 7.888)	0.0778
9. Allocation concealment	0.04 (3)	0.04 (1)	0.04 (2)	1.0 (1)	1.022 (0.088 – 11.876)	1.0000

<b>10. Implementation of randomization</b>	0.01 (1)	0.00 (0)	0.02 (1)	n.a. (1)	0	1.0000
<b>11. Blinding (masking)</b>	0.00 (0)	n.a.	0.00 (0/2)	n.a. (0)	n.a.	n.a.
<b>12. Statistical methods</b>	0.90 (64)	0.79 (19)	0.96 (45)	1.36 (26)	5.921 (1.055 – 33.242)	0.0396
<b>RESULTS</b>						
<b>13. Participant flow</b>	0.93 (66)	0.83 (20)	0.99 (46)	1.30 (26)	9.200 (0.966 – 87.580)	0.0416
<b>14. Periods: a. Recruitment</b>	0.61 (43)	0.33 (8)	0.74 (35)	3.38 (27)	5.833 (1.996 – 17.048)	0.0017
<b>b. Follow-up</b>	0.01 (1)	0.00 (0)	0.02 (1)	n.a. (1)	0	1.0000
<b>15. Baseline data</b>	0.97 (69)	1.00 (24)	0.96 (45)	0.92 (22)	Infinity	0.5461
<b>16. "Intention-to-treat" analysis</b>	0.82 (58)	0.67 (16)	0.89 (42)	1.63 (26)	4.200 (1.195 – 14.763)	0.0264
<b>17. a. Outcomes and</b>	0.69 (49)	0.63 (15)	0.72 (34)	1.27 (19)	1.569 (0.552 – 4.461)	0.2597
<b>b. Estimation of effects</b>	0.32 (23)	0.13 (3)	0.43 (20)	5.66 (17)	5.185 (1.357 – 19.819)	0.0150
<b>18. Ancillary analyses</b>	0.54 (38)	0.33 (8)	0.64 (30)	2.75 (22)	3.529 (1.252 – 9.951)	0.0231
<b>19. Adverse events</b>	0.77 (55)	0.63 (15)	0.85 (40)	1.66 (25)	3.429 (1.083 – 10.8532)	0.0397
<b>DISCUSSION</b>						
<b>20. Interpretation of the results</b>	0.80 (57)	0.83 (20)	0.79 (37)	0.85 (17)	0.740 (0.206 – 2.664)	0.7594
<b>21. Generalizability</b>	0.76 (54)	0.63 (15)	0.83 (39)	1.60 (24)	2.925 (0.951 – 8.994)	0.0784
<b>22. Overall evidence</b>	0.68 (48)	0.50 (12)	0.77 (36)	2.00 (24)	3.2727 (1.1487 – 9.3241)	0.0327

\* CONSORT = Consolidated Standards of Reporting Trials.

† The percentage of articles reporting the CONSORT item.

¥ Odds ratio of reporting an item at post-CONSORT period relative to pre-CONSORT.

‡ *P*-values from Fisher's exact test for testing the association between reporting an item and publication period.

**Supplementary Table 5: Definition of treatments**

No	Description of regimen
<b>C1</b>	Doxorubicin + Etoposide + Vincristine (intensified)
<b>C2</b>	Doxorubicin + Etoposide + Vincristine
<b>C3</b>	Cyclophosphamide + Doxorubicin + Vincristine
<b>C4</b>	Cyclophosphamide + Epirubicin + Vincristine
<b>C5</b>	Cyclophosphamide + Epirubicin + Vincristine + Low Molecular Weight Heparin
<b>C6</b>	Cisplatin + Etoposide
<b>C7</b>	Cisplatin + Etoposide + Ifosfamide
<b>C8</b>	Cyclophosphamide + Doxorubicin + Etoposide
<b>C9</b>	Cyclophosphamide + Doxorubicin + Etoposide (intensified)
<b>C10</b>	Cisplatin + Epirubicin
<b>C11</b>	Cisplatin + Etoposide / Topotecan (sequential adm.: 4 cycles of EP followed by 4 cycles of T)
<b>C12</b>	Topotecan \ Cisplatin + Etoposide (alternate adm.: alternating cycles of EP and T)
<b>C13</b>	Teniposide
<b>C14</b>	Etoposide
<b>C15</b>	CCNU + Cyclophosphamide + Doxorubicin + Methotrexate
<b>C16</b>	CCNU + Cyclophosphamide + Doxorubicin + Methotrexate + Warfarin
<b>C17</b>	Cisplatin+ Etoposide + Hexamethylamine + Mitomycin / CCNU + Cyclophosphamide + Doxorubicin + Methotrexate (alternating administration)
<b>C18</b>	Cisplatin + Etoposide + Gemcitabin
<b>C19</b>	Cisplatin + Gemcitabin
<b>C20</b>	Cisplatin + Topotecan
<b>C21</b>	Ifosfamide + MESNA
<b>C22</b>	Cisplatin + Doxorubicin + Etoposide + Vincristine
<b>C23</b>	Cisplatin + Doxorubicin + Etoposide + Vincristine (intensified)
<b>C24</b>	Cisplatin + Etoposide / Cyclophosphamide + Doxorubicin + Vincristine ; (alt. administration)
<b>C25</b>	Cyclophosphamide + Etoposide + Methotrexate+ Vincristine
<b>C26</b>	Etoposide + Vincristine
<b>C27</b>	Etoposide + Carboplatin+ Paclitaxel
<b>C28</b>	Paclitaxel + Topotecan
<b>C29</b>	Paclitaxel + Topotecan (intensified)
<b>C30</b>	Cisplatin + Etoposide (intensified)
<b>C31</b>	Carboplatin + Etoposide
<b>C32</b>	Carboplatin + Etoposide (intensified)

<b>C33</b>	CCNU + Cyclophosphamide + Vincristine + Methotrexate
<b>C34</b>	CCNU + Cyclophosphamide + Vincristine + Etoposide (delayed administration of etoposide)
<b>C35</b>	CCNU + Cyclophosphamide + Vincristine + Etoposide
<b>C36</b>	Cyclophosphamide + Etoposide + Vincristine
<b>C37</b>	Cyclophosphamide + Vincristine
<b>C38</b>	Cisplatin + Etoposide / Cyclophosphamide + Doxorubicin + Vincristine (intensified)
<b>C39</b>	Carboplatin + Teniposide
<b>C40</b>	Cisplatin + Doxorubicin
<b>C41</b>	Doxorubicin + Cyclophosphamide
<b>C42</b>	Carboplatin + Gemcitabin
<b>C43</b>	Carboplatin + Etoposide + Ifosfamide + Mesna (intensified)
<b>C44</b>	Carboplatin + Etoposide + Ifosfamide + Mesna
<b>C45</b>	Cyclophosphamide + Doxorubicin + Etoposide + Vincristine
<b>C46</b>	Cisplatin + Etoposide (per os administration)
<b>C47</b>	Cyclophosphamide + Doxorubicin + Etoposide + Vincristine + Verapamil
<b>C48</b>	Cisplatin + Doxorubicin + Etoposide
<b>C49</b>	Carboplatin + Ifosfamide + Vincristine
<b>C50</b>	Cisplatin + Irinotecan
<b>C51</b>	Carboplatin + Ifosfamide
<b>C52</b>	Cisplatin + Cyclophosphamide + Etoposide + Epirubicin
<b>C53</b>	Cisplatin + Topotecan(intensified iii)
<b>C54</b>	Carboplatin + Etoposide + Vincristine
<b>C55</b>	Carboplatin + Etoposide + Paclitaxel
<b>C56</b>	Cisplatin + Etoposide + Megestrol acetate
<b>C57</b>	Cisplatin + Etoposide + nIFNA-a
<b>C58</b>	Cisplatin + Etoposide + rIFNA-a
<b>C59</b>	Carboplatin + Irinotecan
<b>C60</b>	Cisplatin + Etoposide + Vindesine
<b>C61</b>	Etoposide + Vindesine
<b>C62</b>	Epirubicin + Ifosfamide + Vindesine
<b>C63</b>	Epirubicin + Ifosfamide + Vindesine (intensified)
<b>C64</b>	Cisplatin + Topotecan (intensified i)
<b>C65</b>	Cisplatin + Topotecan (intensified ii)
<b>C66</b>	Carboplatin + Topotecan (intensified)
<b>C67</b>	Carboplatin + Topotecan
<b>C68</b>	Cisplatin + Etoposide + Irinotecan (intensified i)
<b>C69</b>	Cisplatin + Etoposide + Irinotecan (intensified ii)

<b>C70</b>	Cisplatin + Etoposide + Irinotecan
<b>C71</b>	Etoposide (intensified)
<b>C72</b>	Carboplatin + Pemetrexeb
<b>C73</b>	Cisplatin + Pemetrexeb
<b>C74</b>	Cisplatin + Etoposide / Doxorubicin + Ifosfamide; (alternating cycles)
<b>C75</b>	Carboplatin + Etoposide + Ifosfamide + Vincristine
<b>C76</b>	Carboplatin + Etoposide + Ifosfamide + Vincristine (intensified)
<b>C77</b>	Cisplatin + Cyclophosphamide + Doxorubicin + Etoposide
<b>C78</b>	Cyclophosphamide + Doxorubicin + Vincristine / Cisplatin + Etoposide + Methotrexate (alternated every 3 weeks)
<b>C79</b>	Cisplatin
<b>C80</b>	Etoposide + Ifosfamide
<b>C81</b>	Doxorubicin+ Cisplatin+ Etoposide /CCNU + Cyclophosphamide + Methotrexate + Vincristine
<b>C82</b>	Cyclophosphamide + Doxorubicin + Vincristine / Etoposide+ Hexamethylamine + Methotrexate; (alt. administration)
<b>C83</b>	Carboplatin + Teniposide + Vincristine for cycles 1-3, 7,11 / Cyclophosphamide + CCNU + Vincristine + Etoposide for cycles 4,8 / Doxorubicin + Vincristine for cycles 5,9 / Cisplatin + Hexamethylmelamine + Vindesine for cycles 6,10
<b>C84</b>	Cisplatin + Teniposide + Vincristine for cycles 1-3, 7,11 / Cyclophosphamide + CCNU + Vincristine + Etoposide for cycles 4,8 / Doxorubicin + Vincristine for cycles 5,9 / Cisplatin + Hexamethylmelamine + Vindesine for cycles 6,10
<b>C85</b>	Cyclophosphamide + Doxorubicin + Etoposide for cycles 1,3,5 / Carboplatin + Ifosfamide (+ Vincristine for cycles 2,4
<b>C86</b>	Carboplatin + Cisplatin + Teniposide for cycles 1-3, 6 / Cyclophosphamide + Messnafor cycle 4 / Epirubicin for cycle 5
<b>C87</b>	Carboplatin + Teniposide for cycles 1-3, 6 / Cyclophosphamide + Messna for cycle 4 / Epirubicin for cycle 5
<b>C88</b>	Cyclophosphamide+ CCNU + Vincristine + Etoposide for cycles 1,4,7,10/ Doxorubicin + Vincristine for cycles 2,5,8,11/ Cisplatin + Hexamethylmelamine + Vindesine for cycles 3,6,9
<b>C89</b>	Doxorubicin+ Cisplatin + Etoposide for cycles 1-3 / CCNU + Cyclophosphamide + Methotrexate + Vincristine for cycles 4-6
<b>C90</b>	Doxorubicin + Cisplatin + Cyclophosphamide + Etoposide + Methotrexate + Vincristine + Vindesine
<b>C91</b>	Cisplatin + Etoposide + Paclitaxel

**Supplementary Table 6: Results of all direct and indirect comparisons**

Type of analysis (D=Direct, I=Indirect)	Treatment	Comparator	OR	95LL	95UL
<b>Patients with objective response rate</b>					
D	C56	C6	0.55422	0.309755	0.991621
D	C3	C6	0.665814	0.414409	1.06974
D	C46	C6	1.28571	0.816817	2.02377
D	C50	C6	0.593876	0.132406	2.66369
D	C30	C6	1.1772	0.692344	2.00161
D	C31	C6	1.27801	0.510157	3.20158
D	C9	C6	2.35832	0.093697	59.3584
D	C52	C6	3.38354	0.033455	342.2
D	C20	C6	0.459808	0.023609	8.95511
D	C42	C6	1.2746	0.455121	3.56961
D	C80	C6	1.14231	0.542099	2.40707
D	C57	C6	0.890407	0.404427	1.96037
D	C91	C6	1.08538	0.543949	2.16574
D	C24	C6	0.947479	0.586228	1.53134
D	C7	C6	1.05174	0.599475	1.84521
D	C58	C6	0.974952	0.481172	1.97545
D	C10	C6	0.983117	0.402191	2.40313
I	C13	C6	0.347879	0.213037	0.568071
I	C79	C6	0.333012	0.180685	0.613757
I	C23	C6	3.7901	1.76812	8.1244
I	C21	C6	0.429696	0.26345	0.700848
I	C14	C6	0.403549	0.237971	0.684336
I	C71	C6	0.005622	6.93E-05	0.456306
I	C2	C6	0.398636	0.169074	0.939889
I	C22	C6	1.59698	0.652116	3.9109
I	C48	C6	5.90883	0.13127	265.972
I	C51	C6	10.1892	0.048849	2125.34
I	C32	C6	1.66141	0.425502	6.48715
I	C27	C6	4.26485	0.079789	227.963
I	C77	C6	2.79199	0.100352	77.6788
I	C3	C6	0.48124	0.03984	5.81309
I	C73	C6	0.258308	0.001081	61.7478
I	C72	C6	0.302168	0.001273	71.7514



I	C70	C6	3.50033	0.004328	2830.98
I	C45	C6	0.587723	0.034084	10.1344
I	C82	C6	0.584416	0.023647	14.4434
I	C38	C6	0.74922	0.129755	4.3261
I	C57	C6	0.898971	0.440411	1.83499
I	C90	C6	1.5339	0.053603	43.8936
I	C47	C6	0.655205	0.017476	24.5648
I	C78	C6	0.653166	0.008836	48.281
I	C59	C6	0.826347	0.085383	7.99746
I	C85	C6	1.269	0.043263	37.2225
I	C28	C6	1.40845	0.004804	412.893
I	C8	C6	1.1823	0.043473	32.1541
I	C58	C6	0.965664	0.435719	2.14016
I	C29	C6	0.754956	0.000155	3681.22
I	C49	C6	0.743288	6.27E-05	8817.25
I	C74	C6	1.30706	1.55E-07	1.10E+07
I	C24	C6	1.33116	4.97E-12	3.57E+11
I	C1	C6	0.979463	1.99E-05	48127.3
<b>Patients with complete response</b>					
D	C7	C6	2.27129	0.967299	5.33316
D	C52	C6	1.84394	0.902652	3.76681
D	C9	C6	0.63962	0.359906	1.13672
D	C56	C6	1.30813	0.718025	2.38321
D	C3	C6	0.692308	0.296584	1.61604
D	C24	C6	0.703125	0.301135	1.64174
D	C20	C6	1.22318	0.672741	2.22399
D	C80	C6	0.500818	0.033487	7.48997
D	C57	C6	1.41566	0.348882	5.74433
D	C58	C6	1.8024	0.153905	21.1081
D	C50	C6	0.65623	0.108926	3.95348
D	C91	C6	0.716851	0.169476	3.03215
D	C10	C6	1.35999	0.320729	5.76678
D	C42	C6	1.16473	0.52872	2.56581
D	C46	C6	0.937063	0.488282	1.79832
D	C30	C6	1.00589	0.510862	1.9806
D	C31	C6	0.990476	0.278457	3.52314
I	C71	C6	0.125615	0.024157	0.653186
I	C14	C6	0.355232	0.142912	0.882991
I	C51	C6	0.308476	0.108566	0.876492

I	C37	C6	0.423695	0.17917	1.00194
I	C49	C6	0.370171	0.133258	1.02828
I	C13	C6	0.421213	0.1653	1.07333
I	C2	C6	0.39996	0.139544	1.14637
I	C48	C6	4.73667	0.456798	49.1159
I	C77	C6	1.79612	0.695196	4.64047
I	C3	C6	0.416939	0.080079	2.17083
I	C32	C6	1.91264	0.304896	11.9982
I	C58	C6	1.82484	0.325135	10.242
I	C72	C6	0.330886	0.010552	10.3761
I	C8	C6	0.77119	0.335864	1.77076
I	C45	C6	0.484484	0.02923	8.03039
I	C36	C6	0.586134	0.059741	5.75067
I	C73	C6	0.314546	0.001671	59.2029
I	C29	C6	0.478655	0.015518	14.7645
I	C90	C6	1.23117	0.381965	3.96835
I	C85	C6	1.21462	0.359969	4.09845
I	C23	C6	1.90383	0.032101	112.912
I	C27	C6	1.67256	0.029925	93.4827
I	C57	C6	1.39825	0.097942	19.9619
I	C21	C6	0.707792	0.037851	13.2352
I	C79	C6	1.29821	0.09855	17.1015
I	C53	C6	1.22318	0.069404	21.5575
I	C28	C6	0.799539	0.003736	171.118
I	C38	C6	1.18658	0.013187	106.768
I	C82	C6	0.602298	5.63E-09	6.44E+07
I	C1	C6	0.914652	0.011098	75.3794
I	C22	C6	1.06126	0.024267	46.4125
I	C74	C6	1.43536	1.42E-10	1.45E+10
I	C24	C6	1.50243	5.26E-14	4.29E+13
I	C78	C6	0.659352	2.53E-31	1.72E+30
I	C70	C6	1.00443	0.10781	9.35794

**Supplementary Table 7: Common grade 3-4 hematological toxicities described by the included RCTs in the network meta-analysis, according to the grouping of chemotherapy treatments**

Anemia			Leukopenia			Neutropenia			Febrile neutropenia			Thrombocytopenia		
Treat ment	Sta ge	No of pati ents (%)	Treat ment	Sta ge	No of patie nts (%)	Treat ment	Sta ge	No of patie nts (%)	Treat ment	Sta ge	No of patie nts (%)	Treat ment	Sta ge	No of patie nts (%)
C68	ED	17 (56.7 )	C51	LD , ED	51 (100. 0)	C51	LD , ED	50 (98.0 )	C7	ED	42 (52.5 )	C52	ED	90 (76.9 )
C52	ED	59 (50.4 )	C62	ED	66 (84.6 )	C52	ED	113 (96.6 )	C6	ED	31 (37.8 )	C9	ED	77 (56.2 )
C69	ED	14 (46.7 )	C63	ED	66 (84.6 )	C70	ED	52 (94.5 )	C70	ED	17 (30.9 )	C64	ED	22 (52.4 )
C7	LD , ED	59 (46.5 )	C85	ED	56 (80.0 )	C9	ED	125 (91.2 )	C59	ED	4 (11.4 )	C27	ED	29 (48.3 )
C70	ED	25 (45.5 )	C7	LD , ED	91 (71.7 )	C53	ED	36 (87.8 )	C50	ED	5 (9.3)	C32	LD , ED	35 (46.1 )
C64	ED	18 (42.9 )	C27	ED	42 (70.0 )	C69	ED	26 (86.7 )	C31	ED	26 (5.3)	C63	ED	35 (44.9 )
C20	LD , ED	161 (38.7 )	C8	LD , ED	162 (67.2 )	C27	ED	49 (81.7 )	C72	ED	7 (1.4)	C20	LD , ED	185 (44.5 )
C32	ED	13 (32.5 )	C53	ED	27 (65.9 )	C6	LD , ED	1155 (66.6 )				C18	LD , ED	29 (41.4 )
C85	ED	20 (28.6 )	C64	ED	27 (64.3 )	C28	ED	39 (65.0 )				C85	ED	29 (41.4 )
C9	ED	38 (27.7 )	C28	ED	36 (60.0 )	C20	LD , ED	255 (61.3 )				C65	ED	17 (40.5 )
C24	LD , ED	36 (26.1 )	C90	LD , ED	63 (58.9 )	C68	ED	17 (56.7 )				C7	ED	28 (35.0 )
C67	ED	11 (22.4 )	C69	ED	16 (53.3 )	C31	ED	358 (56.2 )				C59	ED	11 (31.4 )
C66	ED	11 (21.6 )	C70	ED	29 (52.7 )	C11	ED	102 (55.7 )				C19	LD , ED	19 (27.1 )
C6	LD , ED	422 (21.5 )	C68	ED	15 (50.0 )	C12	ED	99 (54.7 )				C79	ED	16 (27.1 )
C65	ED	9 (21.4 )	C65	ED	20 (47.6 )	C50	ED	154 (46.2 )				C68	ED	8 (26.7 )
C53	ED	8 (19.5 )	C60	LD , ED	38 (44.7 )	C91	LD , ED	153 (44.3 )				C70	ED	13 (23.6 )
C79	ED	10 (16.9 )	C91	ED	114 (40.3 )	C10	LD , ED	85 (42.5 )				C12	ED	42 (23.2 )
C91	LD , ED	58 (16.8 )	C3	LD , ED	87 (36.9 )	C18	LD , ED	29 (41.4 )				C67	ED	11 (22.4 )
C50	ED	52 (15.6 )	C61	LD , ED	34 (34.7 )	C42	LD , ED	47 (38.8 )				C42	LD , ED	26 (21.5 )
C10	LD	29	C6	LD	495	C32	ED	15				C91	LD	74

	, ED	(14.5 )
C59	ED	5 (14.3 )
C42	LD , ED	17 (14.0 )
C27	ED	8 (13.3 )
C31	ED	84 (13.2 )
C11	ED	24 (13.1 )
C18	LD , ED	9 (12.9 )
C3	LD , ED	23 (11.6 )
C12	ED	21 (11.6 )
C72	ED	53 (10.8 )
C19	LD , ED	7 (10.0 )
C28	ED	6 (10.0 )
C8	ED	7 (9.6)
C73	ED	2 (5.0)

	, ED	(34.5 )
C24	LD , ED	70 (33.2 )
C20	LD , ED	131 (31.5 )
C59	ED	11 (31.4 )
C42	LD , ED	38 (31.4 )
C66	ED	14 (27.5 )
C79	ED	15 (25.4 )
C67	ED	10 (20.4 )
C31	LD , ED	125 (19.7 )
C50	ED	43 (12.9 )
C18	LD , ED	9 (12.9 )
C32	LD , ED	4 (11.1 )
C49	LD , ED	6 (10.3 )
C1	LD	4 (9.3)
C38	ED	4 (4.5)
C19	LD , ED	3 (4.3)
C72	ED	18 (4.0)

		(37.5 )
C49	LD , ED	20 (34.5 )
C8	LD , ED	17 (28.3 )
C59	ED	9 (25.7 )
C7	LD , ED	12 (25.5 )
C19	LD , ED	17 (24.3 )
C73	ED	7 (17.5 )
C5	LD , ED	5 (11.9 )
C72	ED	52 (10.6 )
C4	LD , ED	1 (2.4)


	, ED	(21.4 )
C31	LD , ED	138 (20.5 )
C73	ED	8 (20.0 )
C11	ED	36 (19.7 )
C66	ED	10 (19.6 )
C53	ED	8 (19.5 )
C24	LD , ED	23 (16.7 )
C62	ED	12 (15.4 )
C6	LD , ED	263 (15.4 )
C28	ED	9 (15.0 )
C8	LD , ED	23 (12.7 )
C69	ED	3 (10.0 )
C72	ED	46 (9.4)
C61	LD , ED	7 (7.1)
C60	LD , ED	6 (7.1)
C90	LD , ED	7 (6.5)
C50	ED	17 (5.1)
C3	LD , ED	7 (3.5)
C24	ED	2 (2.7)
C38	ED	2 (2.2)

**Supplementary Table 8: Additional studies identified after the initial literature research**


Author, Year	Description	Journal	Citation	DOI	Pub. Date	Registry
<b>Belani 2016</b>	Cisplatin + Etoposide, Cisplatin + Etoposide + Vismodegib, Cisplatin + Etoposide + Cixutumumab	Cancer	Cancer. 2016 Aug 1;122(15):2371-8	10.1002/cncr.30062	Aug-16	NCT00887159
<b>Fink 2012</b>	Cisplatin + Etoposide, Cisplatin + Topotecan	Journal of Thoracic Oncology	J Thorac Oncol. 2012 Sep;7(9):1432-9.	10.1097/JTO.0b013e318260de75	Sep-12	NCT00320359
<b>Langer 2014</b>	Carboplatin + Etoposide, Carboplatin + Etoposide + aBcl2	Lung Cancer	Lung Cancer. 2014 Sep;85(3):420-8	10.1016/j.lungcan.2014.05.003	Sep-14	NCT00682981
<b>Lu 2015</b>	Carboplatin + Etoposide, Carboplatin + Etoposide + Endostatin	Journal of Thoracic Oncology	J Thorac Oncol. 2015;10: 206–211)	10.1097/JTO.0000000000000343	Jan-15	NCT00912392
<b>O'Brien 2011</b>	Cisplatin + Etoposide, Amrubicin + Cisplatin, Amrubicin	European Journal of Cancer	Eur J Cancer. 2011 Oct;47(15):2322-30.	10.1016/j.ejca.2011.05.020	Oct-11	NCT00388960
<b>Oh 2016</b>	Cisplatin + Etoposide, Cisplatin + Belotecan	BMC Cancer	BMC Cancer (2016) 16:690	10.1186/s12885-016-2741-z	Aug-16	NCT00826644
<b>Owonikoko, 2014</b>	Cisplatin + Etoposide + Topotecan, Cisplatin + Etoposide + Irinotecan	Cancer Chemother Pharmacol	Cancer Chemother Pharmacol. 2014 Jan;73(1):171-80	10.1007/s00280-013-2338-z	Jan-14	NCT00057837
<b>Reck 2013</b>	Carboplatin + Paclitaxel, Carboplatin + Paclitaxel + Ipilimumab	Annals of Oncology	Annals of Oncology 24: 75–83, 2013	10.1093/annonc/mds213	Aug-12	NCT00527735
<b>Satoushi 2014</b>	Cisplatin + Irinotecan, Cisplatin + Amrubicin	Journal of Clinical Oncology	J Clin Oncol. 2014 Apr 20;32(12):1262-8	10.1200/JCO.2013.53.5153	Apr-14	UMIN000000720

<b>Sekine 2014</b>	Carboplatin + Etoposide Amrubicin	Clinical Lung Cancer	Clin Lung Cancer. 2014 Mar;15(2):96-102.	10.1016/j.clc.2013.11.006	Mar-15	NCT00286169
<b>Shi 2015</b>	Cisplatin + Etoposide, Cisplatin + Irinotecan	Thoracic Cancer	Thoracic Cancer 6 (2015) 785–791	10.1111/1759-7714.12303	Jul-15	NCT02323737
<b>Spiegel 2011*</b>	Platin + Etoposide, Platin + Etoposide + Bevacizumab	Journal of Clinical Oncology	J Clin Oncol 29:2215- 2222		Jun-11	NCT00403403
<b>Sun 2016</b>	Cisplatin + Etoposide, Cisplatin + Amrubicin	BMC Cancer	BMC Cancer (2016) 16:265	10.1186/s12885-016-2301-6	Apr-16	NCT00660504
<b>Tiseo 2016</b>	Cisplatin + Etoposide, Cisplatin + Etoposide + Bevacizumab	1) Clinical Lung Cancer, 2) ASCO 2016	1) Clin Lung Cancer. 2015 Jan;16(1):67-70, 2) Journal of Clinical Oncology, 2016 ASCO Annual Meeting (June 3- 7, 2016). Vol 34, No 15_suppl (May 20 Supplement), 2016: 8513	1) 10.1016/j.clc.2014.09.001	Jun-16	2007-007949-13.

**Supplementary Figure 1: Network of treatments for the outcome “patients with complete response”.**




**Supplementary Figure 2: Network of treatments for the outcome “patients with neutropenia”.**





**Supplementary Figure 3: Network of treatments for the outcome “patients with febrile neutropenia”.**



## Manuscripts

*Bakalos G, Miligos M, Doxani C et al. Assessing the relative effectiveness and tolerability of treatments in small cell lung cancer: a network meta-analysis. Cancer Epidemiol. 2013 Oct;37(5):675-82*



## Assessing the relative effectiveness and tolerability of treatments in small cell lung cancer: A network meta-analysis

Georgios Bakalos<sup>a</sup>, Michael Miligkos<sup>b</sup>, Chrysoula Doxani<sup>a,c</sup>, Ioanna Mpoulmari<sup>a</sup>, Paraskevi Rodopoulou<sup>a</sup>, Elias Zintzaras<sup>a,d,\*</sup>

<sup>a</sup> Evidence-Based Medicine Unit, Department of Biomathematics, University of Thessaly, School of Medicine, Larissa, Greece

<sup>b</sup> Clinical and Translational Science Program, Sackler School of Graduate Biomedical Sciences, Tufts University, Boston, MA, USA

<sup>c</sup> Division of Hematology, University Hospital of Larissa, Biopolis, Larissa, Greece

<sup>d</sup> Center for Clinical Evidence Synthesis, The Institute for Clinical Research and Health Policy Studies, Tufts Medical Center, Tufts University School of Medicine, Boston, MA, USA

### ARTICLE INFO

#### Article history:

Received 30 May 2012

Received in revised form 27 June 2013

Accepted 28 June 2013

Available online 2 August 2013

#### Keywords:

Small cell lung cancer

Network

Indirect analysis

Randomized control trials

Chemotherapy

### ABSTRACT

**Background:** The combination of Cisplatin plus Etoposide (EP) is currently the standard treatment for small cell lung cancer (SCLC). However, a large number of alternative treatments (monotherapies and combinations) have been studied in randomized controlled trials (RCTs) to identify more effective treatments. **Aim of the present study** was to assess the relative effectiveness and tolerability of these treatments. **Methods:** PubMed, EMBASE and Cochrane Central Register of Controlled Trials were systematically searched to identify all RCTs that compared treatments for SCLC. Then, effectiveness of the treatments relative to the combination of Cisplatin plus Etoposide, reference treatment) was estimated by performing a network of treatments analysis. The analysis evaluated two efficacy outcomes (complete response – CR and objective response rate – ORR) and two tolerability outcomes (neutropenia and febrile neutropenia). All RCTs that provided data for calculating the odds ratios (OR) for the selected outcomes were considered. The network analysis involved direct and indirect analyses. **Results:** We identified 71 articles eligible for inclusion, involving 91 different treatments. In total, 16,026 patients were included in the analysis. In the direct analysis the combination of Cisplatin plus Cyclophosphamide plus Etoposide plus Epirubicin showed better response than EP for the ORR outcome, but with worse tolerability (presence of neutropenia). The indirect analysis revealed that the combination of Cisplatin plus Doxorubicin plus Etoposide (plus Vincristine) showed better response than EP for the ORR outcome. **Conclusions:** No therapy shows better response for the two efficacy outcomes (CR and ORR); though, Cisplatin plus Doxorubicin plus Etoposide plus Vincristine might be a promising therapy for SCLC. The results should be interpreted with caution because the network was dominated by indirect comparisons. Large scale head-to-head RCTs are needed to confirm the present findings.

© 2013 Elsevier Ltd. All rights reserved.

### 1. Introduction

Small cell lung cancer (SCLC) accounts for about 15% of all lung cancers [1] and is characterized by a rapid tumor growth rate and early dissemination to regional lymph nodes and to distant sites [2]. At the time of diagnosis, one third of patients diagnosed with SCLC have tumors confined to the hemithorax of origin, the mediastinum, or the supraclavicular lymph nodes (limited-stage disease, LD) and the remaining patients have tumors spread beyond the supraclavicular areas (extensive-stage disease, ED) [3].

Patients with SCLC typically develop distant metastases and thus, localized forms of treatment (e.g. surgical resection or radiation therapy) may not be effective [4]. Thus, chemotherapy remains the standard treatment of SCLC. The most used agents in SCLC are alkylating agents (cisplatin, carboplatin, ifosfamide and cyclophosphamide), antimitotic agents (vincristine and paclitaxel) and topoisomerase inhibitors (etoposide, irinotecan, topotecan and doxorubicin). In both ED and LD SCLC, the combination of cisplatin and etoposide remains the most widely used standard chemotherapeutic regimen [5]. However, the selection of the optimal chemotherapy agent or combination of chemotherapy agents is a difficult task since no studies have estimated the relative effectiveness and safety of all alternative treatments [6,7]. Thus, an integration of the current evidence and quantification of the relative effectiveness and safety of all these treatments based on published RCTs are needed.

\* Corresponding author at: Center for Clinical Evidence Synthesis, The Institute for Clinical Research and Health Policy Studies, Tufts Medical Center, 800 Washington Street, #63, Boston, MA 02111, USA. Tel.: +1 617 6361692.  
E-mail address: [ezintzaras@tuftsmedicalcenter.org](mailto:ezintzaras@tuftsmedicalcenter.org) (E. Zintzaras).

In order to evaluate the relative merits of the different treatments for SCLC based on the mode of action of each chemotherapy agent (or combination of individual chemotherapy agents), we systematically searched and cataloged all available published RCTs in SCLC. Then, we performed a network of multiple treatments analysis (network meta-analysis), involving direct analysis (synthesis of RCTs with the same treatment comparisons) and indirect analysis (comparison between treatments using an intermediate comparator) [8,9]. In the absence of direct comparison between treatments, the effect size can only be estimated only using an indirect comparison approach [8,9]. A network of treatments can be constructed by considering all investigated comparisons (direct and indirect) between treatments. The aim of the network meta-analysis is to synthesize all evidence originated from direct and indirect comparisons and to assess the effectiveness and tolerability of treatments using as reference treatment a standard first line treatment (such as the combination of cisplatin and etoposide). The present methodology has already been applied in ranking the relative effectiveness of treatments in acute myeloid leukemia [8] and multiple sclerosis [9].

## 2. Materials and methods

### 2.1. Search strategy-selection of RCTs

We searched PubMed, EMBASE, and the Central Registry of Controlled Trials of the Cochrane Library to identify all RCTs that investigated chemotherapy regimens in adult patients with histologically proven SCLC. The search was limited to English language, RCTs, adults, and concerned the time period from 1980 until end of May 2011. The articles were identified using as search criterion the terms: “small cell lung cancer” and “chemotherapy”. The reference lists of the retrieved articles were also reviewed to identify additional publications. The search strategy for the selection of the eligible RCTs is shown in Fig. 1.

### 2.2. Eligibility criteria

RCTs that compared at least two arms of different chemotherapy regimens in chemotherapy naïve patients with histologically proven SCLC were included in the network analysis. Only studies that provided sufficient data to calculate odds ratios (ORs) for estimating the magnitude of difference between treatments, and the corresponding precision were considered.

The following studies were excluded: (i) studies comparing second line chemotherapy treatments; (ii) studies reporting radiotherapy interventions, i.e. radical radiotherapy in combination with chemotherapy or chemotherapy administration for sensitization to radiation; (iii) studies reporting surgical interventions; (iv) studies reporting adjuvant chemotherapy (i.e. chemotherapy following radical surgical intervention) or neo-adjuvant chemotherapy (i.e. chemotherapy prior to radical surgical interventions); (v) studies reporting supportive care interventions or comparison of chemotherapy with chemotherapy plus conventional supportive care and (iv) follow-up and extension studies. In addition, studies with a crossover design, meeting abstracts and conference proceedings were excluded.

In RCTs involving more than two treatment arms, each pairwise treatment comparison was considered as different study. Also, RCTs providing data for different SCLC stages were considered as separate studies in the analysis. In order to avoid the inclusion of duplicated data, the retrieved studies were appraised by geographic location, author names and period of study. Then, in studies with overlapping patients, the largest one

was included in the analysis. Only studies conducted after approval from national ethical committees were considered.

### 2.3. Data extraction and outcomes definition

The following information was extracted from each eligible article: name of first author, year of publication, country of origin, reported stage of SCLC, sample size (randomized patients, totally and per arm), types and intensity (dose and duration) of chemotherapies, effect size of each outcome of interest and chemotherapy regimen. Data extraction was undertaken by 2 investigators (GB and CD), independently. The overall agreement rate was 89%. Any disagreement was resolved by a third independent investigator (EZ).

Two primary outcomes were considered for the network analysis: the CR and the ORR. Complete Response (CR) is achieved when all tumor lesions are disappeared after treatment initiation. Objective Response Rate (ORR) is the portion of patients with a predefined amount of tumor size reduction; ORR is defined as the sum of CR and partial response and it is a direct measure of drug antitumor activity. Among the many adverse events after treatment with chemotherapy, we chose to record the neutropenia (NP) and febrile neutropenia (FNP) because they are considered the most important ones.

### 2.4. Treatment definition

Chemotherapy regimens containing the same chemotherapy agents, irrespective of dosage scheme and maximum duration of each chemotherapy cycle, were defined as the same treatment since we are interested in the assessment of the relative effectiveness of the different agent-based therapies. In addition, the effect of different dosage schemes and chemotherapy cycle intensity remains unresolved [5]. Furthermore, the current grouping allows the definition of a less complicated and analyzable network. The combination of cisplatin and etoposide (EP) was set as the reference treatment in the subsequent treatment comparisons since it is the standard first line treatment and the most commonly investigated chemotherapy regimen.

### 2.5. Statistical methods

Treatments were compared using odds ratios (ORs) with their respective 95% confidence intervals (CI). When more than two studies compared the same treatments, a random effects (RE) pooled OR was calculated [10]. The RE model incorporates the between study variability and it is more conservative than the fixed effects model [11].

Indirect comparison was performed for treatments not compared directly [12]. Then, in comparing two treatments, A and B, where each treatment was compared directly with treatment C, the OR for comparing A and B was calculated using the following principle [8]:  $\ln(OR_{AvsB}) = \ln(OR_{AvsC}) - \ln(OR_{BvsC})$ , and the respective 95% CI was estimated assuming asymptotic normality and lack of covariance [12–16] (Fig. 2). The network of treatments was constructed based on all investigated comparisons between treatments and the indirect analysis was performed utilizing all the possible pathways provided by the network. The OR was considered significant when the 95% CI included the one (1).

The network graph was built using S-PLUS 8 (Seattle, WA, USA, <http://www.insightful.com>) [17] and the network analysis was carried out using NET-MS (<http://netms.med.uth.gr>) [8,9]. The algorithm was implemented using Compaq Visual Fortran90 with the IMSL library (Hewlett Packard, Avondale, PA) [18]. MetaAnalyst (Evidence-Based Practice Center, Tufts Medical Center, Boston, MA,




Fig. 1. Flow diagram of the screening process and RCTs selection for multiple-treatments meta-analysis of treatments for SCLC.




Fig. 2. Direct evidence comes from the synthesis of trials of A versus B. Indirect evidence comes through an intermediate/common comparator C (many intermediate comparators are possible) by combining trials of A versus C and of C versus B (prior to combination the trials were synthesized).



USA, [http://tuftscaes.org/meta\\_analyst](http://tuftscaes.org/meta_analyst)) [19] was used to validate the findings of data syntheses.

### 3. Results

#### 3.1. Eligible studies and summary characteristics

The literature search in PubMed, EMBASE and Cochrane Central Register of Controlled Trials identified 243 articles that met the search criterion. After title selection and abstract reviewing of the articles, 98 articles from all databases were judged to be potentially relevant and they were reviewed in their entirety. Finally, 71 articles were selected for inclusion the network analysis: 71 articles for the outcome CR, 69 for ORR, 23 for NP and 5 for FNP. The articles were published between 1980 and 2011. A flow diagram of included/excluded articles is shown in Fig. 1.

The summary characteristics of included RCTs in the multiple treatments meta-analysis is shown in Table 1. The characteristics of the individual RCTs including efficacy/tolerability results and their quality assessment are shown in Supplementary Table 1 and the definition of the treatments is given in Supplementary Table 2. In total, the eligible studies involved 16,026 randomly assigned patients with SCLC and the majority of them were male (72%) while the median age was 61 (55–74) years. Most of the articles involved studies carried out in the US (21.1%) and 18.3% of them were multicenter trials, involving several countries. 83.7% (63/71) of the studies included two comparing treatment arms and only 11.3% of them compared more than two arms. More than half of the studies (37/71) included patients with extended disease (ED), while two studies included patients with limited disease (LD) and 32 patients with ED and LD. The median sample size was 230 (12–455)

patients. Overall survival was reported in 67 articles. Overall, the median ORR was 65.2% (10.0–96.9%) while the median overall survival was 10.3 (1.0–27.7) months. Adverse events of grade 3–4 were reported for 58.5% (9371) of the patients. Almost half patients experienced grade 3–4 neutropenia (53.1%) and leucopenia (44.3%). Thrombocytopenia and anemia was reported in 22.0% and 19.8% of the patients, respectively.

In assessing the quality of reporting, seven items were considered: (1) precise details of the interventions in each arm; (2) description of study end-points; (3) description of sample size estimation; (4) method of randomization (sequence generation); (5) implementation of randomization; (6) blinding and (7) participant flow. The majority of studies were open-label and only two studies were blinded. The precise details of the interventions in each arm were reported in all studies, while the study end-points and the sample size estimation were reported in 50 (70.4%) and 54 (76%) studies respectively. Despite the fact that the method of randomization was described in 32 (45%) reports, only two reports provided information about the implementation of randomization (2.8%). The participant flowchart was described in 66 studies (93%).

#### 3.2. The networks

For the outcome CR, we identified 91 different treatments, representing 4095 theoretically possible direct comparisons, but only 90 (1.1%) had been performed in the retrieved RCTs. The 90 direct comparisons were utilized in network analysis, i.e. in performing both the direct and indirect analyses. The geometry of the network of comparisons for CR is depicted in Fig. 3; the other outcomes are represented in Supplementary Figures 1–3.

In the network figure, the size of the circles was directly related to the number of RCTs investigated each treatment, while the thickness of connecting lines was directly related to the number of available direct comparisons. More specifically, common treatments [e.g. EP] that were compared by more RCTs were drawn with larger circles whereas infrequently investigated regimens (e.g. Cisplatin plus Doxorubicin) were represented by smaller circles. However, most of the treatments were compared against EP, which represented the most commonly used treatment in the RCTs (26 direct comparisons). Carboplatin plus Etoposide involved the biggest sample size of randomized patients (455 patients). All regimens are listed in Supplementary Table 2.


#### 3.3. Direct analysis for comparing treatments with EP

Sixteen treatments were compared directly with EP in 18 trials [20–37]: (1) Cyclophosphamide plus Doxorubicin (CAV); (2) Cisplatin plus Etoposide plus Ifosfamide (VIP); (3) Cyclophosphamide plus Doxorubicin plus Etoposide plus GCSF [ACE (intensified)]; (4) Cisplatin plus Epirubicin (PEP); (5) Cisplatin plus Topotecan (TC); (6) Cisplatin plus Etoposide/Cyclophosphamide plus Doxorubicin plus Vincristine (CAV/EP); (7) Cisplatin plus Etoposide plus GCSF [EP (intensified)]; (8) Carboplatin (AUC5) plus Etoposide (EC); (9) Carboplatin (AUC5) plus Gemcitabine (GEM-CAR); (10) Cisplatin plus Irinotecan (IP); (11) Cisplatin plus Cyclophosphamide plus Etoposide plus Epirubicin (CCEE); (12) Cisplatin plus Etoposide plus Megestrol acetate (EP + Ma); (13) Cisplatin plus Etoposide plus natural interferon alpha (EP + nIFN- $\alpha$ ); (14) Cisplatin plus Etoposide plus recombinant interferon alpha (EP + rIFN- $\alpha$ ); (15) Etoposide plus Ifosfamide (IE) and (16) Cisplatin plus Etoposide plus Paclitaxel (PET).

The numbers of direct comparisons with EP for the outcomes CR, ORR, NP and FNP were 18, 17, 9 and 1, respectively. None of the treatments showed better response compared to EP for both efficacy outcomes. The significant results derived from the direct

**Table 1**  
Characteristics of 71 included randomized controlled trials (RCTs) in the network of treatments meta-analysis. Data are given as number (percentage) except where indicated otherwise.

Characteristics	RCTs included
(n = 71)	
No. of eligible patients	16,026
Median sample size	230
Year of publication	
1981–1990	8 (11.3%)
1991–2000	29 (40.8%)
2001–2011	34 (47.9%)
Number of eligible arms	
Two	63 (88.7%)
Three	8 (11.3%)
Outcomes extracted	
Tumor response	
Patients with complete response	71 (100%)
Patients with objective response	71 (100%)
Safety	
Patients with adverse events	71 (100.0)
Response	
No. of patients with complete response	2720 (16.9%)
No. of patients with objective response	10,201 (63.7%)
Countries involved (investigator affiliations)	
Multiple countries	13 (18.3%)
United States	15 (21.1%)
United Kingdom	11 (15.5%)
Germany	7 (9.9%)
Japan	7 (9.9%)
Belgium	3 (4.2%)
France	3 (4.2%)
Other countries	12 (16.9%)



**Fig. 3.** Network of direct comparisons for the outcome “patients with complete response”. Thickness of connecting lines is proportional to the number of available direct comparisons. The size of each node is proportional to the number of trials investigated each treatment. The definition of the treatments is given in Supplementary Table 2.

analysis are shown in Table 2; the treatments are sorted according to their effect sizes. The results of all direct comparisons are shown in Supplementary Table 3.

### 3.3.1. Patients with complete response

No regimen was significantly different than EP ( $P \geq 0.05$ ).

### 3.3.2. Patients with objective response

The treatment combination Cisplatin plus Cyclophosphamide plus Etoposide plus Epirubicin (CCEE) produced better response [OR = 2.07 (1.17–3.67)], whereas Cisplatin plus Etoposide plus Megestrol acetate (EP + Ma) derived worse response [OR = 0.55 (0.31–0.99)].

### 3.3.3. Tolerability

Two treatment combinations produced worse tolerability (in terms of NP) than EP (Cyclophosphamide plus Doxorubicin plus Etoposide plus GCSF and Cisplatin plus Cyclophosphamide plus Etoposide plus Epirubicin) and two treatments shown better tolerability ( $P < 0.01$ ) (Table 2). Regarding FNP, only one treatment (Cisplatin plus Etoposide plus Ifosfamide) was compared directly to EP, producing non-significant result [OR = 1.81 (0.97–3.40)] [20].

### 3.4. Indirect analysis for comparing treatments with EP

Table 3 shows the indirect analysis significant results ( $P < 0.05$ ); the treatments are sorted according to their effect size.

**Table 2**

Direct analysis results for comparing treatments for small cell lung cancer (SCLC) with reference treatment (Cisplatin + Etoposide, EP) by outcome. The treatments were sorted according to their significance and magnitude of effect size.

Treatment	Stage	Patients, No.	OR (95% CI)	P-value
<i>Patients with objective response</i>				
Cisplatin + Cyclophosphamide + Etoposide + Epirubicin	ED	226	2.07 (1.17–3.67)	0.01
Cisplatin + Etoposide + Megestrol acetate	LD, ED	243	0.55 (0.31–0.99)	0.05
<i>Patients with neutropenia</i>				
Cyclophosphamide + Doxorubicin + Etoposide + GCSF	ED	280	7.37 (3.72–14.56)	<0.01
Cisplatin + Cyclophosphamide + Etoposide + Epirubicin	ED	226	5.59 (1.83–17.1)	<0.01
Cisplatin + Epirubicin	LD, ED	402	0.54 (0.37–0.81)	<0.01
Cisplatin + Topotecan	LD, ED	784	0.27 (0.2–0.39)	<0.01

ED, extensive disease; LD, limited disease.



**Table 3**  
Indirect analysis results for comparing treatments for small cell lung cancer (SCLC) with reference treatment (Cisplatin + Etoposide, EP) by outcome, for treatments that produced significantly ( $P < 0.05$ ) different response than reference treatment. The treatments were sorted according to their significance and magnitude of effect size.

Treatment	OR (95% CI)	P-value
<i>Patients with complete response</i>		
Etoposide	0.36 (0.14–0.88)	0.03
Carboplatin + Ifosfamide	0.31 (0.11–0.88)	0.03
Etoposide (intensified)	0.13 (0.02–0.65)	0.01
<i>Patients with objective response</i>		
Cisplatin + Doxorubicin + Etoposide + Vincristine (intensified)	3.79 (1.77–8.12)	<0.01
Ifosfamide + Mesna	0.43 (0.26–0.70)	<0.01
Carboplatin + Pemetrexeb	0.41 (0.21–0.79)	<0.01
Etoposide	0.40 (0.24–0.68)	<0.01
Doxorubicin + Etoposide + Vincristine	0.40 (0.17–0.94)	0.04
Cyclophosphamide + Doxorubicin + Etoposide	0.38 (0.16–0.93)	0.03
Teniposide	0.35 (0.21–0.57)	<0.01
Cisplatin	0.33 (0.18–0.61)	<0.01
Carboplatin + Ifosfamide	0.25 (0.066–0.94)	0.04
Etoposide (intensified)	0.006 (0.00–0.46)	0.02
<i>Patients with neutropenia</i>		
Carboplatin + Pemetrexeb	0.26 (0.09–0.76)	0.01

None of the treatments derived a better response than EP for both efficacy outcomes. However, one treatment [Cisplatin plus Doxorubicin plus Etoposide plus Vincristine (intensified)] showed better response for the outcome ORR but, this treatment showed worse tolerability in the direct analysis. The results of all indirect comparisons are shown in Supplementary Table 3.

#### 3.4.1. Patients with complete response

None treatment showed better outcome than EP ( $P \geq 0.05$ ). However, the analysis indicated that monotherapy with etoposide (either standard or intensified) and combination therapy with Carboplatin plus Ifosfamide have less comparative effectiveness [OR = 0.36 (0.14–0.88), OR = 0.13 (0.02–0.65) and OR = 0.31 (0.11–0.88), respectively].

#### 3.4.2. Patients with objective response

Only one treatment combination yielded better response: Cisplatin plus Doxorubicin plus Etoposide plus Vincristine (intensified) [OR = 3.79 (1.77–8.12)]. However, nine treatments revealed worse response: (i) Ifosfamide plus Mesna; (ii) Carboplatin plus Pemetrexeb; (iii) Etoposide; (iv) Doxorubicin plus Etoposide plus Vincristine; (v) Cyclophosphamide plus Doxorubicin plus Etoposide; (vi) Teniposide; (vii) Cisplatin; (viii) Carboplatin plus Ifosfamide and (ix) Etoposide (intensified) (Table 3).

#### 3.4.3. Tolerability

Only the treatment Carboplatin plus Pemetrexeb indicated a better tolerability for the outcome NP [OR = 0.26 (0.09–0.76)]. For the outcome FNP, there is only one study [20] reporting this outcome (see direct analysis section) and thus, the comparative tolerability of treatments was not evaluated further.

### 4. Discussion

Herein, we present a comprehensive and systematic assessment of the current status of treating SCLC. The primary aim of the present study was to provide an assessment of the relative effectiveness of treatments in SCLC, especially in the absence of head-to-head comparisons, and to direct future research in SCLC treatment. In order to achieve this scope, we carried out a network analysis of all published RCTs in SCLC. The network analysis

involved the following steps: direct comparison of treatments, indirect comparison and combination of direct and indirect comparison. The secondary aim was to reveal the necessity to performing large RCTs for head-to-head comparisons of treatments. There are no studies involved more than 500 patients and the various chemotherapy combinations have not compared to a standard treatment such as EP. The network consisted of 91 treatments, involving 18 direct comparisons for the outcome CR, 17 for the outcome ORR and 10 for the tolerability.

The analysis of the network indicated that only two regimens have shown effectiveness compared to EP: the application of network analysis of treatments makes optimal use of all available published data and provides insight in the relative effectiveness of different treatments (monotherapies and combination therapies) [38]. However, the selection of the optimal treatment is a difficult task and network analysis may assist in quantifying the rank order of treatments in terms of efficacy/tolerability and outcomes. The direct and indirect analyses revealed two treatments with better effectiveness compared to the reference treatment (EP) for the outcome ORR: (1) combination of Cisplatin, Cyclophosphamide, Etoposide and Epirubicin and (2) combination of Cisplatin, Doxorubicin and Etoposide with Vincristine (intensified), respectively. But, the former combination showed worst tolerability than EP.

On the contrary, six other regimens showed worse effectiveness for the ORR outcome (Ifosfamide plus Mesna, Carboplatin plus Pemetrexeb, Doxorubicin plus Etoposide plus Vincristine, Cyclophosphamide plus Doxorubicin plus Etoposide, Teniposide and Cisplatin) and three regimens for the both outcomes (ORR and CR) (Etoposide standard, Etoposide intensified and Carboplatin plus Ifosfamide).

In the present study, the differences of the dosage schemes and/or treatment cycle maximum duration were ignored since we focused to the antitumor activity of each treatment based on the mode of action of each chemotherapy agent (or combination of individual chemotherapy agents). We adopted this approach since the scientific evidence of the relative anti-tumor activity of each chemotherapy agent, or combination of individual chemotherapy agents is relative scarce.

In the network analysis, possible effect modifiers were not taken into account and only the unadjusted pooled ORs were calculated since data that affect the response were not provided in the individual studies. In addition, the estimated effect sizes were unadjusted for treatment dosage levels. Nevertheless, the developed methodology (and of course, the NET-MS system) cannot estimate adjusted effect sizes; though, it has the capability of subgroup analyses. In addition, the existence of publication bias (defining as the differential magnitude of effect in large versus small studies) cannot totally be excluded [39]. However, a valid method for testing publication bias in network analysis does not exist. Also, in the network analysis, adjustments for multiple comparisons may not be applicable since the purpose of the analysis was to explore the relative significance of risk effect [40]. Data were synthesized with an objective (to assess the relative effectiveness of treatments) but not with a prespecified key hypothesis [40–42]. An appropriate multiple test adjustment is difficult or even impossible because the investigated comparisons in the network are not independent and a clear structure in the multiple tests is missing [42]. Finally, the existence of false positive results may not be totally excluded since heterogeneity between studies within the network cannot be assessed (lack of valid methodology) and the network analysis cannot adjust for possible effect modifiers; though, synthesis of data from many studies usually is expected to reduce false discovery rate. In reporting the network analysis we did not adopt the PRISMA statement [43] since the present approach is



relative novel and it is not considered a typical meta-analysis of RCTs; though, some of the items of the statement are reported adequately. Although the quality of reporting of the studies included in the network-analysis was assessed, a sensitivity analysis involving the studies with high reporting quality was not considered since the aim of the assessment was to obtain an indication of the reporting quality of the current evidence in SCLC treatment; in addition, there is no established quality scales to divide “high-quality” from “low-quality” studies. Furthermore, it has been shown that individual quality measures are not associated with treatment effect size across studies and medical areas [44].

In the network, many treatments were compared in a small number of RCTs and could not be linked to the EP pathway, thus they could not be compared to the reference treatment. In addition, there was not enough replication of treatment comparisons. Therefore, we adopted an adjusted indirect analysis method and not a mixed effects logit hierarchical model nor a complicated Bayesian approach [38,45–47]. Also, the analysis was not restricted to specific subpopulations (e.g. limited and extensive-stage SCLC) due to lack of replication and to achieve greater power in detecting significant results. Since the indirect comparisons are not randomized but observational studies across trials the differences in study populations and prognostic factors across RCTs may lead to overestimation of the treatment effects [48,49]. In addition, the network analysis was based on grouped data from published RCTs and not on individual patient data, assuming that the relative effectiveness of a treatment is consistent in different RCTs. Therefore, the results regarding the superiority of a particular treatment should be interpreted with great caution. However, when the previous basic assumption may not be met, the results of one RCT can be not generalizable to another; though, the identification of factors that may influence the generalizability of an RCT is rather difficult [50].

In general, the network meta-analysis can be useful when there is no direct evidence of the relative effectiveness of treatments and the direct evidence is not sufficient [51]. The results from the direct and indirect analyses may be combined to provide more reliable estimates of the effect sizes when there is no discrepancy between the two analyses [52], which is not the case in the present analysis, and therefore, the combined analysis was not considered. Although, the indirect analysis provides more evidence (since utilizes more information from the network) than the direct, the results from the direct analysis are always more reliable in drawing inferences (since it is based on randomization). Considering that indirect comparisons dominate the network and that there is variability of SCLC patients in terms of demographic and clinical characteristics, the generalized decision about the choice of treatment in SCLC patients should be considered with caution and the network results cannot be extrapolated beyond them.

In conclusion, none therapy shows better response for the two efficacy outcomes (CR and ORR); though, Cisplatin plus Doxorubicin plus Etoposide plus Vincristine might be a promising therapy for SCLC. However, large scale head-to-head RCTs are needed to confirm the present findings.

#### Conflict of interest

The authors declare that they have no competing interests.

#### Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.canep.2013.06.008>.

#### References

- [1] Jemal A, Siegel R, Ward E, Hao Y, Xu J, Murray T, et al. Cancer statistics, 2008. *CA Cancer J Clin* 2008;58(2):71–96.
- [2] Masters G. The clinical presentation of small cell lung cancer. In: Pass HI, Carbone DP, Johnson DH, et al., eds. Lung cancer: principles and practice. Philadelphia: Lippincott Williams & Wilkins, 2005: 304–14.
- [3] Murray N, Coy P, Patier JL, Hodson I, Arnold A, Zee BC, et al. Importance of timing for thoracic irradiation in the combined modality treatment of limited-stage small-cell lung cancer. The National Cancer Institute of Canada Clinical Trials Group. *J Clin Oncol* 1993;11(2):336–44.
- [4] Prasad US, Naylor AR, Walker WS, Lamb D, Cameron EW, Walbaum PR. Long term survival after pulmonary resection for small cell carcinoma of the lung. *Thorax* 1989;44(10):784–7.
- [5] Sorensen M, Pijls-Johannasma M, Felip E, on behalf of the ESMO Guidelines Working Group. Small-cell-lung cancer: ESMO clinical practice guidelines for diagnosis, treatment and follow up. *Ann Oncol* 2010;21(Suppl 5):v120–5.
- [6] Comis RL, Friedland DM, Good BC. Small-cell lung cancer: a perspective on the past and a preview of the future. *Oncology (Williston Park)* 1998 Jan;12(1 Suppl 2):44–50.
- [7] Agra Y, Pelayo M, Sacristan M, Sacristán A, Serra C, Bonfill X. Chemotherapy versus best supportive care for extensive small cell lung cancer. *CDS Rev* 2003;4(4):CD001990.
- [8] Zogas DC, Voulgaris M, Zintzaras E. A network meta-analysis of randomized controlled trials of induction treatments in acute myeloid leukemia in the elderly. *Clin Ther* 2011;33(3):254–79.
- [9] Zintzaras E, Dozani C, Mproitis T, Schmid CH, Hadjigeorgiou GM. Network analysis of randomized controlled trials in multiple sclerosis. *Clin Ther* 2012;34(4):857–869.e9. Epub 2012 Mar 22.
- [10] DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials* 1986;7(3):177–88.
- [11] Zintzaras E, Lau J. Synthesis of genetic association studies for pertinent gene-disease associations requires appropriate methodological and statistical approaches. *J Clin Epidemiol* 2008;61(7):634–45.
- [12] Bucher HC, Guyatt GH, Griffith LE, Walter SD. The results of direct and indirect treatment comparisons in meta-analysis of randomized controlled trials. *J Clin Epidemiol* 1997;50(6):683–91.
- [13] Glenny AM, Altman DG, Song F, Sakarovich C, Deeks JJ, D'Amico R, et al. International Stroke Trial Collaborative Group. Indirect comparisons of competing interventions. *Health Technol Assess* 2005;9(26):1–134. iii–iv.
- [14] Song F, Loke YK, Walsh T, Glenny AM, Eastwood AJ, Altman DG. Methodological problems in the use of indirect comparisons for evaluating healthcare interventions: survey of published systematic reviews. *BMJ* 2009;338:b1147.
- [15] Caldwell DM, Ades AE, Higgins JP. Simultaneous comparison of multiple treatments: combining direct and indirect evidence. *BMJ* 2005;331(7521):897–900.
- [16] Whitehead A. Meta-analysis of controlled clinical trials. Chichester, UK: John Wiley & Sons, Ltd., 2002.
- [17] S-PLUS® 8. Function guide. Seattle, WA: Insightful Corporation, 2002.
- [18] Compaq visual Fortran programmer's guide. Houston, TX: Compaq Computer Corporation, 2001.
- [19] Wallace BC, Schmid CH, Lau J, Trikalinos TA. Meta-analyst: software for meta-analysis of binary, continuous and diagnostic data. *BMC Med Res Methodol* 2009;9:80.
- [20] Loehrer Sr PJ, Ansari R, Gonin R, Monaco F, Fisher W, Sandler A. Cisplatin plus etoposide with and without ifosfamide in extensive small-cell lung cancer: a Hoosier Oncology Group study. *J Clin Oncol* 1995;13(10):2594–9.
- [21] Artal-Cortés A, Gomez-Codina J, Gonzalez-Larriba JL, Barneto I, Carrato A, Isla D, et al. Prospective randomized phase III trial of etoposide/cisplatin versus high-dose epirubicin/cisplatin in small-cell lung cancer. *Clin Lung Cancer* 2004;6(3):175–83.
- [22] Baka S, Califano R, Ferraldeschi R, Aschcroft L, Thatcher N, Taylor P, et al. Phase III randomised trial of doxorubicin-based chemotherapy compared with platinum-based chemotherapy in small-cell lung cancer. *Br J Cancer* 2008;99(3):442–7.
- [23] Eckardt JR, von Pawel J, Papai Z, Tomova A, Tzekova V, Crofts TE, et al. Open-label, multicenter, randomized, phase III study comparing oral topotecan/cisplatin versus etoposide/cisplatin as treatment for chemotherapy-naïve patients with extensive-disease small-cell lung cancer. *J Clin Oncol* 2006;24(13):2044–51.
- [24] Grote T, Yelding AL, Castillo R, Butler D, Fishkin E, Henry DH, et al. Efficacy and safety analysis of epoetin alfa in patients with small-cell lung cancer: a randomized, double-blind, placebo-controlled trial. *J Clin Oncol* 2005;23(36):9377–85.
- [25] Lee SM, James LE, Qian W, Spiro S, Eisen T, Gower NH, et al. Comparison of gemcitabine and carboplatin versus cisplatin and etoposide for patients with poor-prognosis small cell lung cancer. *Thorax* 2009;64(1):75–80.
- [26] Mavroudis D, Papadakis E, Veslemes M, Tsiakaki X, Stavrakakis J, Kouroussis C, et al. Greek Lung Cancer Cooperative Group. A multicenter randomized clinical trial comparing paclitaxel-cisplatin-etoposide versus cisplatin-etoposide as first-line treatment in patients with small-cell lung cancer. *Ann Oncol* 2001;12(4):463–70.
- [27] Miller AA, Herndon II JE, Hollis DR, Ellerton J, Langbein A, Richards II F. Schedule dependency of 21-day oral versus 3-day intravenous etoposide in combination with intravenous cisplatin in extensive-stage small-cell lung cancer: a randomized phase III study of the Cancer and Leukemia Group B. *J Clin Oncol* 1995;13(8):1871–9.

- [28] Miyamoto H, Nakabayashi T, Isobe H, Akita H, Kawakami Y, Arimoto T, et al. A phase III comparison of etoposide/cisplatin with or without added ifosfamide in small-cell lung cancer. *Oncology* 1992;49(6):431–5.
- [29] Niell HB, Herndon II JE, Miller AA, Watson DM, Sandler AB, Kelly K, Cancer and Leukemia Group. Randomized phase III intergroup trial of etoposide and cisplatin with or without paclitaxel and granulocyte colony-stimulating factor in patients with extensive-stage small-cell lung cancer: Cancer and Leukemia Group B Trial 9732. *J Clin Oncol* 2005;23(16):3752–9.
- [30] Noda K, Nishiaki Y, Kawahara M, Negoro S, Sugiura T, Yokoyama A, et al. Japan Clinical Oncology Group. Irinotecan plus cisplatin compared with etoposide plus cisplatin for extensive small-cell lung cancer. *N Engl J Med* 2002;346(2):85–91.
- [31] Okamoto H, Watanabe K, Kunikane H, Yokoyama A, Kudeh S, Asakawa T, et al. Randomised phase III trial of carboplatin plus etoposide vs split doses of cisplatin plus etoposide in elderly or poor-risk patients with extensive disease small-cell lung cancer. *JCOG 9702*. *Br J Cancer* 2007;97(2):162–9.
- [32] Pujol JL, Daurès JP, Rivière A, Quoix E, Westeel V, Quantin X, et al. Etoposide plus cisplatin with or without the combination of 4'-epidoxorubicin plus cyclophosphamide in treatment of extensive small-cell lung cancer: a French Federation of Cancer Institutes multicenter phase III randomized study. *J Natl Cancer Inst* 2001;93(4):300–8.
- [33] Roth BJ, Johnson DH, Einhorn LH, Schacter LP, Cherrig NC, Cohen HJ, et al. Randomized study of cyclophosphamide, doxorubicin, and vincristine versus etoposide and cisplatin versus alternation of these two regimens in extensive small-cell lung cancer: a phase III trial of the Southeastern Cancer Study Group. *J Clin Oncol* 1992;10(2):282–91.
- [34] Rowland Jr KM, Loprinzi CL, Shaw EG, Maksymiuk AW, Kurose SA, Jung SH. Randomized double-blind placebo-controlled trial of cisplatin and etoposide plus megestrol acetate/placebo in extensive-stage small-cell lung cancer: a North Central Cancer Treatment Group study. *J Clin Oncol* 1996;14(1):135–41.
- [35] Ruotsalainen TM, Halme M, Tamminen K, Szopinski J, Niiranen A, Pyrhönen S, et al. Concomitant chemotherapy and IFN- $\alpha$  for small cell lung cancer: a randomized multicenter phase III study. *J Interferon Cytokine Res* 1999;19(3):253–9.
- [36] Wolf M, Havemann K, Holle R, Gropp C, Drings P, Hans K, et al. Cisplatin/etoposide versus ifosfamide/etoposide combination chemotherapy in small-cell lung cancer: a multicenter German randomized trial. *J Clin Oncol* 1987;5(12):1880–9.
- [37] Zatloukal P, Cardenal F, Szczesna A, Gorbunova V, Moiseyenko V, Zhang X, et al. A multicenter international randomized phase III study comparing cisplatin in combination with irinotecan or etoposide in previously untreated small-cell lung cancer patients with extensive disease. *Ann Oncol* 2010;21(9):1810–6.
- [38] van der Valk R, Webers CA, Lumley T, Hendrikse F, Prins MH, Schouten JS. A network meta-analysis combined direct and indirect comparisons between glaucoma drugs to rank effectiveness in lowering intraocular pressure. *J Clin Epidemiol* 2009;62(12):1279–83.
- [39] Zintzaras E, Voulgarelis M, Moutsopoulos HM. The risk of lymphoma development in autoimmune diseases: a meta-analysis. *Arch Intern Med* 2005 Nov 14;165(20):2337–44.
- [40] Rothman KJ. No adjustments are needed for multiple comparisons. *Epidemiology* 1990;1:43–6.
- [41] Savitz DA, Olshan AF. Multiple comparisons and related issues in the interpretation of epidemiologic data. *Am J Epidemiol* 1995;142:904–8.
- [42] Bender R, Lange S. Adjusting for multiple testing – when and how. *J Clin Epidemiol* 2001;54:343–9.
- [43] Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *J Clin Epidemiol* 2009. <http://dx.doi.org/10.1016/j.jclinepi.2009.06.005>.
- [44] Balk EM, Bonis PA, Moskowitz H, Schmid CH, Ioannidis JP, Wang C, et al. Correlation of quality measures with estimates of treatment effect in meta-analyses of randomized controlled trials. *JAMA* 2002;287:2973–82.
- [45] Lu G, Ades AE. Assessing evidence inconsistency in mixed treatment comparisons. *J Am Stat Assoc* 2006;101(474):447–59.
- [46] Hasselblad V. Meta-analysis of multitreatment studies. *Med Decis Making* 1998;18(1):37–43.
- [47] Lumley T. Network meta-analysis for indirect treatment comparisons. *Stat Med* 2002;21(16):2313–24.
- [48] Higgins JP, Green S. *Cochrane handbook for systematic reviews of interventions*. Cochrane book series. Chichester, UK: John Wiley & Sons, Ltd., 2008.
- [49] Higgins JP, Whitehead A. Borrowing strength from external trials in a meta-analysis. *Stat Med* 1996;15(24):2733–49.
- [50] Black N. Why we need observational studies to evaluate the effectiveness of health care. *BMJ* 1996;312(7040):1215–8.
- [51] Song F, Altman DG, Glenny AM, Deeks JJ. Validity of indirect comparison for estimating efficacy of competing interventions: empirical evidence from published meta-analyses. *BMJ* 2003;326(7387):472.
- [52] Ziogas DC, Zintzaras E. Analysis of the quality of reporting of randomized controlled trials in acute and chronic myeloid leukemia, and myelodysplastic syndromes as governed by the CONSORT statement. *Ann Epidemiol* 2009;19(7):494–500.