



University of Thessaly
Department of Physical Education and Sport Sciences

Title

Physical Activity, Sleep Apnea and Quality of Life

By

Elise Löwenthal

A Thesis submitted in Partial Fulfillment of the Requirements for the Degree of Master in
Psychology of Exercise at the University of Thessaly in June 2019.

Approved by supervising committee:

Professor Antonis Hatzigeorgiadis
Professor Nikos Comoutos
Professor Yannis Theodorakis

Trikala, June 2019
Declaration by Author

This thesis is composed of my original work, and contains no material previously published or written by another person except where due reference has been made in the text. I have clearly stated the contribution by others to jointly-authored works that I have included in my thesis.

I have clearly stated the contribution of others to my thesis as a whole, including laboratory training to handle instruments, statistical assistance, data analysis, survey design, significant technical procedures, professional editorial advice, and any other original research work used or reported in my thesis.

I acknowledge that electronic and hard copies of my thesis must be lodged with the University Library.

I acknowledge that copyright of all material contained in my thesis resides with the copyright holder(s) of that material.

Statement of contributions by others to the thesis as a whole

The author of this thesis had a remarkable amount of help from Dr. Ioannis D. Morres, University of Thessaly.

20 June 2019

Name & signature of the author

Elise Löwenthal

Acknowledgements

My sincere appreciation and gratitude goes to the members of the Sleep Apnea project of the University of Thessaly, Professor Antonis Hatzigeorgiadis and Dr Ioannis D. Morres. Also, to Mrs. Beatrice Laurent from the Andre Vesal Hospital, Mrs. Sophie Lanoy from LUSS association and my master thesis supervisors and advisors: Dr. Ioannis D. Morres and Professor Anthonis Harzigeorgiadis.

Contents

Abstract	5
Introduction	6
Methods.....	14
Results.....	15
Discussion.....	19
Conclusion	28
References	29

Abstract

Obstructive sleep apnea is a medical condition characterized by nocturnal breathing cessation due to airway occlusion. Consequences and symptoms of the disease such as impaired quality of sleep can drastically impaired the quality of life of patients. The most commonly used treatment of OSA is positive airway pressure (PAP), however compliance is low and a range of 46 to 83% of patients do not adhere. A treatment alternative is physical activity (PA). It has been shown to improve sleep apnea condition. However, there are a limited number of studies examining the association of physical activity and quality of life in obstructive sleep apnea patients. The purpose of the present study was to investigate the relation between physical activity and quality of life in sleep apnea patients.

Questionnaires were used to assess the different variables of the study: The IPAQ was used to assess physical activity. Epworth Sleepiness Scale was used to assess daytime sleepiness and give an indication of the sleep apnea severity. Finally, the SF-12 was used to measure the health related quality of life. Seventy-eight sleep apnea patients completed the questionnaires. Results showed a positive relationship between low intensity physical activity and quality of life. A relationship has been found between quality of life and sleep apnea severity. Finally, low intensity physical activity predicted quality of life in sleep apnea patients.

The findings of this thesis support that physical activity could be an adequate treatment option, improving the quality of life and the sleep apnea symptoms in sleep apnea patients.

Keywords: sleep apnea, physical activity, quality of life, obstructive sleep apnea.

Introduction

Sleep apnea is a medical condition which causes repetitive episodes of nocturnal breathing cessation (during from a few seconds to a few minutes) (Spicuzza et al., 2015) and can be Obstructive Sleep Apnea (OSA), and be caused by airway occlusion. Finally, sleep apnea can be central (CSA), and lead to lack of breathing control. This study focus on obstructive sleep apnea which is the most common form. Both OSA and CSA lead to negative consequences at different levels: symptoms such as daytime sleepiness, loud snoring and restless sleep are characteristics of the disorder (Motamedi et Al., 2009). Other negative manifestations accompanying OSA can be depression, irritability, mood disorders, impaired cognition, difficulty of concentration or short-term memory loss. It affects the personal life at a familial, professional and social level (Coman, 2016).

Prevalence

Based on a systematic review including 24 studies, the overall population prevalence of sleep apnea ranges from 9 percent to 38 percent and is higher in men. Based on 11 epidemiological studies published between 1993 and 2013, the prevalence of OSA for men is estimated to be from 9 to 37 percent (Franklin & Lindberg, 2015). For women, the prevalence is estimated to be from 4 to 40 percent. However, another study estimates the prevalence for OSA being from 3 to 7 percent of the population (Punjabi, 2008). Some factors make people more at risk: the age, being a male, obesity, family history, menopause, craniofacial abnormalities, and certain health behaviors such as cigarette smoking and alcohol use.

Sleep Apnea and Health related Quality of Life

Health related quality of life regroups aspects of a patient's life such as his emotional and mental health, his physical condition and energy (Gandek et al., 1998) It can be defined as the patient's perception of their position in life, related to their particular disease and treatment (Sosnowski et al., 2017). The state of health has an influence on the everyday functioning of individuals.

Symptoms, consequences and treatments of diseases have their consequences on someone's life and functioning (Schipper et al., 1990). The World Health Organization defines HRQoL as something that has to be evaluate in the context of the culture and value systems of individuals., taking goals, expectations, standards and concerns into account.” (Sosnowski et al., 2017).

HRQoL has an objective component, including the direct physical symptoms of a particular disease, as well as a subjective component, including the patient's personal opinions, emotions and feelings related to their state. Different aspects: such as general health, physical functioning, body pain, mental health and vitality, social functioning, emotional and mental health are components of the HRQoL. (Huo et al., 2018). All those aspects can be measured by both the SF-36 and the SF-12.

OSA patients' state of health has different repercussions on the everyday functioning. Obstructive sleep apnea impairs QoL at different level. First of all, the patient's quality of life is impaired by daytime symptoms (Lacasse et al., 2002): One of OSA's main symptom and consequence is daytime sleepiness. Patients have an unrefreshing sleep, they can have difficulty staying awake during daytime and lack control over their ability to stay awake. In addition to having episodes of suddenly falling asleep, they might have consequences such as lack of concentration, attention and memory deficit. (Naegele et al, 1995). Sleep apnea can lead to physical symptoms such as congested nose, dry mouth or sorethroat or headaches upon wakening, directly impairing health related quality of life (Clarenbach et al., 2008; Mannarino et al, 2012). Patients are limited in their everyday life functioning as they might not have the control on the activities that can do; they might have difficulty doing activities such as reading or driving. They can have the feeling of having to fight a urge to sleep while doing an activity. They might need to take naps during daytime, preventing them from doing what they want. Another way quality of life of OSA patients is affected is by the disturbed quality of sleep (Colt et al., 1991): snoring is a common symptom and consequence. Patients tent to often wake up

during the night and suffer from restless sleep. Sometimes patients wake up during the night with a sensation of choking (Mannarino et al., 2012). Due to the impair quality of sleep, OSA patients tends to wake up early in the morning, limiting their duration of sleep. It is common for them to wake up with a dry mouth. Sleep apnea patients are also affected at an emotional level: consequences such as depression and anxiety are common. OSA patients might have concerns about their health and weight. (Wittels et al., 1990) Finally, quality of life of OSA patients is impaired at a social level. Patients might be worried of their snoring impairing their partner and friends' sleep. Due to their lack of energy, sleep apnea patients might participate in less social activities (Moore et al., 2001).

Consequences of lack of quality sleep

Sleep apnea prevents from quality sleep. Sleep disruption leads to both short- and long-term consequences: increased stress responsivity, emotional distress, mood disorders, impact on cognition and memory, and performance deficits. Somatic problems, as weekly headache or abdominal pain are others possible repercussions of sleep deprivation (Luntamo et Al.,2012). In the long run, sleep disruption consequences include hypertension, dyslipidemia, weight-related issues, metabolic syndrome, poor appetite regulation, poor functioning of immune, hormonal and cardiovascular systems (Medic, 2017).

Cardiovascular consequences

Obstructive sleep apnea is seen in 37 percent of 450 and 11 percent of 81 patients with heart failure resulting from systolic dysfunction (Somers et al., 2008). Also, OSA is associated with a significant cardiovascular morbidity and mortality. Cardiovascular consequences, such as hypertension, resistant systemic hypertension, chronic heart failure, arrhythmia, myocardial infarction and stroke can be caused by sleep apnea. Specifically, OSA is characterized by intermittent hypo xemia and CO₂ retention during sleep, with oxygen saturation dropping up to 60 percent. These factors cause cardiac and vascular diseases.

Depression

Depression is a mood disorder, it leads to different symptoms such as experience of persistent feelings of sadness, loss of interest in activities and hopelessness (American Psychiatric Association,2013). Depression in OSA patients is quite prevalent (Ejaz et al., 2011): specifically, the prevalence of depression is higher in patients with OSA as compared to the general population according to a study with 182 patients which found that 25 percent of participants (47 patients) were suffering from depression (Shoib, 2017). Studies favor a mutual relationship between OSA and depression. The relationship remains unclear, as different mechanisms could explain how OSA can worsen depression and vice versa (Ejaz et al., 2011).

Anxiety

Generalized anxiety disorder is characterized by different symptoms: excessive anxiety and worry about various topics, difficulty to control the anxiety, impaired concentration, irritability, tiredness. A research study that examined the association between OSA and other comorbid disorders in Veterans Health Administration beneficiaries (Sharafkhaneh et al., 2005) found that 16,7 percent of the people suffering from sleep apnea were also suffering from anxiety. Also, 11,9 percent were suffering from posttraumatic stress disorder, 21,8 percent were suffering from depression (Sharafkhaneh et al., 2005). This is a significantly greater prevalence in comparison to patients not diagnosed with sleep apnea (Sharafkhaneh et al., 2005). Also, disruption and discontinuity of sleep has a bigger role than the duration of sleep deprivation for increased stress responsivity (Tiemeier et Al., 2002)

Treatment

The most common treatment used for OSA is positive airway pressure (PAP), which is typically recommended for mild, moderate, and severe OSA (Epstein et al., 2009). PAP can be delivered

in continuous (CPAP), bilevel (BPAP), or autotitrating (APAP) modes (Epstein et al., 2009). BPAP machines differ from CPAP: the first one have two pressure settings: one for inhalation and one for exhalation.

Treatment consequences

A range from 46 to 83% of patients with obstructive sleep apnea do not adhere to PAP treatment. Nasal continuous airway pressure can have side effects as getting allergy to the face, air leaks, abrasions of the ridge of the nose (Pépin et al., 1995). PAP has been shown to have negative side effects at different levels (Gay and al., 2006). The material and design of the mask itself can cause allergy, claustrophobia, mask leak, skin abrasion or conjunctivitis eyes. PAP can also have pressure-related side effects, such as rhinitis, sinusitis, headache or pressure intolerance (Gay and al., 2006). The equipment can be perceived as noisy, cumbersome and invading the intimacy of the patient, moreover it needs to be taking care of and cleaned (Gay and al., 2006)..

Sleep apnea and physical activity

People with untreated obstructive sleep apnea may engage in less physical activity. Fatigue and daytime sleepiness that results from sleep apnea might influence on the physical activity participation (Jean et al., 2017). Specifically, OSA impair exercise capacity and increase cardiovascular risks by several mechanisms (Beitler et al., 2014): a difference in the lactate concentration and a delay in the elimination of lactate have been observed in OSA subjects in comparison to age and BMI-controlled group. Patients with OSA have been shown to have abnormal cardiovascular responses during exercise or recovery. Increasing of the diastolic blood pressure, decrease of the stroke volume, attenuation of the heart rate during peak exercise and recovery or global ventricular dysfunction have been shown.

Sleep apnea and sedentariness

The level of physical activity of OSA patients is low. A meta-analysis (Mendelson et al., 2018) explored on one side the objective physical activity levels of patients with OSA and on the other side, the effects of exercise training on OSA severity, body mass index, sleepiness and cardiovascular fitness. The meta-analysis showed that OSA patients were not physically active. The second part of the meta-analysis found a significant decrease in apnea-hypopnea index following exercise training (mean decrease of 8.9 events/h; 95% CI: -13.4 to -4.3; $p < 0.01$). Also, a reduction in subjective sleepiness, an increase in VO₂peak and no change in BMI were observed.

Spicuzza (2015) reported that weight loss improves symptoms and morbidity in all OSA patients and claimed that a multidisciplinary approach is necessary for an accurate management of the disease.

Obesity is associated with sleep apnea (Schwartz et al., 2008). Adiposity around the pharynx and torso might narrow the upper airway and increase the chance of upper airway collapsibility. Also, obesity is associated with a reduction in lung volumes. A reduction in weight is an effective strategy for treating sleep apnea in obese patients. Physical activity is an effective treatment that can be used in the aim of reducing the weight of obese people. It can solve the anatomic alterations of obesity people and reduce sleep apnea symptoms. However, physical activity may also be beneficial without weight loss consequences.

Physical Activity as A Treatment of OSA

Positive Airway Pressure have been shown to have negative side effects and can increase comorbidity. A range of 46 to 83 percent of patients do not comply to this treatment, the symptoms of OSA could be difficult to manage by those people (Kenneth et al., 2016). Other treatments for OSA and its symptoms that are less invasive are essential: exercise is an option to improve OSA indices, it would constitute a low-cost and easy-to-use treatment modality (de Andrade et al., 2016).

Exercise training has moderate treatment effects on the reduction of apnea-hypopnea index (Kline, 2011) Results suggested that exercise may be beneficial for the management of OSA beyond simply facilitating weight loss. The study included forty-three sedentary and overweight adults with at least moderate-severity untreated OSA, the exercise treatment consisted in 150 minutes of exercise per week for 12 weeks. Exercise consisted in moderate-intensity aerobic activity once a week followed by resistance training twice a week. The control group met twice weekly for 12 weeks and perform low-intensity exercises designed to increase whole-body flexibility.

Sengul et al. (2009) in their RCT found that breathing exercise and other types of exercise have a beneficial effect on apnea-hypopnea index, quality of sleep, exercise capacity and health related quality of life in patients with mild to moderate OSA. The exercise group followed a program consisting in 1.5 hours of exercise, three days a week for 12 weeks whereas the control group did not receive any treatment.

A meta-analysis (Iftikhar and al.,2014), using 5 studies with 129 participants in total, showed a statistically significant effect of exercise in reducing the severity of sleep apnea in patients with OSA with minimal changes in body weight. The potential value of exercise was due to its effect on cardiorespiratory fitness, daytime sleepiness, and sleep efficiency.

Another meta-analysis (Aiello, 2016) using 8 articles with 182 participants also conclude that exercise has an effect on the severity of apnea. Exercise induces a decrease on the apnea/hypopnea index and decreases sleepiness during day time.

Physical activity effects on Quality of Life in OSA patients

Conn et al. (2009) have shown with a meta-analysis that interventions to increase physical activity increase quality of life: the mean of quality of life effect size was .11 for two groups comparison and .27 for pre-post comparison. The study included Eighty-fives samples with 7 291 subjects.

Physical activity increases the quality of life of OSA patients ($P < 0.01$) (Ueano et al., 2009). The study included twenty-five patients with heart failure with obstructive, central or no sleep apnea. The interventions consisted of four months of no-training followed by four months of an exercise training program.

The effects of Physical activity on Depression in OSA patients

Physical exercise is associated with reduction of depression symptoms. Specifically, a systematic review and meta-analysis that included 455 patients across 11 trials found that aerobic exercise is an effective antidepressant intervention that has a large overall effect (Morres et al., 2018).

Compared to control group, physical exercise showed higher effects on depression symptoms in OSA patients (Kline et al., 2012). The clinical trial included forty-three overweight/obese and sedentary adults (18-55 years old) with untreated OSA.

Method

Participants

Recruitment was conducted through the Sleep Apnea Project of the University of Thessaly. A total of 51 participants were patients of the University of Thessaly Hospital (Larissa, Greece). A total of 27 patients were referred by a nurse from the André Vésale Hospital (Charleroi, Belgium). A written consent form was acquired before participation by all patients.

Participants were 78 adults with obstructive sleep apnea (51 males and 27 females), with a mean age of 48.45 (SD= 10.63) years. Participants had to be a minimum of 17 years old to be included in this study.

Procedure

Questionnaires were used to assess the variables of the present study. The French version of the SF-12 Survey, a shorter version of the SF-36, was used to measure functional health and well-being from the patient's point of view (Ware et al., 1996). The Epworth Sleepiness Scale was used to assess daytime sleepiness (Johns, 1991). The IPAQ (International Physical Activity Questionnaire) was used to measure physical activity (Craig et al., 2003).

Results

With respect to Quality of Life and METS variables, correlations showed a moderate significant relationship between quality of Life and low intensity physical activity (METs low) (.23, $p=.046$), no significant relationship have been found with the variables METS moderate (.20, $p=.087$) and METS high (.15, $p=.743$). Mets low predicted quality of life.

A significant relationship has been shown between quality of life and Ipaq item 1 (.24, $p=0.035$) and Ipaq item 5 (.29, $p=.014$). Item 1 refers to high intensity physical activity (self-reported days of high intensity physical activity), item 5 refers to the number of days someone walked for at least 10 minutes.

Finally, a significant relationship has been found between quality of life and METS total (.28, $p=.015$). Mets total refers to an estimated metabolic equivalents of total intensity physical activity, it includes low, moderate and high intensity.

A significant relationship has been found between Quality of Life and Epworth, this variable showed a moderate negative correlation (-.41, $p=.000$).

With respect to Epworth and IPAQ, a significant relationship has been found with IPAQ_Item 7 (.27, $p=.021$). Ipaq Item 7 refers to sedentariness (time spent seated). No significant correlations have been found between the variables Epworth and IPAQ total (-.11, $p=.285$).

Results can be found in Table 1 and Table 2.

Table 1. Descriptive Statistics for studied variables

	Minimum	Maximum	Mean	Standard Deviation
Age	17	77	48,45	10,633
IPAQ-1	0	6	,58	1,274
IPAQ-2	0	240	47,09	67,623
IPAQ-3	0	7	1,53	2,106
IPAQ-4	0	180	61,03	50,380
IPAQ-5	0	7	4,30	2,408
IPAQ-6	0	480	47,92	65,680
IPAQ-7	2	720	87,60	148,866
METs IPAQ Low	,00	1680,00	208,1081	278,54527
METs IPAQ Moderate	,00	980,00	91,4359	170,27515
METs IPAQ High	,00	1200,00	48,7564	164,09802
METs IPAQ Total	,00	2600,00	355,8784	421,78187
Sleepiness	,00	18,00	6,7564	4,42597
QoLife Physical	6,00	25,00	19,1667	4,38835
QoLife Mental	10,00	29,00	21,1026	4,79066
QoLife Total	18,00	54,00	40,2692	8,13151

QoLife : Quality of Life; IPAQ : International Physical Activity Questionnaire; METs: Metabolic Equivalents

Table 2. Correlation coefficients for studied variables

	Variable	1	2	3	4	5	6	7	8	9	10	11	12	13	14
1	IPAQ-1														
2	IPAQ-2	.48**													
3	IPAQ-3	.32**	.11												
4	IPAQ-4	.10	.31	.08											
5	IPAQ-5	.10	.07	.22	.18										
6	IPAQ-6	-.00	.47**	-.13	.24	.04									
7	IPAQ-7	.16	-.35	.21	-.03	-.19	-.13								
8	METs Low	-.00	.36*	-.09	.23	.35**	.86**	-.19							
9	METs Moderate	.31**	.43*	.66**	.64**	.23*	-.02	.20	.05						
10	METs High	.68**	.84**	.30**	.26	.07	.11	.07	.08	.51**					
11	METs Total	.39**	.70**	.32**	.50**	.35**	.60**	-.02	.71**	.65**	.66**				
12	Sleepiness	-.08	-.08	.06	-.15	-.19	-.02	.27*	-.11	-.08	-.04	-.13			
13	QoLife Physical	.28*	.29	.19	.21	.16	.08	.10	.14	.19	.17	.23	-.29*		
14	QoLife Mental	.15	.14	.14	-.03	.34**	.17	-.08	.26*	.16	.10	.16	-.43**	.57**	
15	QoLife Total	.24*	.23	.19	.09	.29*	.15	.01	.23*	.20	.15	.28*	-.41*	.88**	.90**

QoLife : Quality of Life; IPAQ: International Physical Activity Questionnaire; METs: Metabolic Equivalent;

*: Statistically Significant <.05 **: Statistically Significant <.001

Table 3. Regression analysis for quality of life

	Beta	t	p	F	R ² change
Significant predictor				4.12*	.23
Mets Low (Light Physical Activity)	.23	2.03	.046		
Excluded Predictors					
Mets High (Vigorous Physical Activity)	.12	1.08	.28		
Mets Moderate (Moderate Physical Activity)	.17	1.45	.15		

*: Statistically Significant at .05

Discussion

Physical activity and quality of life in OSA patients

The results of this study found that low intensity physical activity was significantly correlated and predicted quality of life. Although the numbers of days of high intensity physical activity was associated with an increase in quality of life, METs high was not correlated with quality of life. This suggests that low intensity physical activity could be an adequate treatment to increase quality of life of OSA patients.

Low physical activity is an easier treatment option in comparison to moderate or high intensity physical activity. Walking or doing activities such as house cleaning or gardening are some example of low intensity physical activity. This suggests that simply a change of lifestyle could be effective in improving the quality of life of sleep apnea patients. Moreover, those activities and this type of physical activity does not necessitate a gym.

In term of motivation, it could be assumed that it is easier to motivate sleep apnea patients to do low intensity physical activity in comparison to moderate or high intensity. A study from Perri et al. (2002) found that increasing exercise intensity may lower adherence while increasing exercise frequency did not show any decline in adherence. The same study found that a high intensity exercise was correlate to higher exercise-reported injuries. Low intensity physical activity might be a good treatment option for sleep apnea patients as there is less chances to get injured. (Perri et al., 2002)

In the literature, physical activity has been found to be associated with quality of life in various populations: in a 2007 meta-analysis including 55 studies, Bize et al.(2007) showed a consistently positive association between physical activity and health related quality of life in the general population. A study by Rejeski et al. (2001) analysed studies on physical activity and quality of life among older adults in various settings: physical activity and exercise were found to be related to better quality of life in retirees men (Peppers, 1976), in adults with

physical disabilities (Jette et al., 1999), in adults with chronic obstructive pulmonary disease (Emery et al., 1998) in patients with myocardial infarction and other heart diseases (Tyni-Lenné et al., 1998). Finally, physical activity have been found to be related with improved quality of life in older adults with minor to major depression (Singh et al., 1997).

Specifically, in OSA patients, studies have found that physical activity and exercise improves quality of life. Studies addressed physical activity in the aspect of exercise such as aerobic and resistance training or physical activity in a general way. These effects of exercise on quality of life favored CPAP. (Ackel-D'Elia et al., 2012).. Other studies assessed exercise in association with a weight-reduction diet (Barnes et al., 2009).

Exercise training induce small to significant improvements of the quality of life components in OSA patients. That is what has been found in a study by Kline et al. (2012) on 43 participants using the SF-36 questionnaire. The exercise training consisted in moderate-intensity aerobic and resistance training, six days a week for 12 weeks. The control group was doing stretching. Significant improvements have been found in physical functioning, vitality and mental health. Small to moderate sized improvements have been found in all aspects of quality of life except for bodily pain, social functioning and role limitations due to emotional health.

Regular physical activity has been found to be a better predictor than sleep apnea severity in predicting perceptions of energy and fatigue; regular physical activity is significantly correlated with higher SF-36 vitality. (Hong et al., 2003). The results remain significant after controlling for BMI.

A study by Barnes et al. (2009) on 21 OSA patients following a weight reduction diet and an exercise program found a significant effect on quality of life, however no effects have been found for the sleep-specific quality of life questionnaire. (Functional Outcomes of Sleep Questionnaire). The intervention consisted in 16 weeks of resistance, aerobic training (80% VO₂PEAK) and a low-energy meal diet.

In comparison to a CPAP treatment only, a 2-month exercise training (running and walking, 85 percent of the anaerobic threshold) combined with CPAP treatment has been found to have a significant and better impact on quality of life (physical functioning and general health perception) as well as on subjective daytime sleepiness and mood states (Ackel-D'Elia et al., 2012).

Sleep Apnea and Quality of Life

In accordance with the literature, a significant moderate relationship has been found between quality of life and sleep apnea severity. Quality of life can be defined as the patient's perception of their position in life, related to their particular disease and its treatment. (Sosnowski et al., 2017).

The results of this thesis aligned with the current literature: poor quality of life has been found to be one of the most common complaint among obstructive sleep apnea patients. Impaired daytime functioning, excessive daytime sleepiness, fatigue and other consequences in cognitive function, and mood are characteristics of OSA (Kline et al., 2012). Subjective distress and poor quality of life in general are frequently reported by OSA patients, even after adjusting for body mass index (BMI), gender, age and comorbidity (Hong et al., 2003).

Mediators and moderators

The relation between physical activity and quality of life can be explained by different variables: studies have looked at the mediators and moderators between physical activity and quality of life (Joseph et al., 2014; Rejeski et al., 2001). In general population and in sleep apnea patients, physical activity can increase quality of life by different mechanisms including enjoyment of an activity, self-efficacy feeling, self-esteem enhancement and increase in energy. Some of the variables might play a bigger role for the sleep apnea patients; increase in energy might be particularly beneficial for them as their condition implies that they suffer from low energy and fatigue.

One explanation that has been found in the literature is that physical activity is an enjoyable activity in itself and that the repeated enjoyment that someone gets from the activity enhance his quality of life. Fox et al. (2000) has found that manipulating social interaction in an exercise activity such as aerobic dance impacts on the enjoyment perceived. The authors suggest that the enjoyment created by a socially enhanced environment could improve quality of life.

Rejeski et al. (2001) suggests that the feeling of self-efficacy, the individual belief in its capacity can mediate the relation between physical activity and quality of life. Self-satisfaction and the feeling of being proud of yourself can also be a mediator.

Physical activity has been shown to enhance self-esteem, which can also be a mediator. Self-esteem is a predictor of quality of life. A study by Elavsky et al. (2009) has found that both self-esteem and positive feeling state induced by physical activity were correlated to long-term quality of life in older adults.

Another potential mediator is energy. Physical exercise has been show to increase feeling of energy and decrease feeling of fatigue. (Rejeski et al., 2001)

The value someone placed on physical activity can be a moderator. (Rejeski et al., 2001).

Those mediators seem to be comparable with the self-determination theory which suggests that the satisfactions of the three psychological needs of autonomy, relatedness and competence leads to autonomous motivation to be physically active.

Physical Activity and Epworth

The results of this thesis shows that sleep apnea severity (measured by the Epworth) was correlated to sedentariness and the amount of time in a day a patient spent seated.

Impaired day-time functioning is one of the main complain of sleep apnea patients. One of the major symptom of OSA is day-time sleepiness; sleep apnea patients feel sleepy during day-time; they are at risks of falling asleep during various activities (Kline et al., 2012). The likelihood of falling asleep is assess by the Epworth Sleepiness Scale: patients have to report if

they fall asleep in eight different sedentary activities. Various studies have demonstrated a positive impact of physical activity on sleepiness: Hong et al. (2003) demonstrated that physical activity was significantly correlated with daytime sleepiness in sleep apnea patients.

The literature has demonstrated that physical activity was associate with reduction of day-time sleepiness in sleep apnea patients and in other medical population.

A study by Kline et al. (2012) including forty-three sedentary overweight OSA patients demonstrated that exercise training was positively correlated to improvement in day-time functioning, including day-time sleepiness. Exercise training may be helpful for improving aspects of day-time functioning of adults with OSA: exercise training produces a moderate size reduction in day-time sleepiness measured on both the ESS and FOSQ-10. Participants of the study were randomized into two groups: one following a 12-weeks plan of moderate-intensity aerobic and resistance training, the other following a low-intensity stretching control treatment. A study on insomnia patients compared aerobic exercise and sleep hygiene, to sleep hygiene only. The combination of aerobic exercise and sleep hygiene was associate with a reduction in daytime sleepiness. (Reid K.J. et al., 2010)

A meta-analysis on 12 population-based studies has found that physical activity was associated with a 40 percent reduced risk of experiencing feeling of low energy and fatigue. (Puetz, 2006) The duration and frequency of exercise might have an impact on fatigue perception. Andreassen et al. (2011) have studied the effect of exercise therapy on fatigue in patients with multiple sclerosis (MS). The study demonstrated that longer and low-frequency protocols might be more efficient to reduce fatigue in comparison to high-frequency and short-duration protocols for MS patients. The authors suggest that a social component could influence the fatigue's level of perception. Group exercise and contacts with other participants might play a role.

A randomized controlled trial from Puet et al. (2008) reported that low-intensity exercise had a positive effect on the level of perceived fatigue in sedentary young adults with persistent

fatigue. Moderate exercise has been found to have a positive impact on health outcome such as reduction of depression but did not have an impact on fatigue perception. These authors suggested that moderate exercise might be too demanding for a population reporting persistent fatigue. The same study assessed the perception of energy and found that both low and moderate intensity has a positive impact. The two different patterns for the variables of fatigue and energy perception remained unclear.

Self-determination Theory

The literature review and the results of this study suggests that physical activity is an effective treatment to improve the condition of sleep apnea patients and improve their quality of life. Motivating sleep apnea patients to be physically active might be an important issue. The self-determination theory is a theory of motivation that could be effective to motivate sleep apnea patients to be active.

The SDT is a theory of human motivation and personality, according to the self-determination theory, the continuum of motivation is described by lack of motivation (amotivation), then by extrinsic motives and is finalized by intrinsic motives. The extrinsic motives include external motivation which refers to behavior for rewards and awards and by introjected motivation which refers to a behavior that is implemented by the person to avoid feeling of guilt. As far as the intrinsic motive is concern, this includes internal motivation which refers to a behavior that is implemented for joy and pleasure, and the identified motivation which refers to a behavior that is implement due to personal values and reasons. In SDT, also, both external and introjected motives comprise the so called "controlled motivation", whereas both internal and identified motives comprise the so called "autonomous motivation".

The self-determination theory also includes three dendrites which are the three basic psychological needs of competence, autonomy and relatedness. Competence refers to the need of feeling effective and developing, autonomy refers to the need of having the feeling of self-

direction and self-endorsement in action, relatedness refers to the need of feeling meaningfully connected to others.

The self-determination theory also reports that when these three psychological needs are satisfied, they catalyze the facilitation of autonomous motivation (the internal and identified motives). As far as the satisfaction of the three psychological needs is concern, SDT suggests the synergetic satisfaction of all three psychological needs.

The SDT have a great reach, encompass concepts that have personal meaning, leads to empirical methodologies and use principles that can be apply on different life context: parenting, health care, education, work, sport and psychotherapy for example.

The theory postulates that all individuals naturally want to be active, motivated, curious and willing to succeed, as success is gratifying in itself. The theory also recognizes that some individuals can be passive and unmotivated. Those difference in motivation are explained by the different type of motivation: autonomous and controlled. The type of motivation that an individual has results from the interaction between the inherent active nature of individuals and the environment, which can be supporting or thwarting.

Physical Activity and SDT

Various studies demonstrated that the self-determination theory is efficient in a physical activity and exercise context. In a meta-analysis including 184 independent data set, Johan et al. (2012) demonstrated an existent relation between psychological need satisfaction, autonomous motivation and beneficial health outcomes. The meta-analysis included various study in health care and health promotion contexts. Specifically, a study by Silva et al. (2011) showed the links between psychological needs satisfaction, autonomous motivation and exercise adherence: satisfaction of competence, relatedness and autonomy is related to autonomous motivation and exercise adherence. A systematic review by Teixeira et al. (2012) showed a positive relation

between autonomous forms of motivation and exercise; in most of the included studies, identified regulation was predicting initial/short term adoption and intrinsic motivation was predictive of long-term exercise adherence. Competence satisfaction positively predict exercise participation across a range of samples and settings.

A study by Edmunds et al. (2006) investigated the adherence of obese/overweight people on physical activity. Individuals who adhered more reported more self-efficacy to overcome barriers to exercise and demonstrated an increase in relatedness need satisfaction over time. The study also found that need satisfaction predicted self-determined regulation.

Sleep Apnea and SDT

The self-determination theory could also be beneficial to sleep apnea patients in other context that are not related to physical activity. The satisfaction of the three psychological needs in life can have an impact on sleep; a study by Campbell et al. (2015) including 215 adults found that the satisfaction of the three psychological needs of competence, relatedness and autonomy was positively related to two components of sleep: perceived sleep quantity and quality (Pittsburgh Quality Index) and was negatively related to day-time dysfunction, which was assessed with the Insomnia and Lassitude subscales as well as the Fatigue Severity scale and the General Vitality scale.

According to Campbell et al. (2015), the positive relation between psychological needs satisfaction and improved sleep could be related to two facts. First, people who satisfy their psychological needs are more likely to encounter positive daily experiences, second, to have positive thoughts when falling asleep.

Another finding of the study (Campbell et al., 2015) is that the psychological needs satisfaction are partial mediators of the relation between sleep quality and quantity and daytime dysfunction with financial strain and mindfulness. Mindfulness is positively related to sleep quality and quantity and negatively related to day-time dysfunction. This relation could be explained by the

partial mediating effect of psychological needs satisfaction. As far as the financial strain is concerned, it seems that it is negatively related to sleep quality and quantity and day-time function. This infers that the relationship is maybe due to the threatening impact of financial strain on psychological needs.

SDT and quality of life

According to the SDT model, the patient's experience of autonomy, competence and relatedness is associated with better mental health (fewer depressive symptoms, anxiety and somatization), better health related outcomes, including more physical activity, as well as greater quality of life (Ryan et al., 2008). A study by Chen et al. (2018) found a positive relation between hemodialysis patient's perceptions of autonomy support and health related quality of life (physical and mental component) through basic psychological needs satisfaction. This infers that the autonomy support provided by physicians and nurses contributes to the improvement of the health-related quality of life of patients through basic need satisfaction. A study by Duda et al. (2014) comparing an SDT-grounded physical activity consultation with a standard provision exercise referral have found a difference in term of participant's psychological well-being outcomes. The study concluded that it could be due to the basic needs' satisfaction promoting environment.

Studies and meta-analysis have shown that the satisfaction of psychological needs is related to quality of life in different populations and contexts. Studies in school contexts, health behavior contexts, on adolescents, on people suffering from different diseases such as HIV or diabetes have investigated this relation and found that psychological needs satisfaction was positively related to quality of life improvement. (NG et al., 2012; Gillison et al., 2006; Shah et al., 2016; Standage et al., 2012)

Conclusion

The results of this study supports the existing literature review on physical activity and quality of life. In this Thesis we found that a relation exists between physical activity and quality of life and in sleep apnea patients. Specifically, low intensity physical activity predicted better quality of life. An association between sleep apnea severity and sedentariness has been found. Finally, the results showed a significant relation between quality of life and sleep apnea severity. These findings support the fact that low intensity of physical activity could be an effective option to improve quality of life of sleep apnea patients.

References

- Ackel-D'Elia C, da Silva AC, Silva RS, et al. Effects of exercise training associated with continuous positive airway pressure treatment in patients with obstructive sleep apnea syndrome. *Sleep Breath*. 2012;16(3):723–735.
- Aiello, K. D., Caughey, W. G., Nelluri, B., Sharma, A., Mookadam, F., & Mookadam, M. (2016). Effect of exercise training on sleep apnea: a systematic review and meta-analysis. *Respiratory medicine*, 116, 85-92.
- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed., text rev.).doi:10.1176/appi.books.9780890423349.
- Andreasen, A. K., Stenager, E., & Dalgas, U. (2011). The effect of exercise therapy on fatigue in multiple sclerosis. *Multiple Sclerosis Journal*, 17(9), 1041-1054.
- Barnes, M., Goldsworthy, U. R., Cary, B. A., & Hill, C. J. (2009). A diet and exercise program to improve clinical outcomes in patients with obstructive sleep apnea--a feasibility study. *Journal of clinical sleep medicine : JCSM : official publication of the American Academy of Sleep Medicine*, 5(5), 409–415.
- Beitler, J. R., Awad, K. M., Bakker, J. P., Edwards, B. A., DeYoung, P., Djonlagic, I., ... & Malhotra, A. (2014). Obstructive sleep apnea is associated with impaired exercise capacity: a cross-sectional study. *Journal of Clinical Sleep Medicine*, 10(11), 1199-1204.
- Campbell, R., Vansteenkiste, M., Delesie, L. M., Mariman, A. N., Soenens, B., Tobbac, E., ... & Vogelaers, D. P. (2015). Examining the role of psychological need satisfaction in sleep: A Self-Determination Theory perspective. *Personality and Individual Differences*, 77, 199-204.
- Chen, M. F., Chang, R. E., Tsai, H. B., & Hou, Y. H. (2018). Effects of perceived autonomy support and basic need satisfaction on quality of life in hemodialysis patients. *Quality of Life Research*, 27(3), 765-773.

- Clarenbach, C. F., Kohler, M., Senn, O., Thurnheer, R., & Bloch, K. E. (2008). Does nasal decongestion improve obstructive sleep apnea?. *Journal of sleep research*, 17(4), 444-449.
- Colt HG, Haas H, Rich GB. Hypoxemia versus sleep fragmentation as a cause of excessive daytime sleepiness in obstructive sleep apnea. *Chest* 1991; 100: 1542–1548
- Conn, V. S., Hafdahl, A. R., & Brown, L. M. (2009). Meta-analysis of quality-of-life outcomes from physical activity interventions. *Nursing research*, 58(3), 175-83.
- Craig, C. L., Marshall, A. L., Sjöström, M., Bauman, A. E., Booth, M. L., Ainsworth, B. E., ... & Oja, P. (2003). International physical activity questionnaire: 12-country reliability and validity. *Medicine & science in sports & exercise*, 35(8), 1381-1395.
- De Andrade FMD, Pedrosa RP. The role of physical exercise in obstructive sleep apnea. *Jornal Brasileiro de Pneumologia*. 2016;42(6):457-464. doi:10.1590/S1806-37562016000000156.
- Duda, J. L., Williams, G. C., Ntoumanis, N., Daley, A., Eves, F. F., Mutrie, N., ... & Jolly, K. (2014). Effects of a standard provision versus an autonomy supportive exercise referral programme on physical activity, quality of life and well-being indicators: a cluster randomised controlled trial. *International Journal of Behavioral Nutrition and Physical Activity*, 11(1), 10.
- Edmunds, J., Ntoumanis, N., & Duda, J. L. (2007). Adherence and well-being in overweight and obese patients referred to an exercise on prescription scheme: A self-determination theory perspective. *Psychology of Sport and Exercise*, 8(5), 722-740.
- Ejaz, S. M., Khawaja, I. S., Bhatia, S., & Hurwitz, T. D. (2011). Obstructive Sleep Apnea and Depression: A Review. *Innovations in Clinical Neuroscience*, 8(8), 17–25.
- Elavsky, S., McAuley, E., Motl, R. W., Konopack, J. F., Marquez, D. X., Hu, L., ... & Diener, E. (2005). Physical activity enhances long-term quality of life in older adults: efficacy, esteem, and affective influences. *Annals of Behavioral Medicine*, 30(2), 138-145.

- Epstein, L. J., Kristo, D., Strollo, P. J., Friedman, N., Malhotra, A., Patil, S. P., ... & Weinstein, M. D. (2009). Clinical guideline for the evaluation, management and long-term care of obstructive sleep apnea in adults. *Journal of clinical sleep medicine*, 5(03), 263-276.
- Emery, C. F., Schein, R. L., Hauck, E. R., & MacIntyre, N. R. (1998). Psychological and cognitive outcomes of a randomized trial of exercise among patients with chronic obstructive pulmonary disease. *Health Psychology*, 17(3), 232.
- Fox, L. D., Rejeski, W. J., & Gauvin, L. (2000). Effects of leadership style and group dynamics on enjoyment of physical activity. *American Journal of Health Promotion*, 14(5), 277-283.
- Gay, P., Weaver, T., Loube, D., & Iber, C. (2006). Evaluation of positive airway pressure treatment for sleep related breathing disorders in adults. *Sleep*, 29(3), 381-401.
- Gillison, F. B., Standage, M., & Skevington, S. M. (2006). Relationships among adolescents' weight perceptions, exercise goals, exercise motivation, quality of life and leisure-time exercise behaviour: a self-determination theory approach. *Health education research*, 21(6), 836-847.
- Hong S, Dimsdale JE. Physical activity and perception of energy and fatigue in obstructive sleep apnea. *Med Sci Sports Exerc*. 2003;35(7):1088–1092.
- Huo, T., Guo, Y., Shenkman, E., & Muller, K. (2018). Assessing the reliability of the short form 12 (SF-12) health survey in adults with mental health conditions: a report from the wellness incentive and navigation (WIN) study. *Health and quality of life outcomes*, 16(1), 34. doi:10.1186/s12955-018-0858-2
- Iftikhar, I. H., Kline, C. E., & Youngstedt, S. D. (2014). Effects of exercise training on sleep apnea: a meta-analysis. *Lung*, 192(1), 175-184.

Jean, R. E., Duttuluri, M., Gibson, C. D., Mir, S., Fuhrmann, K., Eden, E., & Supariwala, A. (2017). Improvement in physical activity in persons with obstructive sleep apnea treated with continuous positive airway pressure. *Journal of Physical Activity and Health, 14*(3), 176-182.

Jette AM, Lachman M, Giorgetti MM, et al. 1999. Exercise—it's never too late: the Strong-for-Life program. *Am J Public Health.*89:66-72.

Johns MW. A new method for measuring daytime sleepiness: the Epworth Sleepiness Scale. *Sleep* 1991;14:540–545

Kline, C. E., Crowley, E. P., Ewing, G. B., Burch, J. B., Blair, S. N., Durstine, J. L., ... & Youngstedt, S. D. (2011). The effect of exercise training on obstructive sleep apnea and sleep quality: a randomized controlled trial. *Sleep, 34*(12), 1631-1640.

Kline, C. E., Ewing, G. B., Burch, J. B., Blair, S. N., Durstine, J. L., Davis, J. M., & Youngstedt, S. D. (2012). Exercise training improves selected aspects of daytime functioning in adults with obstructive sleep apnea. *Journal of clinical sleep medicine : JCSM : official publication of the American Academy of Sleep Medicine, 8*(4), 357-65. doi:10.5664/jcsm.2022

Lacasse, Y., Godbout, C., & Series, F. (2002). Health-related quality of life in obstructive sleep apnoea. *European Respiratory Journal, 19*(3), 499-503.

LaGrotte, C., Fernandez-Mendoza, J., Calhoun, S. L., Liao, D., Bixler, E. O., & Vgontzas, A. N. (2016). The relative association of obstructive sleep apnea, obesity and excessive daytime sleepiness with incident depression: a longitudinal, population-based study. *International Journal of Obesity, 40*(9), 1397.

Luntamo T, Sourander A, Rihko M, et al. Psychosocial determinants of headache, abdominal pain, and sleep problems in a community sample of Finnish adolescents. *Eur Child Adolesc Psychiatry.* 2012;21(6):301–313

Mannarino, M. R., Di Filippo, F., & Pirro, M. (2012). Obstructive sleep apnea syndrome. *European journal of internal medicine, 23*(7), 586-593.

Medic, G., Wille, M., & Hemels, M. E. (2017). Short- and long-term health consequences of sleep disruption. *Nature and Science of Sleep*, *9*, 151–161.

<http://doi.org/10.2147/NSS.S134864>

Mendelson, M., Bailly, S., Marillier, M., Flore, P., Borel, J. C., Vivodtzev, I., Doutreleau, S., Verges, S., Tamisier, R., ... Pépin, J. L. (2018). Obstructive Sleep Apnea Syndrome, Objectively Measured Physical Activity and Exercise Training Interventions: A Systematic Review and Meta-Analysis. *Frontiers in neurology*, *9*, 73. doi:10.3389/fneur.2018.00073

Moore, P., Bardwell, W. A., Ancoli-Israel, S., & Dimsdale, J. E. (2001). Association between polysomnographic sleep measures and health-related quality of life in obstructive sleep apnea. *Journal of sleep research*, *10*(4), 303-308.

Motamedi, K. K., McClary, A. C., & Amedee, R. G. (2009). Obstructive Sleep Apnea: A Growing Problem. *The Ochsner Journal*, *9*(3), 149–153.

Naegele B, Thouvard V, Pepin JL. Deficits of cognitive executive functions in patients with sleep apnea syndrome. *Sleep* 1995; *18*: 43–52.

Ng, J. Y., Ntoumanis, N., Thøgersen-Ntoumani, C., Deci, E. L., Ryan, R. M., Duda, J. L., & Williams, G. C. (2012). Self-determination theory applied to health contexts: A meta-analysis. *Perspectives on Psychological Science*, *7*(4), 325-340.

Paquet, Y., & Vallerand, R. (2016). La théorie de l'autodétermination: Aspects théoriques et appliquées. De Boeck supérieur.

Pépin, J. L., Leger, P., Veale, D., Langevin, B., Robert, D., & Lévy, P. (1995). Side effects of nasal continuous positive airway pressure in sleep apnea syndrome: study of 193 patients in two French sleep centers. *Chest*, *107*(2), 375-381.

Peppers LG, 1976. Patterns of leisure and adjustment to retirement. *Gerontologist*. *16*:441-446.

Perri, M. G., Anton, S. D., Durning, P. E., Ketterson, T. U., Sydeman, S. J., Berlant, N. E., ... & Martin, A. D. (2002). Adherence to exercise prescriptions: effects of prescribing moderate versus higher levels of intensity and frequency. *Health Psychology, 21*(5), 452.

Puetz, T. W. (2006). Physical activity and feelings of energy and fatigue. *Sports medicine, 36*(9), 767-780.

Puetz, T. W., Flowers, S. S., & O'Connor, P. J. (2008). A randomized controlled trial of the effect of aerobic exercise training on feelings of energy and fatigue in sedentary young adults with persistent fatigue. *Psychotherapy and psychosomatics, 77*(3), 167-174.

Punjabi, N. M. (2008). The Epidemiology of Adult Obstructive Sleep Apnea. *Proceedings of the American Thoracic Society, 5*(2), 136–143. <http://doi.org/10.1513/pats.200709-155MG>

Reid, K. J., Baron, K. G., Lu, B., Naylor, E., Wolfe, L., & Zee, P. C. (2010). Aerobic exercise improves self-reported sleep and quality of life in older adults with insomnia. *Sleep medicine, 11*(9), 934-940.

Rejeski, W. J., & Mihalko, S. L. (2001). Physical activity and quality of life in older adults. *The Journals of Gerontology Series A: Biological sciences and medical sciences, 56*(suppl_2), 23-35.

Ryan, R. M., Patrick, H., Deci, E. L., & Williams, G. C. (2008). Facilitating health behaviour change and its maintenance: Interventions based on self-determination theory. *European Health Psychologist, 10*(1), 2-5.

Schipper H. Quality of life: principles of the clinical paradigm. *J Psychosocial Oncology. 1990;8:171–185*

Schwartz, A. R., Patil, S. P., Laffan, A. M., Polotsky, V., Schneider, H., & Smith, P. L. (2008). Obesity and obstructive sleep apnea: pathogenic mechanisms and therapeutic approaches. *Proceedings of the American Thoracic Society, 5*(2), 185–192.

doi:10.1513/pats.200708-137MG

Sebire, S. J., Kesten, J. M., Edwards, M. J., May, T., Banfield, K., Tomkinson, K., ... & Jago, R. (2016). Using self-determination theory to promote adolescent girls' physical activity: Exploring the theoretical fidelity of the Bristol Girls Dance Project. *Psychology of sport and exercise, 24*, 100-110.

Singh, N. A., Clements, K. M., & Fiatarone, M. A. (1997). A randomized controlled trial of progressive resistance training in depressed elders. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences, 52*(1), M27-M35.

Sosnowski, R., Kulpa, M., Ziętalewicz, U., Wolski, J. K., Nowakowski, R., Bakuła, R., & Demkow, T. (2017). Basic issues concerning health-related quality of life. *Central European journal of urology, 70*(2), 206–211. doi:10.5173/ceju.2017.923

Senaratna, C. V., Perret, J. L., Lodge, C. J., Lowe, A. J., Campbell, B. E., Matheson, M. C., ... & Dharmage, S. C. (2017). Prevalence of obstructive sleep apnea in the general population: a systematic review. *Sleep medicine reviews, 34*, 70-81.

Sengul, Y. S., Ozalevli, S., Oztura, I., Itil, O., & Baklan, B. (2011). The effect of exercise on obstructive sleep apnea: a randomized and controlled trial. *Sleep and Breathing, 15*(1), 49-56.

Shah, K. N., Majeed, Z., Yoruk, Y. B., Yang, H., Hilton, T. N., McMahon, J. M., ... & Ryan, R. M. (2016). Enhancing physical function in HIV-infected older adults: A randomized controlled clinical trial. *Health Psychology, 35*(6), 563.

Sharafkhaneh, A., Giray, N., Richardson, P., Young, T., & Hirshkowitz, M. (2005). Association of psychiatric disorders and sleep apnea in a large cohort. *Sleep, 28*(11), 1405-1411.

Shoib, S., Malik, J. A., & Masoodi, S. (2017). Depression as a Manifestation of Obstructive Sleep Apnea. *Journal of Neurosciences in Rural Practice, 8*(3), 346–351.

http://doi.org/10.4103/jnrp.jnrp_462_16

Silva, M. N., Markland, D., Carraça, E. V., Vieira, P. N., Coutinho, S. R., Minderico, C. S., ... & Teixeira, P. J. (2011). Exercise autonomous motivation predicts 3-yr weight loss in women. *Medicine & Science in Sports & Exercise*, 43(4), 728-737.

Somers, V. K., White, D. P., Amin, R., Abraham, W. T., Costa, F., Culebras, A., ... & Pickering, T. G. (2008). Sleep apnea and cardiovascular disease: An american heart association/american college of cardiology foundation scientific statement from the american heart association council for high blood pressure research professional education committee, council on clinical cardiology, stroke council, and council on cardiovascular nursing in collaboration with the national heart, lung, and blood institute national center on sleep disorders research (national institutes of health). *Journal of the American College of Cardiology*, 52(8), 686-717.

Spicuzza, L., Caruso, D., & Di Maria, G. (2015). Obstructive sleep apnoea syndrome and its management. *Therapeutic Advances in Chronic Disease*, 6(5), 273–285.

<http://doi.org/10.1177/2040622315590318>

Standage, M., Gillison, F. B., Ntoumanis, N., & Treasure, D. C. (2012). Predicting students' physical activity and health-related well-being: A prospective cross-domain investigation of motivation across school physical education and exercise settings. *Journal of Sport and Exercise Psychology*, 34(1), 37-60.

Teixeira, P. J., Carraça, E. V., Markland, D., Silva, M. N., & Ryan, R. M. (2012). Exercise, physical activity, and self-determination theory: a systematic review. *International journal of behavioral nutrition and physical activity*, 9(1), 78.

Teychenne, M., Ball, K., & Salmon, J. (2008). Physical activity and likelihood of depression in adults: a review. *Preventive medicine*, 46(5), 397-411.

Tiemeier H, Pelzer E, Jonck L, Moller HJ, Rao ML. Plasma catechol-amines and selective slow wave sleep deprivation. *Neuropsychobiology*. 2002;45(2):81–86.

Tyni-Lenné, R., Gordon, A., Jansson, E., & Sylven, C. (1998). Exercise-based rehabilitation improves skeletal muscle capacity, exercise tolerance, and quality of life in both women and men with chronic heart failure. *Journal of cardiac failure*, 4(1), 9-17.

Ware J Jr, Kosinski M, Keller SD. A 12-item short-form health survey: construction of scales and preliminary tests of reliability and validity. *Med Care* 1996;34:220–233

Wittels, E. H., & Thompson, S. (1990). Obstructive sleep apnea and obesity. *Otolaryngologic Clinics of North America*, 23(4), 751-760.